## Sampling Scheme & Question Design

By: Emily Butler, Meggie Hayes, Douglas Heckmann, Christopher Peter Makris, & Thomas Todd

## K. Sampling Scheme

Since we will be conducting a "man on the street" survey, it will be difficult to collect a please do a completely random sample. Therefore, we plan to collect a pseudo-random sample by attempting little "dry run" to sample every third person that passes by. We will also be implementing stratification to try to somewhere to gain a demographically accurate sample of our target population. This stratification will be done get an idea of what's going by soliciting survey responses in different locations where we expect certain demographic groups to be to be commonly located. By using this method of stratification, we hope that the demographics workable in of our sample population will be representative of our target population.

what variable(s) will you stratify on?

## L. Proposed Questions

1

Before individuals take our survey, we will ask them whether or not they are an undergraduate student. If they are an undergraduate student, we will ask them to fill out our survey. We believe that by doing so our coverage error will be reduced since we completely eliminate receiving responses from non-undergraduate students who are all outside the scope of our target population. Therefore, we consider the question "Are you an undergraduate student?" an initial question that is not technically part of our survey.

- 0. (Verbally) Are you an undergraduate student?
  - -If response is "yes," offer the individual a survey to complete.
- 1. Do you consume alcoholic beverages?
  - -If response is "no," please stop this survey and return it to your surveyor.
- 2. What university do you attend? Please choose one:
  - -Carnegie Mellon University
  - -The University of Pittsburgh
  - -Other
- 3. Are you a full-time or part-time student?
- 4. What is your age?
- 5. What is your gender?
- 6. Are you affiliated with the Greek life of your University?
- 7. How frequently do you consume at least one alcoholic beverage? Please choose one:
  - -Less than 1 time per week, on average
  - -From 1 to 2 times per week, on average
  - -From 3 to 5 times per week, on average
  - -Greater than 5 times per week, on average
- 8. Do you consume energy drinks in conjunction with alcohol?
- 9. Have you ever experienced complete or partial memory loss (i.e. a "blackout") during an event that involved drinking?
  - -If response is "no," please skip to question #11.
- 10. Were you consuming an alcoholic energy drink or alcohol in conjunction with an energy drink in the events shortly prior to any blackouts?

not clear what you're using to stratify on or monitor coverage and unbiasedness here. person might

be too frequent to

- 11. Have you ever directly or indirectly purchased alcoholic energy drinks?
  - -If response is "no," please skip to question #13.
- 12. Why did you choose to purchase alcoholic energy drinks? Please choose all that apply. -Price

-Taste

-Alcoholic content

-Availability

-Effects of consumption

-Other reason(s)

these are related. you might consider instead a check-all-that-apply "what effect attracted you" question, along with a blanik for any effects you didn't think of.

- 13. Have you ever heard of the alcoholic energy product Four Loko?
  - -If response is "no," please skip to question #18.
- 14. How did you first hear about Four Loko? Please choose one:
  - -A friend or relative
  - -A print advertisement
  - -News media
  - -An online source
  - -Other outlet (give blank)
- 15. Have you ever consumed the product Four Loko?
  - -If response is "no," please skip to question #17.
- 16. The caffeine and energy stimulants were removed from Four Loko by January 1, 2011. When did you consume Four Loko?
  - -Before January 1, 2011
  - -After January 1, 2011
  - -Both before & after January 1, 2011
- 17. Would you consider drinking the product Four Loko since the removal of its caffeine and energy stimulants?
- 18. In general, would you consider drinking alcoholic energy drinks in the future?
- 19. Have the recent health risks associated with alcoholic energy drinks influenced your
- answer to the previous question?

some "why" or "what risks" followups might be interesting

## M. Initial<sup>es</sup> ample Size Calculation

Because our survey consists of primarily "yes" or "no" answers, we consider most of our parameters to be from a Bernoulli distribution. Therefore, we will consider our standard deviation to be of the worst case scenario, by setting p = .5:

$$SD_{Worst \ Case \ Scenario} = \sqrt{(.5)(1-.5)} = .5$$

It may be difficult to receive a large sample size since there are many groups conducting surveys within our class. Therefore, we will allow our margin of error to be up to 5%. We can calculate  $n_0$  as follows:

$$n_0 = \frac{\left(\frac{Z\alpha}{2}\right)^2 (SD)^2}{(ME)^2}$$



Overall, the total number of undergraduate students attending both Carnegie Mellon University and the University of Pittsburgh is approximately 25,000. Because we are under the assumption that we are taking a random sample without replacement, we must make the following adjustment to our calculation above:

 $n \ge \frac{Nn_0}{N+n_0}$  $n \ge \frac{(25,000)(270.6025)}{(25,000) + (270.6025)}$ 

 $n \geq 267.7048361$ 

Therefore, to make inferences about our population concerning the questions we plan to ask with a margin of error of 5%, we must sample at least 268 total individuals.

Since we are conducting a "face to face" survey, with practice we may be able to get a response rate up to 70% (as noted in the lecture slides). Therefore, for the worst case scenario we will consider our response rate to be approximately 50%. If we take our response rate into account, we may need to physically ask more individuals for their participation in our survey. Specifically:

$$n_{With, 5 Response Rate} \geq \frac{267.7048361}{.5}$$
 this is great  
$$n_{With, 5 Response Rate} \geq 535.4096722$$

Thus, for a worst case scenario calculation, we need to ask approximately 536 individuals to take our survey, but only have approximately 268 of those individuals completely fill out our survey, to be able to make inferences with a 5% margin of error.