

36-303: Sampling, Surveys & Society

Post-survey Processing
Brian Junker
132E Baker Hall
brian@stat.cmu.edu

31 March 2011

1

Handouts

- These Lecture Notes
- Handout on post-stratification challenges
- Remember: HW 5 is due Apr 7

31 March 2011

2


Outline

- Post-survey Processing [Groves Ch 10]
 - Coding
 - Weighting
 - Imputation
 - Variance Estimation
- Review

31 March 2011

3

Post-survey Processing



- Top row: Raw data collection process
 - The order of Coding, Data Entry and Editing will depend on the data collection design (FTF, phone, www, computer assisted, ...)
 - Computer-based surveys require you to design the Data Entry and Edit Checks when you build the form in surveymonkey.com, questionpro.com, etc.
- Bottom row: Calculations based on the data and/or design

31 March 2011

4

Coding

- Translating non-quantitative or non-categorical data into quantities and categories
- M/C and Likert items usually require no coding
 - Indicate your status (check one box only):
 - Full-time student
 - Part-time student
 - Applicant, acceptance letter received
 - Applicant, acceptance letter not received
- When might an M/C item require some coding anyway?

31 March 2011

5

Coding

- Short answer, long answer, graphical response, performance, etc., all require some coding
 - Team A: Students Changing Majors: "Why did you choose/change your major?"
 - Team I: Kid's Architecture Knowledge: "Draw a plan of your school"
- On the job crime in the NCVS:
 - What is the name of the (company/government agency/business/nonprofit organization) for which you worked at the time of the accident?
 - What kind of business or industry is this? (*What do they make or do where you worked at the time of the incident?*)
 - What kind of work did you do; that is, what was your occupation at the time of the incident? (*For example: plumber, typist, farmer*)
 - What were your usual activities or duties at this job?

31 March 2011

6

Coding System (Code Structure, Rubric)

- **Each code should include:**
 - A number or category, used in statistical analysis
 - A text label, describing all answers in that category
- **The set of codes for a response should be:**
 - Exhaustive: Every response should be codable into one of the categories
 - Separate codes are needed for skipped, not-asked, off-topic, etc.
 - Exclusive: No response should be codable into more than one category
 - Appropriate to the purposes of the research
 - Use codes that help you answer your research question(s)
 - If you have more than one research question, you might code the same response using different coding systems
- **Different coders using the same system will produce different codes for the same response**
 - Introduces a kind of "cluster structure" (by coder) into the data
 - Want to construct codes to minimize this (reduce ICC!)

31 March 2011

7

Standard Classification Systems

- If the survey will be compared with other surveys, they should use the same coding scheme
- An ongoing longitudinal or panel survey like NCVS tries to use the same categories in each survey cycle or wave.
- Government or international agencies maintain standard coding systems for common types of information
 - Race/Ethnicity (US Census)
 - Standard Occupational Classification (US Dept of Labor)
 - North American Industry Classification System (US Economic Classification Policy Committee)

31 March 2011

8

Weighting

- Many different sources of weighting in a survey, e.g.
 - Survey design weights
 - In stratified sampling, weights are used to combine stratum means & variance into overall means & variances
 - Other survey designs require weights to account for unequal sampling probabilities, etc.
 - Compute these weights *before the data is collected*
 - Nonresponse and post-stratification weights
 - Try to adjust sample proportions to equal population proportions
 - Compute these weights *after you see the data*
- These are discussed in detail in Groves; below I will only talk about post-stratification weights

31 March 2011

9

Post-Stratification Weights

- As part of survey data collection it is a good idea to get general demographic information (e.g. in our surveys: sex, age, class, major, hometown, etc.)
- After data collection we compare the proportions in each of these categories in our sample with the same proportions in the population
- If they agree, great. If they disagree, we may re-weight the sample to make them agree
- These categories are called “post-strata”, and the weights are called “post-stratification weights”

31 March 2011

10

Post-Stratification Example

- The 2007 HSS advising satisfaction survey was a simple (no strata, no clusters) web survey of all 986 students in HSS.
- We can separate the responding students by major to see how representative the survey was of each department in HSS.
- If the representativeness was not the same in each department, and if we assume that the nonresponse is ignorable (??) within department, we can re-weight the sample data to get more accurate estimates of population quantities.

31 March 2011

11

HSS Response Rate in Dept Post-Strata

| Post-Stratum | Sample | Population | Resp Rate |
|--------------|--------|------------|-----------|
| Economics | 40 | 126 | 0.32 |
| English | 39 | 115 | 0.34 |
| History | 21 | 48 | 0.44 |
| ModLang | 8 | 16 | 0.50 |
| Philosophy | 4 | 7 | 0.57 |
| Psychology | 37 | 104 | 0.36 |
| SDS | 54 | 161 | 0.34 |
| Statistics | 6 | 8 | 0.75 |
| Interdisc/IS | 76 | 233 | 0.33 |
| Undeclared | 19 | 168 | 0.11 |
| Total | 304 | 986 | 0.31 |

31 March 2011

12

HSS Post-Strata Proportions & Weights

| Post-Stratum | Sample | Prop | Population | Prop | Weights |
|--------------|--------|-------|------------|-------|---------|
| Economics | 40 | 0.132 | 126 | 0.128 | 0.97 |
| English | 39 | 0.128 | 115 | 0.117 | 0.91 |
| History | 21 | 0.069 | 48 | 0.049 | 0.70 |
| ModLang | 8 | 0.026 | 16 | 0.016 | 0.62 |
| Philosophy | 4 | 0.013 | 7 | 0.007 | 0.54 |
| Psychology | 37 | 0.122 | 104 | 0.105 | 0.87 |
| SDS | 54 | 0.178 | 161 | 0.163 | 0.92 |
| Statistics | 6 | 0.020 | 8 | 0.008 | 0.41 |
| Interdisc/IS | 76 | 0.250 | 233 | 0.236 | 0.95 |
| Undeclared | 19 | 0.062 | 168 | 0.170 | 2.73 |
| Total | 304 | | 986 | | |

weight = (Population Proportion) / (Sample Proportion)

31 March 2011

13

Fictional Example: What proportion of students think advising is OK?

| Post-stratum | Total | | Think Advising is OK | |
|--------------|--------|------------|----------------------|------------|
| | Sample | Population | Sample | Population |
| Economics | 40 | 126 | 28 | 88 |
| English | 39 | 115 | 23 | 69 |
| History | 21 | 48 | 10 | 24 |
| ModLang | 8 | 16 | 3 | 6 |
| Philosophy | 4 | 7 | 1 | 2 |
| Psychology | 37 | 104 | 11 | 31 |
| SDS | 54 | 161 | 22 | 64 |
| Statistics | 6 | 8 | 3 | 4 |
| Interdisc/IS | 76 | 233 | 46 | 140 |
| Undeclared | 19 | 168 | 13 | 118 |
| Total | 304 | 986 | 160 | 546 |

31 March 2011

14

Population proportion, vs. Unweighted and Weighted sample proportion

- Population proportion:

$$p_{pop} = 546/986 = 0.553$$

- Unweighted Sample proportion:

$$\hat{p} = 160/304 = 0.526$$

- Weighted Sample Proportion

$$\text{Weighted Total} = (0.97)(40) + (0.91)(39) + \dots + (2.73)(19) = 304(!)$$

$$\text{Weighted OK's} = (0.97)(28) + (0.91)(23) + \dots + (2.73)(13) = 167.45$$

$$\hat{p} = 167.45/304 = 0.551$$

Post-Stratification Weights – Pros & Cons

- Post-stratification weights can fix
 - disproportionate sampling of post strata
 - disproportionate nonresponse across poststrata
- Only works if the sampling/nonresponse process is **ignorable** within post-strata
 - That is, nonresponse does not depend on the answer you would have gotten if the person had responded
- If the sampling/nonresponse process is non-ignorable then these weights don't work; other weights have to be used**
- The weights are only as good as your model for nonresponse
 - These weights are a very big deal in pre-election phone surveys for example (resp. rate 5-20%; weights account for ignorable and nonignorable nonresponse)

31 March 2011

16

31 March 2011

15

Imputation

- **Weights** are a good solution for unit nonresponse (missed that whole person)
- **Imputation** is a good solution for item nonresponse (person never answered question #17).
- Basic ideas of imputation:
 - Build a model for **what sort of person wouldn't respond**, and use the model to fill in a value for this person
 - Find one or more other people like this person who **did** answer #17, and use their answers for this person
- Alternative to imputation: **Case-wise deletion**
 - Delete this person from the survey so you don't have to deal with the nonresponse to question #17
 - Pro's and con's of case-wise deletion??
 - MCAR: Missing Completely at Random

31 March 2011

17

Mean-value Imputation

- If question #17 is a numerical item, take the average of everyone else's answer to #17, and fill that in for this person
- If question #17 is a yes/no, fill in the proportion of yes's for everyone else (or do a flip of a coin with that probability of "heads")
- Pro's and con's?
- MCAR

31 March 2011

18

Hot-Deck Imputation

- Among all the other people who answered question #17, find the one person who matches this person on important variables
 - age, sex, occupation, answers to other questions, etc. (whatever you think is important to understand this non-response!)
- Fill in that person's answer for this person's #17.
- Pro's? Cons?
- MAR: Missing at Random (within covariates)

31 March 2011

19

Regression Imputation

- Among all the people who answered question #17, fit a regression model (or logistic regression, or whatever) for response to question #17 as a function of other variables:
$$y_{17} = \beta_0 + \beta_1(\text{age}) + \beta_2(\text{sex}) + \beta_3(\text{occupation}) + \beta_4(\text{answer to Q3}) + \dots + \epsilon$$
- Use the fitted model to predict what this person would have answered to #17, and fill that value in
- Pro's? Con's?
- MAR

31 March 2011

20

Limitations of Imputation

- You have to have other variables in the survey that help you build a model for the nonresponse (MAR, or better MCAR)
- Nonignorable missingness (MNAR, missing not at random) is much harder
- After you have filled in the missing data
 - You have NOT increased the sample size; this will matter a lot if you are doing a lot of imputation
 - There is some uncertainty in what value to fill in; this can be accounted for by a technique called **“multiple imputation”**

Variance Calculations

- Final weights in a survey will be a combination (usually multiply together) of
 - Design Weights
 - Nonresponse Weights
 - Post-stratification Weights
- They each improve the point estimate (\bar{y}) but they have different effects on the variance ($Var(\bar{y})$) and standard error
- Additional variance adjustments are made for imputation
- There are not usually closed-form variance formulas (like our simple formulas for strata and clusters)
- More on this next time!

Review

- The entire survey process consists of
 - Collect/Code/Enter/Edit the data
 - Post-processing of the data
- **Coding** is the process of summarizing complex responses into numbers or categories – subject to bias (bad categories) and variability (coder variation)
- **Post-processing** includes (usually in this order)
 - Imputation
 - Weighting
 - Variance Calculation

These are calculations on the data to account for various weaknesses of the data.