
36-303: Sampling, Surveys and Society

Stratified Samples and Sample Size Calculations

Brian Junker

132E Baker Hall

brian@stat.cmu.edu

Handouts

- These Lecture Notes
- Handout on Stratified Sampling
- Handout on Sampling Details
 - Selecting an SRS from C-Book
 - Contacting respondents
 - Nonresponse followup on surveymonkey.com
- Reading:
 - Stratified Sampling: Groves Sect 4.5,
 - Nonresponse: Groves Ch 6

Outline

- Team Projects This Week
- Midterm
- Stratification
 - What is it; Notation
 - Weights and Proportionate Sampling
 - Variances and Design Effect
 - Examples

Team Projects This Week

■ II.5 Due Thursday (Blackboard)

- Include a paragraph or so on your research question
- Pretest a version of your questionnaire (or observation protocol) on a group of possible respondents/units.
- Write 2-4 paragraphs: how many respondents/units you used in the pretest; how similar they were to units in the population; and the changes you made in the survey based on this pretest.
- Include both old and revised questionnaire/protocol

■ IRB Form Due Thursday (Blackboard)

- If you are surveying people, and you have not turned one in to me yet, you need to do this by Thursday.
- Form at <http://www.stat.cmu.edu/~brian/303>. You don't need to include any attachments

■ Peer Evaluations Due Thursday (email to me)

- Each person should email me forms for all other members of their team, in one email with subject "36303 Peer Evaluations"
- Form at <http://www.stat.cmu.edu/~brian/303>

Team Projects After Spring Break

- II.6 Project Plan (Tue Mar 20, Blackboard)
 - Final, full project proposal (items A-M on the “designing a sample survey” handout, except don’t include the IRB form [item I]).
 - This should be easy: copy and update the latest thing you have done for each of the items A-M up to now into a single electronic file to submit on blackboard. for each team.
 - From this proposal, anyone outside our class should be able to read and understand completely what you are proposing to do
- **Get started Collecting Data!**

Midterm Exam ...

■ Seemed to go well...

```
> summary(exam1)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	NA's
64.00	76.00	85.00	82.41	89.00	99.00	3.00

```
> stem(exam1)
```

```
6 | 4
-----
6 | 8999          C (11)
7 | 2
7 | 567799
-----
8 | 04            B (11)
8 | 566778899
-----
9 | 224          A (6)
9 | 559
```

Stratified Sampling

- Strata are just subgroups of the target population that have some feature in common (gender, major, region, income, ...)
- Why stratify?
 - We need to make a separate inference for each stratum (e.g. we want to estimate men's and women's incomes separately)
 - Different sampling schemes would be used in each stratum (PA voters in PA, vs PA voters in Afghanistan)
 - Population is geographically diverse (Minnesota, Illinois, Ohio, Pennsylvania)
 - Reduce variance of estimates (and reduce sample size) by exploiting similarity among members of the same stratum

What is Stratification?

Record	Name	Group	
1	Bradburn, N.	High	
2	Cochran, W.	Highest	
3	Deming, W.	High	
4	Fuller, W.	Medium	
5	Habermann, H.	Medium	
6	Hansen, M.	Low	
7	Hunt, J.	Highest	
8	Hyde, H.	High	
9	Kalton, G.	Medium	→ Kalton, G.
10	Kish, L.	Low	
11	Madow, W.	Highest	
12	Mandela, N.	Highest	
13	Norwood, J.	Medium	→ Norwood, J.
14	Rubin, D.	Low	→ Rubin, D.
15	Sheatsley, P.	Low	
16	Steinberg, J.	Low	
17	Sudman, S.	High	
18	Wallman, K.	High	→ Wallman, K.
19	Wolfe, T.	Highest	
20	Woolsley, T.	Medium	

One SRS of Size 4

Unstratified Sample

Record	Name	Group	
2	Cochran, W.	Highest	
7	Hunt, J.	Highest	
11	Madow, W.	Highest	
12	Mandela, N.	Highest	
19	Wolfe, T.	Highest	→ Wolfe, T.
1	Bradburn, N.	High	→ Bradburn, N.
3	Deming, W.	High	
8	Hyde, H.	High	
17	Sudman, S.	High	
18	Wallman, K.	High	
4	Fuller, W.	Medium	→ Fuller, W.
5	Habermann, H.	Medium	
9	Kalton, G.	Medium	
13	Norwood, J.	Medium	
20	Woolsley, T.	Medium	
6	Hansen, M.	Low	
10	Kish, L.	Low	
14	Rubin, D.	Low	→ Rubin, D.
15	Sheatsley, P.	Low	
16	Steinberg, J.	Low	

One Stratified Random
Sample of Total Size 4

Stratified Sample

Some Basic Notation

■ H strata

- N_h = population size in each stratum $N = \sum_{h=1}^H N_h$
- n_h = sample size in each stratum $n = \sum_{h=1}^H n_h$
- $f_h = n_h/N_h$ = sampling fraction, each stratum

■ The population average

$$\bar{y}_{pop} = \frac{1}{N} \sum_{i=1}^N y_i = \frac{1}{N} \sum_{h=1}^H \sum_{i=1}^{N_h} y_{hi} = \sum_{h=1}^H \frac{N_h}{N} \frac{1}{N_h} \sum_{i=1}^{N_h} y_{hi} = \sum_{h=1}^H \frac{N_h}{N} \bar{y}_{h,pop}$$

■ In stratified sampling we mimic this

$$\bar{y}_{st} = \frac{1}{n} \sum_{i=1}^n y_i = \sum_{h=1}^H \frac{N_h}{N} \bar{y}_h \text{ where } \bar{y}_h = \frac{1}{n_h} \sum_{i=1}^{n_h} y_{hi}$$

Weights, and Proportionate Sampling

- Let $W_h = N_h/N$. Then

$$\bar{y}_{pop} = \sum_{h=1}^H W_h \bar{y}_{h,pop} \text{ and } \bar{y}_{st} = \sum_{h=1}^H W_h \bar{y}_h$$

- In proportionate sampling we let $f_h = n_h/N_h = f$ for all strata h . Then $n_h/n = N_h/N$ (why??)
 - The sample is called “self-weighting”
 - Sample mean is “simple” for self-weighting

$$\begin{aligned} \bar{y}_{st} &= \sum_{h=1}^H W_h \bar{y}_h = \sum_{h=1}^H \frac{N_h}{N} \bar{y}_h = \sum_{h=1}^H \frac{n_h}{n} \bar{y}_h = \\ &= \sum_{h=1}^H \frac{n_h}{n} \frac{1}{n_h} \sum_{i=1}^{n_h} y_{hi} = \frac{1}{n} \sum_{h=1}^H \sum_{i=1}^{n_h} y_{hi} = \frac{1}{n} \sum_{i=1}^n y_i \bar{y}_{sts} \end{aligned}$$

Sampling Variances

(SRS w/o replacement in each stratum)

- Within each stratum it's the same old answer

$$Var(\bar{y}_h) = (1 - f_h) \frac{s_h^2}{n_h} \text{ where } s_h^2 = \frac{1}{n_h - 1} \sum_{i=1}^{n_h} (y_{hi} - \bar{y}_h)^2$$

- Then we combine across strata using weights

$$\begin{aligned} (W_h)^2: \quad Var(\bar{y}_{st}) &= Var\left(\sum_{h=1}^H W_h \bar{y}_h\right) \\ &= \sum_{h=1}^H Var(W_h \bar{y}_h) = \sum_{h=1}^H W_h^2 Var(\bar{y}_h) \\ &= \sum_{h=1}^H W_h^2 (1 - f_h) \frac{s_h^2}{n_h} \end{aligned}$$

Design Effect

- The design effect is a measure of how much better or worse Stratified is than one SRS:

$$d^2 = \frac{Var(\bar{y}_{st})}{Var(\bar{y}_{srs})} = \frac{\sum_{h=1}^H W_h^2 (1 - f_h) \frac{s_h^2}{n_h}}{(1 - f) \frac{s^2}{n}}$$

- Usually, $d^2 < 1$, i.e. stratified does better than one big SRS!
 - Usually best if:
 - Elements are more similar to each other within strata than between (e.g., substantively meaningful strata)
 - Proportionate sampling
 - Cochran (1961) suggests 2-6 strata usually give the best results; greater than 6 OK, but there are diminishing returns

Handout on Stratified Sampling

(Briefly) Handout on Sampling Details

Review

- Team Projects This Week
- Midterm Exam
- Stratification
 - What is it; Notation
 - Weights and Proportionate Sampling
 - Variances and Design Effect
 - Handout on Stratified Sampling
- Handout on Sampling Details
- Reading:
 - Stratified Sampling: Groves Sect 4.5,
 - Nonresponse: Groves Ch 6