
36-303: Sampling, Surveys & Society

Post-survey Processing
Brian Junker
132E Baker Hall
brian@stat.cmu.edu

Handouts


- These Lecture Notes
 - Handout on post-stratification challenges
 - [reading for Thursday]

 - Reminders:
 - HW 5 is due Thu Apr 5
 - But I will accept Fri Apr 6 also due to confusion about due date!
 - There is a place to submit it on Blackboard
 - First Drafty Draft is due Fri Apr 6
 - Submit one pdf or doc file per group, on Blackboard
-

Outline

- Post-survey Processing [Groves Ch 10]
 - Coding
 - Weighting
 - Imputation
 - Variance Estimation
- Review/Summary

Post-survey Processing



- Top row: Raw data collection process
 - The order of Coding, Data Entry and Editing will depend on the data collection design (FTF, phone, www, computer assisted, ...)
 - Computer-based surveys require you to design the Data Entry and Edit Checks when you build the form in surveymonkey.com, questionpro.com, Google Docs, etc.
- Bottom row: Calculations based on the data and/or design

Coding

- Translating non-quantitative or non-categorical data into quantities and categories
- M/C and Likert items usually require no coding
 - Indicate your status (check one box only):
 - Full-time student
 - Part-time student
 - Applicant, acceptance letter received
 - Applicant, acceptance letter not received
- When might an M/C item require some coding anyway?

Coding

- Short answer, long answer, graphical response, performance, etc., all require some coding
 - Team F: Coding sheets for make & type of car, etc.
 - Team A: What is your Religious Affiliation? (open answer)


- On the job crime in the NCVS:
 - What is the name of the (company/government agency/business/nonprofit organization) for which you worked at the time of the accident?
 - What kind of business or industry is this? (*What do they make or do where you worked at the time of the incident?*)
 - What kind of work did you do; that is, what was your occupation at the time of the incident? (*For example: plumber, typist, farmer*)
 - What were your usual activities or duties at this job?

Coding System (Code Structure, Rubric)


- **Each code should include:**
 - A number or category, used in statistical analysis
 - A text label, describing all answers in that category
 - **The set of codes for a response should be:**
 - Exhaustive: Every response should be codable into one of the categories
 - Separate codes are needed for skipped, not-asked, off-topic, etc.
 - Exclusive: No response should be codable into more than one category
 - Appropriate to the purposes of the research
 - Use codes that help you answer your research question(s)
 - If you have more than one research question, you might code the same response using different coding systems
 - **Different coders using the same system will produce different codes for the same response**
 - Introduces a kind of “cluster structure” (by coder) into the data
 - Want to construct codes to minimize this (reduce ICC!)
-

Coding Anchoring Vignettes

Respondent 1



Respondent 2



- Respondent 1 would be coded in category **2**
- Respondent 2 would be coded in category **3**

Standard Classification Systems

- If the survey will be compared with other surveys, they should use the same coding scheme
- An ongoing longitudinal or panel survey like NCVS tries to use the same categories in each survey cycle or wave.
- Government or international agencies maintain standard coding systems for common types of information
 - Race/Ethnicity (US Census)
 - Standard Occupational Classification (US Dept of Labor)
 - North American Industry Classification System (US Economic Classification Policy Committee)

Weighting

- Many different sources of weighting in a survey, e.g.
 - Survey design weights
 - In stratified sampling, weights are used to combine stratum means & variance into overall means & variances
 - Other survey designs require weights to account for unequal sampling probabilities, etc.
 - Compute these weights before the data is collected
 - Nonresponse and post-stratification weights
 - Try to adjust sample proportions to equal population proportions
 - Compute these weights after you see the data
- These are discussed in detail in Groves; **below I will only talk about post-stratification weights**

Post-Stratification Weights

- As part of survey data collection it is a good idea to get general demographic information (e.g. in our surveys: sex, age, class, major, hometown, etc.)
- After data collection we compare the proportions in each of these categories in our sample with the same proportions in the population
- If they agree, great. If they disagree, we may re-weight the sample to make them agree
- These categories are called “post-strata”, and the weights are called “post-stratification weights”

Post-Stratification Example

- The 2007 HSS advising satisfaction survey was a simple (no strata, no clusters) web survey of all 986 students in HSS.
- We can separate the responding students by major to see how representative the survey was of each department in HSS.
- If the representativeness was not the same in each department, and if we assume that the nonresponse is ignorable (??) within department, we can re-weight the sample data to get more accurate estimates of population quantities.

HSS Response Rate in Dept Post-Strata

Post-Stratum	Sample	Population	Resp Rate
Economics	40	126	0.32
English	39	115	0.34
History	21	48	0.44
ModLang	8	16	0.50
Philosophy	4	7	0.57
Psychology	37	104	0.36
SDS	54	161	0.34
Statistics	6	8	0.75
Interdisc/IS	76	233	0.33
Undeclared	19	168	0.11
Total	304	986	0.31

HSS Post-Strata Proportions & Weights

Post-Stratum	Sample	Prop	Population	Prop	Weights
Economics	40	0.132	126	0.128	0.97
English	39	0.128	115	0.117	0.91
History	21	0.069	48	0.049	0.70
ModLang	8	0.026	16	0.016	0.62
Philosophy	4	0.013	7	0.007	0.54
Psychology	37	0.122	104	0.105	0.87
SDS	54	0.178	161	0.163	0.92
Statistics	6	0.020	8	0.008	0.41
Interdisc/IS	76	0.250	233	0.236	0.95
Undeclared	19	0.062	168	0.170	2.73
Total	304		986		

$\text{weight} = (\text{Population Proportion}) / (\text{Sample Proportion})$

Fictional Example: What proportion of students think advising is OK?

Post-stratum	Total		Think Advising is OK	
	Sample	Population	Sample	Population
Economics	40	126	28	88
English	39	115	23	69
History	21	48	10	24
ModLang	8	16	3	6
Philosophy	4	7	1	2
Psychology	37	104	11	31
SDS	54	161	22	64
Statistics	6	8	3	4
Interdisc/IS	76	233	46	140
Undeclared	19	168	13	118
Total	304	986	160	546

Population proportion, vs. Unweighted and Weighted sample proportion

- Population proportion:

$$p_{pop} = 546/986 = 0.553$$

- Unweighted Sample proportion:

$$\hat{p} = 160/304 = 0.526$$

- Weighted Sample Proportion

$$\text{Weighted Total} = (0.97)(40) + (0.91)(39) + \dots + (2.73)(19) = 304(!!)$$

$$\text{Weighted OK's} = (0.97)(28) + (0.91)(23) + \dots + (2.73)(13) = 167.45$$

$$\hat{p} = 167.45/304 = 0.551$$

Post-Stratification Weights – Pros & Cons

- Post-stratification weights can fix
 - disproportionate sampling of post strata
 - disproportionate nonresponse across poststrata
- Only works if the sampling/nonresponse process is **ignorable** within post-strata
 - That is, nonresponse does not depend on the answer you would have gotten if the person had responded
- **If the sampling/nonresponse process is non-ignorable then these weights don't work; other weights have to be used**
- The weights are only as good as your model for nonresponse
 - These weights are a very big deal in pre-election phone surveys for example (resp. rate 5-20%; weights account for ignorable and nonignorable nonresponse)

Imputation

- **Weights** are a good solution for unit nonresponse (missed that whole person)
- **Imputation** is a good solution for item nonresponse (person never answered question #17).
- Basic ideas of imputation:
 - Build a model for **what sort of person wouldn't respond**, and use the model to fill in a value for this person
 - Find one or more other people like this person who **did** answer #17, and use their answers for this person
- Alternative to imputation: **Case-wise deletion**
 - Delete this person from the survey so you don't have to deal with the nonresponse to question #17
 - Pro's and con's of case-wise deletion??
 - MCAR: Missing Completely at Random

Mean-value Imputation

- If question #17 is a numerical item, take the average of everyone else's answer to #17, and fill that in for this person
- If question #17 is a yes/no, fill in the proportion of yes's for everyone else (or do a flip of a coin with that probability of "heads")
- Pro's and con's?
- MCAR

Hot-Deck Imputation

- Among all the other people who answered question #17, find the one person who matches this person on important variables
 - age, sex, occupation, answers to other questions, etc. (whatever you think is important to understand this non-response!)
- Fill in that person's answer for this person's #17.
- Pro's? Cons?
- MAR: Missing at Random (within covariates)

Regression Imputation

- Among all the people who answered question #17, fit a regression model (or logistic regression, or whatever) for response to question #17 as a function of other variables:

$$y_{17} = \beta_0 + \beta_1(\text{age}) + \beta_2(\text{sex}) + \beta_3(\text{occupation}) + \beta_4(\text{answer to Q3}) + \dots + \epsilon$$

- Use the fitted model to predict what this person would have answered to #17, and fill that value in
- Pro's? Con's?
- MAR

Limitations of Imputation

- You have to have other variables in the survey that help you build a model for the nonresponse (MAR, or better MCAR)
- Nonignorable missingness (MNAR, missing not at random) is much harder
- After you have filled in the missing data
 - You have NOT increased the sample size; this will matter a lot if you are doing a lot of imputation
 - There is some uncertainty in what value to fill in; this can be accounted for by a technique called **“multiple imputation”**

Variance Calculations

- Final weights in a survey will be a combination (usually multiply together) of
 - Design Weights
 - Nonresponse Weights
 - Post-stratification Weights
- They each improve the point estimate (\bar{y}) but they have different effects on the variance ($Var(\bar{y})$) and standard error
- Additional variance adjustments are made for imputation
- There are not usually closed-form variance formulas (like our simple formulas for strata and clusters)
- More on this next time!

Review/Summary

- The entire survey process consists of
 - Collect/Code/Enter/Edit the data
 - Post-processing of the data
- **Coding** is the process of summarizing complex responses into numbers or categories – subject to bias (bad categories) and variability (coder variation)
- **Post-processing** includes (usually in this order)
 - Imputation
 - Weighting
 - Variance Calculation

These are calculations on the data to account for various weaknesses of the data.