36-463/663: Multilevel & Hierarchical Models

More on Bayes

Brian Junker
132E Baker Hall
brian@stat.cmu.edu

Outline

- Exponential Likelihood, Gamma Prior, & Prof Smedley
- Prior Influence on Posterior
- Comparing 95% intervals
- Conjugate Priors
- Non-Conjugate Priors & WinBUGS
- Normal Model: Estimate Mean, Variance Known
- Shrinkage and the Minnesota Radon Example
- Homework
  - HW07 due today
  - HW08 is out – due Tues!

Exponential Distribution

- Last week we saw that if $x_1, \ldots, x_n \sim$ Expon (λ), i.e. each x_i is indep., with density $f(x|\lambda) = \lambda\, e^{-\lambda x}$, then

$$L(\lambda) = \prod_{i=1}^{n} \lambda e^{-\lambda x_i} = \lambda^n e^{-\lambda \sum_{i=1}^{n} x_i} = \lambda^n e^{-\lambda n \bar{x}}$$

$$\hat{\lambda}_{MLE} = n \Big/ \sum_{i=1}^{n} x_i = 1/\bar{X}$$

and

$$SE(\hat{\lambda}) = \sqrt{1/I(\hat{\lambda})} = \sqrt{\hat{\lambda}^2/n} = \hat{\lambda}/\sqrt{n}$$

MLE point estimate and interval for Prof. Smedley

- We saw that three randomly-chosen students from Prof. Smedley's class took $x_1 = 3$, $x_2 = 10$ and $x_3 = 8$ days to learn boxplots
- The MLE and SE are:

$$\begin{aligned} \hat{\lambda} &= 1/\bar{x} = 1/7 = 0.014 \\ I(\lambda) &= 3/\lambda^2 \\ SE(\hat{\theta}) &= \hat{\lambda}/\sqrt{3} = (1/7)/\sqrt{3} = 0.082 \end{aligned}$$

- and so a rough 95% confidence interval for λ would be [1/7 – 2(1/7)/sqrt(3), 1/7 + 2(1/7)/sqrt(3)], or (-0.022, 0.308)

How does this look as a Bayesian problem?

- Still have $L(\lambda) = \lambda^n e^{-\lambda n\bar{x}}$
- Need a prior distribution for λ. We'll start with a *gamma distribution*:


$$Gamma(\lambda|\alpha,\beta) = \frac{\beta^\alpha}{\Gamma(\alpha)}\lambda^{\alpha-1}e^{-\lambda\beta}$$

- Since (posterior) $\propto$ (likelihood)$\times$(prior),

$$\begin{aligned} f(\lambda|x\text{'s}) &\propto (\lambda^n e^{-\lambda n\bar{x}})(\lambda^{\alpha-1}e^{-\lambda\beta}) = \lambda^{(\alpha+n)-1}e^{-\lambda(\beta+n\bar{x})} \\ &\propto Gamma(\lambda|\alpha+n,\beta+n\bar{x}) \end{aligned}$$

Prior influence on Posterior

- Here are plots for $n = 3, \bar{x} = 7$,
  - α=1, β=2
  - α=4, β=8
  - α=20, β=20
- We see the "shrinkage" idea again: the posterior is "between" the likelihood and the prior
- We see that the location and spread of the prior influences the location and spread of the posterior distribution



Bayesian point estimate and interval for Prof. Smedley

- We will take α=4, β=8, as an example
- There are formulae, but we will simulate

```
> n <- 3
> xbar <- 7
> alpha <- 4
> beta <- 8
> nsim <- 1000
> simdata <- rgamma(nsim,alpha+n,beta+n*xbar)
> quantile(simdata,c(0.025,0.5,0.975))
      2.5%        50%      9.75%
 0.0975616  0.2308562  0.4435332
```

So a point estimate would be 0.23, and a 95% interval would be (0.098, 0.444)

Comparing the MLE and Bayes intervals

- MLE interval: (-0.022, 0.308)
  - Left endpoint can be unrealistic
  - ***"In 95% of experiments, the procedure we used would produce a CI that contains the true value of λ"***
  - ***CI = "confidence interval"***
- Bayesian interval: (0.098, 0.444)
  - Endpoints always in the parameter space
  - ***P[0.098 < λ < 0.444 \| data] = 0.95***
  - ***CI = "credible interval"***

Choosing priors… Conjugate priors

- For the binomial
  - Beta prior: $f(p|\alpha,\beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} p^{\alpha-1}(1-p)^{\beta-1}$
  - Binomial likelihood: $L(p) \propto p^k (1-p)^{n-k}$
  - Beta posterior: $f(p|\text{data}) = Beta(p|\alpha+k, \beta+n-k)$
- For the exponential
  - Gamma prior: $f(\lambda|\alpha,\beta) = \frac{\beta^\alpha}{\Gamma(\alpha)}\lambda^{\alpha-1}e^{-\lambda\beta}$
  - Exponential likelihood: $L(\lambda) = \lambda^n e^{-\lambda n\bar{x}}$
  - Gamma posterior: $f(\lambda|\text{data}) = Gamma(\lambda|\alpha+n, \beta+n\bar{x})$
- ***Conjugate prior***: iff posterior is in same "family"

Non-conjugate priors & WinBUGS

- Conjugate priors made life easy
  - Even with no formulae, using `rbeta()` or `rgamma()` to simulate from the posterior was easy!
- For exponential model, a ***non-conjugate*** choice:
  - Prior: log-normal: $f(\lambda|\mu,\sigma) = \frac{1}{\lambda\sigma\sqrt{2\pi}} \exp\left\{-\frac{(\log\lambda-\mu)^2}{2\sigma^2}\right\}$
  - Likelihood: exponential: $L(\lambda) = \lambda^n e^{-\lambda n\bar{x}}$
  - Posterior: $f(\lambda|\text{data}) \propto \lambda^n e^{-\lambda n\bar{x}} \frac{1}{\lambda\sigma\sqrt{2\pi}} \exp\left\{-\frac{(\log\lambda-\mu)^2}{2\sigma^2}\right\}$
- *WinBUGS is essentially a program for simulating from the posterior, no matter what prior!*

Normal Model: Estimate μ, with σ^2 Known, One Observation x $\sim$ N(μ,σ^2)

- Easy to "see" conjugate prior

$$f(x|\mu) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2\sigma^2}(x-\mu)^2}$$

$$f(\mu) = \frac{1}{\sqrt{2\pi}\tau_0} e^{-\frac{1}{2\tau_0^2}(\mu-\mu_0)^2}$$

$$f(\mu|x) \propto f(x|\mu)f(\mu) \propto \exp\left\{-\frac{1}{2}\left[\frac{(x-\mu)^2}{\sigma^2} + \frac{(\mu-\mu_0)^2}{\tau_0^2}\right]\right\}$$

- Posterior must be normal for μ (quadratic in μ!); to identify it, complete the square…

- The exponent of p(μ|x) looks like -1/2 times

$$\begin{aligned}\frac{(x-\mu)^2}{\sigma^2} + \frac{(\mu-\mu_0)^2}{\tau_0^2} &= \frac{\tau_0^2+\sigma^2}{\tau_0^2\sigma^2}\left[\mu^2 - \frac{2x\mu\tau_0^2 + 2\mu\mu_0\sigma^2}{\tau_0^2+\sigma^2} + \frac{x^2\tau_0^2+\mu_0^2\sigma^2}{\tau_0^2+\sigma^2}\right] \\ &= \frac{\tau_0^2+\sigma^2}{\tau_0^2\sigma^2}\left[\left(\mu - \frac{x\tau_0^2+\mu_0\sigma^2}{\tau_0^2+\sigma^2}\right)^2 + \text{junk}(x,\sigma^2,\mu_0,\tau_0^2)\right] \\ &= \frac{1}{\tau_1^2}(\mu-\mu_1)^2 + (\text{known junk})\end{aligned}$$

so that $\mu|x \sim N(\mu_1, \tau_1^2)$, where

$$\tau_1^2 = \frac{\tau_0^2\sigma^2}{\tau_0^2+\sigma^2} = \frac{1}{1/\sigma^2 + 1/\tau_0^2}$$

$$\mu_1 = \frac{x\tau_0^2+\mu_0\sigma^2}{\tau_0^2+\sigma^2} = \left(\frac{\tau_0^2}{\tau_0^2+\sigma^2}\right)x + \left(\frac{\sigma^2}{\tau_0^2+\sigma^2}\right)\mu_0$$

- The exponent of p(μ|x) looks like -1/2 times

$$\begin{aligned}\frac{(x-\mu)^2}{\sigma^2}+\frac{(\mu-\mu_0)^2}{\tau_0^2} &= \frac{\tau_0^2+\sigma^2}{\tau_0^2\sigma^2}\left[\mu^2-\frac{2x\mu\tau_0^2+2\mu\mu_0\sigma^2}{\tau_0^2+\sigma^2}+\frac{x^2\tau_0^2+\mu_0^2\sigma^2}{\tau_0^2+\sigma^2}\right]\\ &= \frac{\tau_0^2+\sigma^2}{\tau_0^2\sigma^2}\left[\left(\mu-\frac{x\tau_0^2+\mu_0\sigma^2}{\tau_0^2+\sigma^2}\right)^2+\text{junk}(x,\sigma^2,\mu_0,\tau_0^2)\right]\\ &= \frac{1}{\tau_1^2}(\mu-\mu_1)^2+(\text{known junk})\end{aligned}$$

so that $\mu|x \sim N(\mu_1, \tau_1^2)$, where

$$\tau_1^2 = \frac{\tau_0^2\sigma^2}{\tau_0^2+\sigma^2} = \frac{1}{1/\sigma^2+1/\tau_0^2}$$

$$\mu_1 = \frac{x\tau_0^2+\mu_0\sigma^2}{\tau_0^2+\sigma^2} = \left(\frac{\tau_0^2}{\tau_0^2+\sigma^2}\right)x+\left(\frac{\sigma^2}{\tau_0^2+\sigma^2}\right)\mu_0$$

n Observations $x_i \sim N(\mu,\sigma^2)$

- Since

$$p(x_1,\ldots,x_n|\mu) = N(x_1,\ldots,x_n|\mu,\sigma^2) = \prod_{i=1}^{n}\frac{1}{\sqrt{2\pi}\sigma}e^{-\frac{1}{2\sigma^2}(y_i-\mu)^2}$$

$$\propto N(\bar{x}|\mu,\sigma^2/n) \equiv \frac{1}{\sqrt{2\pi\sigma^2/n}}e^{-\frac{1}{2\sigma^2/n}(\bar{y}-\mu)^2}$$

we can apply the results for one observation

- p(x_1, …, x_n|μ) $\propto$ N($\bar{x}$|μ, σ_n^2), $\sigma_n^2 = \sigma^2/n$
- p(μ) = N(μ|μ_o, τ_o^2)
- p(μ|data) = N(μ|μ_n, τ_n^2) where

$$\tau_n^2 = \frac{1}{1/\sigma_n^2+1/\tau_0^2} = \frac{1}{n/\sigma^2+1/\tau_0^2}$$


$$\mu_n = \frac{\bar{x}/\sigma_n^2+\mu_0/\tau_0^2}{1/\sigma_n^2+1/\tau_0^2} = \left(\frac{\tau_0^2}{\tau_0^2+\sigma^2/n}\right)\bar{x}+\left(\frac{\sigma^2/n}{\tau_0^2+\sigma^2/n}\right)\mu_0$$

Normal Mean, Example

- Suppose we know σ=12, we look at n=169 IQ scores, and we find $\bar{x}$ = 100.
- We use as prior N(μ_o, τ_o^2) with μ_o=90, τ_o^2 = 4
- Shrinkage determined by

$$\mu_n = \left(\frac{\tau_0^2}{\tau_0^2+\sigma^2/n}\right)\bar{x}+\left(\frac{\sigma^2/n}{\tau_0^2+\sigma^2/n}\right)\mu_0$$

- $\frac{\tau_0^2}{\tau_0^2+\sigma^2/n}$ is the ***reliability***
- n larger ⇒ reliability larger ⇒ less shrinkage



Minnesota Radon Example

- Emphasize Distribution Structure

$$\text{Level 2: } \mu_i \overset{iid}{\sim} N(\mu_0,\tau_0^2)$$


$$\text{Level 1: } y_{ij} \overset{indep}{\sim} N(\mu_i,\sigma^2)$$

- Emphasize Bayesian point of view (more later!)


$$\text{Prior: } \mu_i \overset{iid}{\sim} N(\mu_0,\tau_0^2)$$

$$\text{Likelihood: } y_{ij} \overset{indep}{\sim} N(\mu_i,\sigma^2)$$

- Emphasize two-stage (multistage) sampling



Minnesota Radon Example



- In each county i with n_i houses, the posterior mean radon level will be

$$\mu_i^{post} = \left(\frac{\tau_0^2}{\tau_0^2 + \sigma^2/n_i}\right)\bar{y}_i + \left(\frac{\sigma^2/n_i}{\tau_0^2 + \sigma^2/n_i}\right)\mu_0$$


  - When n_i large, $\mu_i^{post} \approx \bar{y}_i$
  - When n_i small, $\mu_i^{post} \approx \mu_o$

Minnesota Radon Example

- In the figure, the grand mean is μ_o
- In each county i with n_i houses, posterior mean is

$$\mu_i^{post} = \left(\frac{\tau_0^2}{\tau_0^2 + \sigma^2/n_i}\right)\bar{y}_i + \left(\frac{\sigma^2/n_i}{\tau_0^2 + \sigma^2/n_i}\right)\mu_0$$

  - When n_i large, $\mu_i^{post} \approx \bar{y}_i$
  - When n_i small, $\mu_i^{post} \approx \mu_o$



Summary

- Exponential Likelihood, Gamma Prior, & Prof Smedley
- Prior Influence on Posterior
- Comparing 95% intervals
- Conjugate Priors
- Non-Conjugate Priors & WinBUGS
- Normal Model: Estimate Mean, Variance Known
- Shrinkage and the Minnesota Radon Example
- Homework
  - HW07 due today
  - HW08 is out – due Tues!