36-617: Applied Linear Models Fall 2021

HW10 - Due Wed Nov 17, 11:59pm

- Please turn the homework in to Gradescope using the appropriate link in our course webspace at canvas.cmu.edu, under Assignments.
- · No new reading.
- No quiz on Mon Nov 22.
- Please get started on this assignment right away. Problem #2 may take a while, for example (it's the technical appendix for Project 02!).

Exercises

1. Continuing with the Project 01 data... The file cdi.dat in the Canvas folder for this hw is taken from Kutner et al. (2005): It provides selected county demographic information (CDI) for 440 of the most populous counties in the United States. Each line of the data set has an identification number with a county name and state abbreviation and provides information on 14 variables for a single county. Counties with missing data were deleted from the data set. The information generally pertains to the years 1990 and 1992. The definitions of the variables are given in Table 1 on p. 3 below. For this exercise we only consider a couple of these variables.

Use the function rescale() from library(arm) to rescale pct.hs.grad and per.capita.income as follows:

```
cdi$pci <- rescale(2*cdi$per.cap.income)
cdi$phg <- rescale(2*cdi$pct.hs.grad)</pre>
```

(This rescales the variables to have mean 0 and standard deviation 1; see help(rescale). The rescaling is necessary because lmer() is a somewhat fragile function, and it can fail to converge correctly if the variables are on vastly different scales.)

- (a) Make a facet plot¹ like that on slide 7, lecture 22, using state as the grouping variable, with the following elements:
 - Scatter plot of x=phg and y=pci within each facet (state)
 - The fitted regression line for the completely pooled regression $pci_i = \beta_0 + \beta_1 phg_i + \epsilon_i$, ignoring states, plotted in each facet (state)
 - The fitted regression lines for the completely unpooled regression $pci_i = \beta_{0j[i]} + \beta_{1j[i]}phg_i + \epsilon_i$ where j[i] is the state that observation i is in, and where β_{0j} and β_{1j} are fixed effects, plotted in each facet.
 - The fitted regression lines for the multilevel model

$$\begin{aligned} & \text{pci}_i &= & \alpha_{0j[i]} + \alpha_{1j[i]} \text{phg}_i + \epsilon_i, & \epsilon_i \overset{iid}{\sim} N(0, \sigma^2) \\ & \alpha_{0j} &= & \beta_0 + \eta_{0j}, & \eta_{0j} \overset{iid}{\sim} N(0, \tau_0^2) \\ & \alpha_{1j} &= & \beta_1 + \eta_{1j}, & \eta_{1j} \overset{iid}{\sim} N(0, \tau_1^2) \end{aligned}$$

¹Hint: see sample code in the file 22 - mlm residuals.r.

and where the correlation between η_{0i} and η_{1i} is zero².

(b) Make a new facet plot with state as the grouping variable again, with x equal to the conditional \hat{y} 's, and yequal to the conditional residuals³, within each facet (state).

Submit your code, and the two plots. You do not need to write any commentary.

- 2. Make and submit sections of the technical appendix for your project 02 paper, using the hints and suggestions under Suggestions for Answering the Research Questions in the project-02.pdf assignment file:
 - (a) Section of your technical appendix for Research Question #1.
 - (b) Section of your technical appendix for Research Question #2.
 - (c) Section of your technical appendix for Research Question #3.
 - (d) Section of your technical appendix for Research Question #4.

²Hint: to force the correlation to be zero, rather than specifying the random effects as (1 + phg | state), you should specify (1 | state) + (0 + phg | state).

3Hint: source("residual-functions.r"), and use the appropriate functions defined in that file.

Table 1: Variable definitions for CDI data from Kutner et al. (2005). *Original source:* Geospatial and Statistical Data Center, University of Virginia.

Variable		
Number	Variable Name	Description
1	Identification number	1–440
2	County	County name
3	State	Two-letter state abbreviation
4	Land area	Land area (square miles)
5	Total population	Estimated 1990 population
6	Percent of population aged 18–34	Percent of 1990 CDI population aged 18–34
7	Percent of population 65 or older	Percent of 1990 CDI population aged 65 or old
8	Number of active physicians	Number of professionally active nonfederal physicians during 1990
9	Number of hospital beds	Total number of beds, cribs, and bassinets during 1990
10	Total serious crimes	Total number of serious crimes in 1990, including murder, rape, robbery, aggravated assault, burglary, larceny-theft, and motor vehicle theft, as reported by law enforcement agencies
11	Percent high school graduates	Percent of adult population (persons 25 years old or older) who completed 12 or more years of school
12	Percent bachelor's degrees	Percent of adult population (persons 25 years old or older) with bachelor's degree
13	Percent below poverty level	Percent of 1990 CDI population with income below poverty level
14	Percent unemployment	Percent of 1990 CDI population that is unemployed
15	Per capita income	Per-capita income (i.e. average income per person) of 1990 CDI population (in dollars)
16	Total personal income	Total personal income of 1990 CDI population (in millions of dollars)
17	Geographic region	Geographic region classification used by the US Bureau of the Census, NE (northeast region of the US), NC (north-central region of the US), S (southern region of the US), and W (Western region of the US)