

Chapter 5

Simulation

You will recall from your previous statistics courses that quantifying uncertainty in statistical inference requires us to get at the **sampling distributions** of things like estimators. When the very strong simplifying assumptions of basic statistics courses do not apply¹, or when estimators are themselves complex objects, like kernel regression curves or even histograms, there is little hope of being able to write down sampling distributions in closed form. We get around this by using simulation to approximate the sampling distributions we can't calculate.

5.1 What Do We Mean by “Simulation”?

A stochastic model is a mathematical story about how the data could have been generated. Simulating the model means implementing it, step by step, in order to produce something which should look like the data — what's sometimes called **synthetic data**, or **surrogate data**, or a **realization** of the model. In a stochastic model, some of the steps we need to follow involve a random component, and so multiple simulations starting from exactly the same initial conditions will not give exactly the same outputs or realizations. Rather, will be a distribution over the realizations. Doing large numbers of simulations gives us a good estimate of this distribution.

For a trivial example, consider a model with three random variables, $X_1 \sim \mathcal{N}(\mu_1, \sigma_1^2)$, $X_2 \sim \mathcal{N}(\mu_2, \sigma_2^2)$, with $X_1 \perp\!\!\!\perp X_2$, and $X_3 = X_1 + X_2$. Simulating from this model means drawing a random value from the first normal distribution for X_1 , drawing a second random value for X_2 , and adding them together to get X_3 . The marginal distribution of X_3 , and the joint distribution of (X_1, X_2, X_3) , are implicit in this specification of the model, and we can find them by running the simulation.

In this particular case, we could also find the distribution of X_3 , and the joint distribution, by probability calculations of the kind you learned how to do in your basic probability courses. For instance, X_3 is $\mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$. These analytical

¹In 36-401, you will have seen results about the sampling distribution of linear regression coefficients when the linear model is true, and the noise is Gaussian with constant variance. As an exercise, try to get parallel results when the noise has a t distribution with 10 degrees of freedom.

probability calculations can usually be thought of as just short-cuts for exhaustive simulations.

5.2 How Do We Simulate Stochastic Models?

5.2.1 Chaining Together Random Variables

Stochastic models are usually specified by sets of conditional distributions for one random variable, given some other variable or variables. For instance, a simple linear regression model might have the specification

$$X \sim \mathcal{N}(\mu_x, \sigma_1^2) \quad (5.1)$$

$$Y|X \sim \mathcal{N}(\beta_0 + \beta_1 X, \sigma_2^2) \quad (5.2)$$

If we knew how to generate a random variable from the distributions given on the right-hand sides, we could simulate the whole model by chaining together draws from those conditional distributions. This is in fact the general strategy for simulating any sort of stochastic model, by chaining together random variables.²

What this means is that we can reduce the problem of simulating to that of generating random variables.

5.2.2 Random Variable Generation

5.2.2.1 Built-in Random Number Generators

R provides random number generators for most of the most common distributions. By convention, the names of these functions all begin with the letter “r”, followed by the abbreviation of the functions, and the first argument is always the number of draws to make, followed by the parameters of the distribution:

```
rnorm(n,mean=0,sd=1) # Gaussian
runif(n,min=0,max=1) # Uniform
rexp(n,rate=1)       # Exponential, rate is 1/mean
rpois(n,lambda)      # Poisson, lambda is mean
rbinom(n,size,prob)  # Binomial
```

etc., etc. A further convention is that these parameters can be *vectorized*. Rather than giving a single mean and standard deviation (say) for multiple draws from the Gaussian distribution, each draw can have its own:

```
rnorm(10,mean=1:10,sd=1/sqrt(1:10))
```

That instance is rather trivial, but the exact same principle would be at work here:

²In this case, we could in principle first generate Y , and then draw from $Y|X$, but have fun finding those distributions. Especially have fun if, say, X has a t distribution with 5 degrees of freedom — a very small change to the specification.

```
rnorm(nrow(x), mean=predict(regression.model, newdata=x),
      sd=predict(volatility.model, newdata=x))
```

where `regression.model` and `volatility.model` are previously-defined parts of the model which tell us about conditional expectations and conditional variances.

Of course, none of this explains how R actually draws from any of these distributions; it's all at the level of a black box, which is to say black magic. Because ignorance is evil, and, even worse, *unhelpful* when we need to go beyond the standard distributions, it's worth open the black box at least a little.

5.2.2.2 Transformations

If we can generate a random variable Z with some distribution, and $V = g(Z)$, then we can generate V . So one thing which gets a lot of attention is writing random variables as transformations of one another — ideally as transformations of easy-to-generate variables.

Example. Suppose we can generate random numbers from the standard Gaussian distribution $Z \sim \mathcal{N}(0, 1)$. Then we can generate from $\mathcal{N}(\mu, \sigma^2)$ as $\sigma Z + \mu$. We can generate χ^2 random variables with 1 degree of freedom as Z^2 . We can generate χ^2 random variables with d degrees of freedom by summing d independent copies of Z^2 .

In particular, if we can generate random numbers uniformly distributed between 0 and 1, we can use this to generate anything which is a transformation of a uniform distribution. How far does that extend?

5.2.2.3 Quantile Method

Suppose that we know the **quantile function** Q_Z for the random variable X we want, so that $Q_Z(0.5)$ is the median of X , $Q_Z(0.9)$ is the 90th percentile, and in general $Q_Z(p)$ is bigger than or equal to X with probability p . Q_Z comes as a pair with the cumulative distribution function F_Z , since

$$Q_Z(F_Z(a)) = a, \quad F_Z(Q_Z(p)) = p \quad (5.3)$$

In the **quantile method** (or **inverse distribution transform method**), we generate a uniform random number U and feed it as the argument to Q_Z . Now $Q_Z(U)$ has the distribution function F_Z :

$$\Pr(Q_Z(U) \leq a) = \Pr(F_Z(Q_Z(U)) \leq F_Z(a)) \quad (5.4)$$

$$= \Pr(U \leq F_Z(a)) \quad (5.5)$$

$$= F_Z(a) \quad (5.6)$$

where the last line uses the fact that U is uniform on $[0, 1]$, and the first line uses the fact that F_Z is a non-decreasing function, so $b \leq a$ is true if and only if $F_Z(b) \leq F_Z(a)$.

Example. The CDF of the exponential distribution with rate λ is $1 - e^{-\lambda x}$. The quantile function $Q(p)$ is thus $-\frac{\log(1-p)}{\lambda}$. (Notice that this is positive, because $1-p < 1$ and so $\log(1-p) < 0$, and that it has units of $1/\lambda$, which are the units of x , as it

should.) Therefore, if $U \sim \text{Unif}(0, 1)$, then $-\frac{\log(1-U)}{\lambda} \sim \text{Exp}(\lambda)$. This is the method used by `rexp()`.

Example. The **Pareto distribution** or **power law** is a two-parameter family, $f(x; \alpha, x_0) = \frac{\alpha-1}{x_0} \left(\frac{x}{x_0}\right)^{-\alpha}$ if $x \geq x_0$, with density 0 otherwise. Integration shows that the cumulative distribution function is $F(x; \alpha, x_0) = 1 - \left(\frac{x}{x_0}\right)^{-\alpha+1}$. The quantile function therefore is $Q(p; \alpha, x_0) = x_0(1-p)^{-\frac{1}{\alpha-1}}$. (Notice that this has the same units as x , as it should.)

Example. The standard Gaussian $\mathcal{N}(0, 1)$ does not have a closed form for its quantile function, but there are fast and accurate ways of calculating it numerically (they're what stand behind `qnorm`), so the quantile method can be used. In practice, there are other transformation methods which are even faster, but rely on special tricks.

Since $Q_Z(U)$ has the same distribution function as X , we can use the quantile method, as long as we can calculate Q_Z . Since Q_Z always exists, in principle this solves the problem. In practice, we need to calculate Q_Z before we can use it, and this may not have a closed form, and numerical approximations may be intractable.³

5.2.2.4 Rejection Method

Another general approach, which avoids needing the quantile function, is the **rejection method**. Suppose that we want to generate Z , with probability density function f_Z , and we have a method to generate R , with p.d.f. ρ , called the **proposal distribution**. Also suppose that $f_Z(x) \leq \rho(x)M$, for some constant $M > 1$. For instance, if f_Z has a limited range $[a, b]$, we could take ρ to be the uniform distribution on $[a, b]$, and M the maximum density of f_Z .

The rejection method algorithm then goes as follows.

1. Generate a proposal R from ρ .
2. Generate a uniform U , independently of R .
3. Is $MU\rho(R) < f_Z(R)$?
 - If yes, “accept the proposal” by returning R and stopping.
 - If no, “reject the proposal”, discard R and U , and go back to (1)

If ρ is uniform, this just amounts to checking whether $MU < f_Z(R)$, with M the maximum density of Z .

Computationally, the idea looks like Example 3.

One way to understand the rejection method is as follows. Imagine drawing the curve of $f_Z(x)$. The total area under this curve is 1, because $\int dx f_Z(x) = 1$. The area between any two points a and b on the horizontal axis is $\int_a^b dx f_Z(x) = F_Z(b) - F_Z(a)$. It follows that if we could uniformly sample points from the area between the curve and the horizontal axis, their x coordinates would have exactly the distribution

³In essence, we have to solve the nonlinear equation $F_Z(x) = p$ for x over and over — and that assumes we can easily calculate F_Z .

```

rrejection.1 <- function(dtarget,dproposal,rproposal,M) {
  rejected <- TRUE
  while(rejected) {
    R <- rproposal(1)
    U <- runif(1)
    rejected <- (M*U*dproposal(R) < dtarget(R))
  }
  return(R)
}

rrejection <- function(n,dtarget,dproposal,rproposal,M) {
  replicate(n,rrejection.1(dtarget,dproposal,rproposal,M))
}

```


Code Example 3: An example of how the rejection method would be used. The arguments `dtarget`, `dproposal` and `rproposal` would all be functions. This is not quite industrial-strength code, because it does not let us pass arguments to those functions flexibly. See online code for comments.

function we are looking for. If ρ is a uniform distribution, then we are drawing a rectangle which just encloses the curve of f_Z , sampling points uniformly from the rectangle (with x coordinates R and y coordinates MU), and only keeping the ones which fall under the curve. When ρ is not uniform, but we can sample from it nonetheless, then we are uniformly sampling from the area under $M\rho$, and keeping only the points which are also below f_Z .

Example. The beta distribution, $f(x; a, b) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1} (1-x)^{b-1}$, is defined on the unit interval⁴. While its quantile function can be calculated and so we could use the quantile method, we could also use the reject method, taking the uniform distribution for the proposals. Figure 5.1 illustrates how it would go for the Beta(5,10) distribution

The rejection method's main drawback is speed. The probability of accepting on any given pass through the algorithm is $1/M$. (EXERCISE: Why?) Thus produce n random variables from it takes, on average, nM cycles. (EXERCISE: Why?) Clearly, we want M to be as small, which means that we want the proposal distribution ρ to be close to the target distribution f_Z . Of course if we're using the rejection method because it's hard to draw from the target distribution, and the proposal distribution is close to the target distribution, it may be hard to draw from the proposal.

⁴Here $\Gamma(a) = \int_0^\infty dx e^{-x} x^{a-1}$. It is not obvious, but for integer a , $\Gamma(a) = (a-1)!$. The distribution gets its name because $\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}$ is called the **beta function** of a and b , a kind of continuous generalization of $\binom{a+b}{a}$. The beta distribution arises in connection with problems about minima and maxima, and inference for binomial distributions.



```

M <- 3.3
curve(dbeta(x,5,10),from=0,to=1,ylim=c(0,M))
r <- runif(300,min=0,max=1)
u <- runif(300,min=0,max=1)
below <- which(M*u*dunif(r,min=0,max=1) <= dbeta(r,5,10))
points(r[below],M*u[below],pch="+")
points(r[-below],M*u[-below],pch="-")

```

Figure 5.1: Illustration of the rejection method for generating random numbers from a Beta(5,10) distribution. The proposal distribution is uniform on the range of the beta, which is $[0, 1]$. Points are thus sampled uniformly from the rectangle which runs over $[0, 1]$ on the horizontal axis and $[0, 3.3]$ on the vertical axis, i.e., $M = 3.3$, because the density of the Beta is < 3.3 everywhere. (This is not the lowest possible M but it is close.) Proposed points which fall below the Beta's pdf are marked $+$ and are accepted; those above the pdf curve are marked $-$ and are rejected. In this case, exactly 70% of proposals are rejected.

5.2.2.5 The Metropolis Algorithm and Markov Chain Monte Carlo

One very important, but tricky, way of getting past the limitations of the rejection method is what's called the **Metropolis algorithm**. Once again, we have a density f_Z from which we wish to sample. Once again, we introduce a distribution for “proposals”, and accept or reject proposals depending on the density f_Z . The twist now is that instead of making independent proposals each time, the next proposal depends on the last accepted value — the proposal distribution is a conditional pdf $\rho(r|z)$.

Assume for simplicity that $\rho(r|z) = \rho(z|r)$. (For instance, we could have a Gaussian proposal distribution centered on z .) Then the Metropolis algorithm goes as follows.

1. Start with value Z_0 (fixed or random).
2. Generate R from the conditional distribution $\rho(\cdot|Z_t)$.
3. Generate a uniform U , independent of R .
4. Is $U \leq f_Z(R)/f_Z(Z_t)$?
 - If yes, set $Z_{t+1} = R$ and go to (2)
 - If not, set $Z_{t+1} = Z_t$ and go to (2)

Mostly simply, the algorithm is run until $t = n$, at which point it returns Z_1, Z_2, \dots, Z_n . In practice, better results are obtained if it's run for $n + n_0$ steps, and the first n_0 values of Z are discarded — this is called “burn-in”.

Notice that if $f_Z(R) > f_Z(Z_t)$, then R is always accepted. The algorithm always accepts proposals which move it towards places where the density is higher than where it currently is. If $f_Z(R) < f_Z(Z_t)$, then the algorithm accepts the move with some probability, which shrinks as the density at R gets lower. It should not be hard to persuade yourself that the algorithm will spend more time in places where f_Z is high.

It's possible to say a bit more. Successive values of Z_t are dependent on each other, but $Z_{t+1} \perp\!\!\!\perp Z_{t-1} | Z_t$ — this is a Markov process. The target distribution f_Z is in fact exactly the stationary distribution of the Markov process. If the proposal distributions have broad enough support that the algorithm can get from any z to any z' in a finite number of steps, then the process will “mix”. (In fact we only need to be able to visit points where $f_Z > 0$.) This means that if we start with an arbitrary distribution for Z_0 , the distribution of Z_t approaches f_Z and stays there — the point of burn-in is to give this convergence time to happen. The fraction of time Z_t is close to x is in fact proportional to $f_Z(x)$, so we can use the output of the algorithm as, approximately, so many draws from that distribution.⁵

It would seem that the Metropolis algorithm should be superior to the rejection method, since to produce n random values we need only n steps, or $n + n_0$ to handle burn-in, not nM steps. However, this is deceptive, because if the proposal distribution is not well-chosen, the algorithm ends up staying stuck in the same spot for, perhaps, a very long time. Suppose, for instance, that the distribution is bimodal. If

⁵ And if the dependence between Z_t and Z_{t+1} bothers us, we can always randomly permute them, once we have them.

Z_0 starts out in between the modes, it's easy for it to move rapidly to one peak or the other, and spend a lot of time there. But to go from one mode to the other, the algorithm has to make a series of moves, all in the same direction, which all reduce f_Z , which happens but is unlikely. It thus takes a very long time to explore the whole distribution. The “best” optimal proposal distribution is make $\rho(r|z) = f_Z(r)$, i.e., to just sample from the target distribution. If we could do that, of course, we wouldn't need the Metropolis algorithm, but trying to make ρ close to f_Z is generally a good idea.

The original **Metropolis algorithm** was invented in the 1950s to facilitate designing the hydrogen bomb. It relies on the assumption that the proposal distribution is symmetric, $\rho(r|z) = \rho(z|r)$. It is sometimes convenient to allow an asymmetric proposal distribution, in which case one accepts R if $U \frac{\rho(R|Z_t)}{\rho(Z_t|R)} \leq \frac{f_Z(R)}{f_Z(Z_t)}$. This is called **Metropolis-Hastings**. Both are examples of the broader class of **Markov Chain Monte Carlo** algorithms, where we give up on getting independent samples from the target distribution, and instead make the target the invariant distribution of a Markov process.

5.2.2.6 Generating Uniform Random Numbers

Everything previously to this rested on being able to generate uniform random numbers, so how do we do that? Well, really that's a problem for computer scientists. . . But it's good to understand a little bit about the basic ideas.⁶

First of all, the numbers we get will be produced by some deterministic algorithm, and so will be merely **pseudorandom** rather than truly random. But we would like the deterministic algorithm to produce extremely convoluted results, so that its output *looks* random in as many ways that we can test as possible. Dependencies should be complicated, and correlations between easily-calculated functions of successive pseudorandom numbers should be small and decay quickly. (In fact, “truly random” can be defined, more or less, as the limit of the algorithm becoming infinitely complicated.) Typically, pseudorandom number generators are constructed to produce a sequence of uniform values, starting with an initial value, called the **seed**. In normal operation, the seed is set from the computer's clock; when debugging, the seed can be held fixed, to ensure that results can be reproduced exactly.

Probably the simplest example is **incommensurable rotations**. Imagine a watch which fails very slightly, but deterministically, to keep proper time, so that its second hand advances $\phi \neq 1$ seconds in every real second of time. The position of the watch after t seconds is

$$\theta_t = \theta_0 + t\alpha \bmod 60 \quad (5.7)$$

If ϕ is **commensurable** with 60, meaning $\alpha/60 = k/m$ for some integers k, m , then the positions would just repeat every $60k$ seconds. If α is **incommensurable**, because it is an irrational number, then θ_t never repeats. In this case, not only does θ_t never repeat, but it is uniformly distributed between 0 and 60, in the sense that the fraction of time it spends in any sub-interval is just proportional to the length of the interval. (EXERCISE: Why?)

⁶This section is optional for Spring 2013.

You *could* use this as a pseudo-random number generator, with θ_0 as the seed, but it would not be a very good one, for two reasons. First, exactly representing an irrational number α on a digital computer is impossible, so at best you could use a rational number such that the period $60k$ is large. Second, and more pointedly, the successive θ_t are really too close to each other, and too similar. Even if we only took, say, every 50th value, they'd still be quite correlated with each other.

One way this has been improved is to use *multiple* incommensurable rotations. Say we have a second inaccurate watch, $\phi_t = \phi_0 + \beta t \bmod 60$, where β is incommensurable with both 60 and with α . We record θ_t when ϕ_t is within some small window of 0.⁷

Another approach is to use more aggressively complicated deterministic mappings. Take the system

$$\begin{aligned}\theta_{t+1} &= \theta_t + \phi_t \bmod 1 \\ \phi_{t+1} &= \theta_t + 2\phi_t \bmod 1\end{aligned}\tag{5.8}$$

This is known as “Arnold’s cat map”, after the great Soviet mathematician V. I. Arnold, and Figure 5.2. We can think of this as the second-hand θ_t advancing not by a fixed amount α every second, but by a varying amount ϕ_t . The variable ϕ_t , meanwhile, advances by the amount $\phi_t + \theta_t$. The effect of this is that if we look at only one of the two coordinates, say θ_t , we get a sequence of numbers which, while deterministic, is uniformly distributed, and very hard to predict (Figure 5.3).

5.2.3 Sampling

A complement to drawing from given distributions is to **sample** from a given collection of objects. This is such a common task that R has a handy built-in function to do it:

```
sample(x,size,replace=FALSE,prob=NULL)
```

Here x is a vector which defines the set of objects we’re going to sample from. `size` is the number of samples we want to draw from x . `replace` says whether the samples are drawn with or without replacement. (If `replace=TRUE`, then `size` can be arbitrarily larger than the length of x . If `replace=FALSE`, having a larger `size` doesn’t make sense.) Finally, the optional argument `prob` allows for *weighted* sampling; ideally, `prob` is a vector of probabilities as long as x , giving the probability of drawing each element of x .⁸

As a convenience for a common situation, running `sample` with one argument produces a random permutation of the input, i.e.,

```
sample(x)
```

is equivalent to

⁷The core idea here actually dates back to a medieval astronomer named Nicholas Oresme in the 1300s, as part of an argument that the universe would not repeat exactly (von Plato, 1994, pp. 279–284).

⁸If the elements of `prob` do not add up to 1, but are positive, they will be normalized by their sum, e.g., setting `prob=c(9,9,1)` will assign probabilities $(\frac{9}{19}, \frac{9}{19}, \frac{1}{19})$ to the three elements of x .




Figure 5.2: Effect of the Arnold cat map. The original image is 300×300 , and mapped into the unit square. The cat map is then applied to the coordinates of each pixel separately, giving a new pixel which inherits the old color. (This can most easily be seen in the transition from the original to time 1.) The original image re-assembles itself at time 300 because all the original coordinates are multiples of $1/300$. If we had sampled every, say, 32 time-steps, it would have taken much longer to see a repetition. In the meanwhile, following the x coordinate of a single pixel from the original image would provide a very creditable sequence of pseudo-random values. (Figure from Wikipedia, s.v. “Arnold’s cat map”. See also <http://math.gmu.edu/~sander/movies/arnold.html>.)


```

arnold.map <- function(v) {
  theta <- v[1]
  phi <- v[2]
  theta.new <- (theta+phi)%%1
  phi.new <- (theta+2*phi)%%1
  return(c(theta.new,phi.new))
}

rarnold <- function(n,seed) {
  z <- vector(length=n)
  for (i in 1:n) {
    seed <- arnold.map(seed)
    z[i] <- seed[1]
  }
  return(z)
}

```

Code Example 4: A function implementing the Arnold cat map (Eq. 5.9), and a second function which uses it as a pseudo-random number generator. See online version for comments.



```

par(mfrow=c(2,1))
z <- rarnold(1000,c(0.11124,0.42111))
hist(z,probability=TRUE)
plot(z[-1000],z[-1],xlab=expression(Z[t]),ylab=expression(Z[t+1]))

```

Figure 5.3: Left: histogram from 1000 samples of the θ coordinate of the Arnold cat map, started from (0.11124,0.42111). Right: scatter-plot of successive values from the sample, showing that the dependence is very subtle.

```
sample(x,size=length(x),replace=FALSE)
```

For example, if we're doing five-fold cross-validation, then

```
sample(rep(1:5,length.out=nrow(df)))
```

will first repeat the numbers 1,2,3,4,5 until we have one number for each row of `df`, and then shuffle the order of those numbers randomly. This then would give an assignment of each row of `df` to one (and only one) of five folds.

5.2.3.1 Sampling Rows from Data Frames

When we have multivariate data (which is the usual situation), we typically arrange it into a data-frame, where each row records one unit of observation, with multiple interdependent columns. The natural notion of sampling is then to draw a random sample of the data points, which in that representation amounts to a random sample of the rows. We can implement this simply by sampling row *numbers*. For instance, this command,

```
df[sample(1:nrow(df),size=b),]
```

will create a new data frame from `b`, by selecting `b` rows from `df` without replacement. It is an easy exercise to figure out how to sample from a data frame *with* replacement, and with unequal probabilities per row.

5.2.3.2 Multinomials and Multinoullis

If we want to draw one value from a multinomial distribution with probabilities $p = (p_1, p_2, \dots, p_k)$, then we can use `sample`:

```
sample(1:k,size=1,prob=p)
```

If we want to simulate a “multinoulli” process⁹, i.e., a sequence of independent and identically distributed multinomial random variables, then we can easily do so:

```
rmultinoulli <- function(n,prob) {
  k <- length(prob)
  return(sample(1:k,size=n,replace=TRUE,prob=prob))
}
```

5.2.3.3 Probabilities of Observation

Often, our models of how the data are generated will break up into two parts. One part is a model of how actual variables are related to each other out in the world. (E.g., we might model how education and racial categories are related to occupation, and occupation is related to income.) The other part is a model of how variables come to be recorded in our data, and the distortions they might undergo in the course of doing so. (E.g., we might model the probability that someone appears in a survey

⁹A handy term I learned from Gustavo Lacerda.

as a function of race and income.) Plausible sampling mechanisms often make the probability of appearing in the data a function of some of the variables. This can then have important consequences for our inferences from the data we happen to see to the whole population or process.

```
income <- rnorm(n,mean=predict(income.model,x),sd=sigma)
capture.proBABILITIES <- predict(observation.model,x)
observed.income <- sample(income,size=b,prob=capture.proBABILITIES)
```

5.2.4 Repeating Simulations

Because simulations are often most useful when they are repeated many times, R has a command to repeat a whole block of code:

```
replicate(n,expr)
```

Here `expr` is some executable “expression” in R, basically something you could type in the terminal without trouble, and `n` is the number of times to repeat it.

For instance,

```
output <- replicate(1000,rnorm(length(x),beta0+beta1*x,sigma))
```

will replicate, 1000 times, sampling from the predictive distribution of a Gaussian linear regression model. Conceptually, this is equivalent to doing something like

```
output <- matrix(0,nrow=1000,ncol=length(x))
for (i in 1:1000) {
  output[i,] <- rnorm(length(x),beta0+beta1*x,sigma)
}
```

but the `replicate` version has two great advantages. First, it is faster, because R processes it with specially-optimized code. (Loops are especially slow in R.) Second, and *far* more importantly, it is *clearer*: it makes it obvious what is being done, in one line, and leaves the computer to figure out the boring and mundane details of how best to implement it.

5.3 Why Simulate?

There are three major uses for simulation: to understand a model, to check it, and to fit it.

5.3.1 Understanding the Model; Monte Carlo

We understand a model by seeing what it predicts about the variables we care about, and the relationships between them. Sometimes those predictions are easy to extract from a mathematical representation of the model, but often they aren’t. With a model we can simulate, however, we can just run the model and see what happens.

Our stochastic model gives a distribution for some random variable Z , which in general is a complicated, multivariate object with lots of interdependent components. We may also be interested in some complicated function g of Z , such as, say, the ratio of two components of Z , or even some nonparametric curve fit through the data points. How do we know what the model says about g ?

Assuming we can make draws from the distribution of Z , we can find the distribution of any function of it we like, to as much precision as we want. Suppose that $\tilde{Z}_1, \tilde{Z}_2, \dots, \tilde{Z}_b$ are the outputs of b independent runs of the model — b different *replicates* of the model. (I am using the tilde to remind us that these are just simulations.) We can calculate g on each of them, getting $g(\tilde{Z}_1), g(\tilde{Z}_2), \dots, g(\tilde{Z}_b)$. If averaging makes sense for these values, then

$$\frac{1}{b} \sum_{i=1}^b g(\tilde{Z}_i) \xrightarrow{b \rightarrow \infty} \mathbf{E}[g(Z)] \quad (5.9)$$

by the law of large numbers. So simulation and averaging lets us get expectation values. This basic observation is the seed of the **Monte Carlo method**.¹⁰ If our simulations are independent, we can even say that $\frac{1}{b} \sum_{i=1}^b g(\tilde{Z}_i)$ has approximately the distribution $\mathcal{N}(\mathbf{E}[g(Z)], \text{Var}[g(Z)]/b)$ by the central limit theorem. Of course, if you can get expectation values, you can also get variances. (This is handy if trying to apply the central limit theorem!) You can also get any higher moments — if you need the kurtosis for whatever reason, you just have to simulate enough.

You can also pick any set s and get the probability that $g(Z)$ falls into that set:

$$\frac{1}{b} \sum_{i=1}^b \mathbf{1}_s(g(\tilde{Z}_i)) \xrightarrow{b \rightarrow \infty} \Pr(g(Z) \in s) \quad (5.10)$$

The reason this works is of course that $\Pr(g(Z) \in s) = \mathbf{E}[\mathbf{1}_s(g(Z))]$, so it's just the central limit theorem. So we can get the whole distribution of any complicated function of the model that we want, as soon as we can simulate the model. It is really only a little harder to get the complete sampling distribution than it is to get the expectation value, and the exact same ideas apply.

5.3.2 Checking the Model

An important but under-appreciated use for simulation is to *check* models after they have been fit. If the model is right, after all, it represents the mechanism which generates the data. This means that when we simulate, we run that mechanism, and the surrogate data which comes out of the machine should look like the real data. More exactly, the real data should look like a typical realization of the model. If it does not, then the model's account of the data-generating mechanism is systematically wrong

¹⁰The name comes from the physicists who used the method to do calculations relating to designing the hydrogen bomb; see Metropolis *et al.* (1953). Folklore among physicists says that the method goes back at least to Enrico Fermi in the 1930s, without the cutesy name.

```

rgeyser <- function() {
  n <- nrow(geyser)
  sigma <- summary(fit.ols)$sigma
  new.waiting <- rnorm(n,mean=fitted(fit.ols),sd=sigma)
  new.geyser <- data.frame(duration=geyser$duration,
    waiting=new.waiting)
  return(new.geyser)
}

```

Code Example 5: Function for generating surrogate data sets from the linear model fit to `geyser`.

in some way. By carefully choosing the simulations we perform, we can learn a lot about how the model breaks down and how it might need to be improved.¹¹

Often the comparison between simulations and data can be done qualitatively and visually. For example, a classic data set concerns the time between eruptions of the Old Faithful geyser in Yellowstone, and how they relate to the duration of the latest eruption. A common exercise is to fit a regression line to the data by ordinary least squares:

```

library(MASS)
data(geyser)
fit.ols <- lm(waiting~duration,data=geyser)


```

Figure 5.4 shows the data, together with the OLS regression line. It doesn't look that great, but if someone insisted it was a triumph of quantitative vulcanology, how could you show they were wrong?

Well, OLS is usually presented as part of a probability model for the response conditional on the input, with Gaussian and homoskedastic noise. In this case, the probability model is $\text{waiting} = \beta_0 + \beta_1 \text{duration} + \epsilon$, with $\epsilon \sim \mathcal{N}(0, \sigma^2)$. If we simulate from this probability model, we'll get something we can compare to the actual data, to help us assess whether the scatter around that regression line is really bothersome. Since OLS doesn't require us to assume a distribution for the input variable (here, `duration`), the simulation function in Code Example 5 leaves those values alone, but regenerates values of the response (`waiting`) according the model assumptions.


A useful principle for model checking is that if we do some exploratory data analyses of the real data, doing the same analyses to realizations of the model should give roughly the same results. This isn't really the case here. Figure 5.5 shows the actual density of `waiting`, plus the density produced by simulating — reality is clearly bimodal, but the model is unimodal. Similarly, Figure 5.6 shows the real data, the OLS line, and a simulation from the OLS model. It's visually clear that the deviations of the real data from the regression line are both bigger and more patterned than those we get from simulating the model, so something is wrong with the latter.

¹¹“Might”, because sometimes we're better off with a model that makes systematic mistakes, if they're small and getting it right would be a hassle.



```
plot(geyser$duration,geyser$waiting,xlab="duration",ylab="waiting")
abline(fit.ols)
```

Figure 5.4: Data for the geyser data set, plus the OLS regression line.




```
plot(density(geyser$waiting),xlab="waiting",main="",sub="")
lines(density(rgeyser())$waiting),lty=2)
```

Figure 5.5: Actual density of the waiting time between eruptions (solid curve) and that produced by simulating the OLS model (dashed).

By itself, just seeing that data doesn't look like a realization of the model isn't super informative, since we'd really like to know *how* the model's broken, and how to fix it. Further simulations, comparing analyses of the data to analyses of the simulation output, are often very helpful here. Looking at Figure 5.6, we might suspect that one problem is heteroskedasticity — the variance isn't constant. This suspicion is entirely correct, and will be explored in §7.3.2.

5.3.3 Sensitivity Analysis

Often, the statistical inference we do on the data is predicated on certain assumptions about how the data is generated. For instance, if we have missing values for some variables and just *ignore* incomplete rows, we are implicitly assuming that data are “missing at random”, rather than in some systematic way. If we are not *totally* confident in such assumptions, we might wish to see what happens they break down. That is, we set up a model where the assumptions are more or less violated, and then run our *original* analysis on the simulation output. Because it's a simulation, we know the complete truth about the data-generating process, and can assess how far off our inferences are. In favorable circumstances, our inferences don't mess up too much even when the assumptions we used to motivate the analysis are badly wrong. Sometimes, however, we discover that even tiny violations of our initial assumptions lead to large errors in our inferences. Then we either need to make some compelling case for those assumptions, or be *very* cautious in our inferences.



```
plot(geyser$duration,geyser$waiting,xlab="duration",ylab="waiting")
abline(fit.ols)
points(rgeyser(),pch=20,cex=0.5)
```

Figure 5.6: As in Figure 5.4, plus one realization of simulating the OLS model (small black dots).

5.4 The Method of Simulated Moments

Checking whether the model's simulation output looks like the data naturally suggests the idea of *adjusting* the model until it does. This becomes a way of estimating the model — in the jargon, **simulation-based inference**. All forms of this involve adjusting parameters of the model until the simulations *do* look like the data. They differ in what “look like” means, concretely. The most straightforward form of simulation-based inference is the **method of simulated moments**.¹²

5.4.1 The Method of Moments

You will have seen the ordinary **method of moments** in earlier statistics classes. Let's recall the general setting. We have a model with a parameter vector θ , and pick a vector m of moments to calculate. The moments, like the expectation of any variables, are functions of the parameters,

$$m = g(\theta) \quad (5.11)$$

for some function g . If that g is invertible, then we can recover the parameters from the moments,

$$\theta = g^{-1}(m) \quad (5.12)$$

The method of moments *estimator* takes the observed, sample moments \hat{m} , and plugs them into Eq. 5.12:

$$\widehat{\theta}_{MM} = g^{-1}(\hat{m}) \quad (5.13)$$

What if g^{-1} is hard to calculate — if it's hard to explicitly solve for parameters from moments? In that case, we can use minimization:

$$\widehat{\theta}_{MM} = \underset{\theta}{\operatorname{argmin}} \|g(\theta) - \hat{m}\|^2 \quad (5.14)$$

For the minimization version, we just have to calculate moments from parameters $g(\theta)$, not vice versa. To see that Eqs. 5.13 and 5.14 do the same thing, notice that (i) the squared¹³ distance $\|g(\theta) - \hat{m}\|^2 \geq 0$, (ii) the distance is only zero when the moments are matched exactly, and (iii) there is only θ which will match the moments.

In either version, the method of moments works *statistically* because the sample moments \hat{m} converge on their expectations $g(\theta)$ as we get more and more data. This is, to repeat, a consequence of the law of large numbers.

It's worth noting that nothing in this argument says that m has to be a vector of *moments* in the strict sense. They could be expectations of any functions of the random variables, so long as $g(\theta)$ is invertible, we can calculate the sample expectations of these functions from the data, and the sample expectations converge. When m isn't just a vector of moments, then, we have the **generalized method of moments**.

¹²This section is optional for spring 2013.

¹³Why squared? Basically because it makes the function we're minimizing smoother, and the optimization nicer.

It is also worth noting that there's a somewhat more general version of the same method, where we minimize

$$(g(\theta) - \hat{m}) \cdot \mathbf{w} (g(\theta) - \hat{m}) \quad (5.15)$$

with some positive-definite weight matrix \mathbf{w} . This can help if some of the moments are much more sensitive to the parameters than others. But this goes beyond what we really need here.

5.4.2 Adding in the Simulation

All of this supposes that we know how to calculate $g(\theta)$ — that we can find the moments exactly. Even if this is too hard, however, we could always *simulate* to approximate these expectations, and try to match the simulated moments to the real ones. Rather than Eq. 5.14, the estimator would be

$$\widehat{\theta}_{SMM} = \operatorname{argmin}_{\theta} \left\| \tilde{g}_{s,T}(\theta) - \hat{m} \right\|^2 \quad (5.16)$$

with s being the number of simulation paths and T being their size. Now consistency requires that $\tilde{g} \rightarrow g$, either as T grows or s or both, but this is generally assured by the law of large numbers, as we talked about earlier. Simulated method of moments estimates like this are generally more uncertain than ones which don't rely on simulation, since it introduces an extra layer of approximation, but this can be reduced by increasing s .¹⁴

5.4.3 An Example: Moving Average Models and the Stock Market

To give a concrete example, we will try fitting a time series model to the stock market: it's a familiar subject which interests *most* students, and we can check the method of simulated moments here against other estimation techniques.¹⁵

Our data will consist of about ten year's worth of daily values for the S&P 500 stock index, available on the class website:

```
sp <- read.csv("SPhistory.short.csv")
# We only want closing prices
sp <- sp[,7]
# The data are in reverse chronological order, which is weird for us
sp <- rev(sp)
# And in fact we only want log returns, i.e., difference in logged prices
sp <- diff(log(sp))
```

¹⁴A common trick is to fix T at the actual sample size n , and then to increase s as much as computationally feasible. By looking at the variance of \tilde{g} across different runs of the model with the same θ , one gets an idea of how much uncertainty there is in \hat{m} itself, and so of how precisely one should expect to be able to match it. If the optimizer has gotten $|\tilde{g}(\theta) - \hat{m}|$ down to 0.02, and the standard deviation of \tilde{g} at constant θ is 0.1, further effort at optimization is probably wasted.

¹⁵Nothing in what follows, or in the homework, could actually be used to make money, however.

Professionals in finance do not care so much about the sequence of **prices** P_t , as the sequence of **returns**, $\frac{P_t - P_{t-1}}{P_{t-1}}$. This is because making \$1000 is a lot better when you invested \$1000 than when you invested \$1,000,000, but 10% is 10%. In fact, it's often easier to deal with the **log returns**, $X_t = \log \frac{P_t}{P_{t-1}}$, as we do here.

The model we will fit is a **first-order moving average**, or MA(1), model:

$$X_t = Z_t + \theta Z_{t-1} \quad (5.17)$$

$$Z_t \sim \mathcal{N}(0, \sigma^2) \text{ i.i.d.} \quad (5.18)$$

The X_t sequence of variables are the returns we see; the Z_t variables are invisible to us. The interpretation of the model is as follows. Prices in the stock market change in response to news that affects the prospects of the companies listed, as well as news about changes in over-all economic conditions. Z_t represents this flow of news, good and bad. It makes sense that Z_t is uncorrelated, because the relevant part of the news is only what everyone hadn't already worked out from older information¹⁶. However, it does take some time for the news to be assimilated, and this is why Z_{t-1} contributes to X_t . A negative contribution, $\theta < 0$, would seem to indicate a "correction" to the reaction to the previous day's news.

Mathematically, notice that since Z_t and θZ_{t-1} are independent Gaussians, X_t is a Gaussian with mean 0 and variance $\sigma^2 + \theta^2 \sigma^2$. The marginal distribution of X_t is therefore the same for all t . For technical reasons¹⁷, we can really only get sensible behavior from the model when $-1 \leq \theta \leq 1$.

There are two parameters, θ and σ^2 , so we need two moments for estimation. Let's try $\text{Var}[X_t]$ and $\text{Cov}[X_t, X_{t-1}]$.

$$\text{Var}[X_t] = \text{Var}[Z_t] + \theta^2 \text{Var}[Z_{t-1}] \quad (5.19)$$

$$= \sigma^2 + \theta^2 \sigma^2 \quad (5.20)$$

$$= \sigma^2(1 + \theta^2) \equiv v(\theta, \sigma) \quad (5.21)$$

(This agrees with our earlier reasoning about Gaussians, but doesn't need it.)

$$\text{Cov}[X_t, X_{t-1}] = \mathbf{E}[(Z_t + \theta Z_{t-1})(Z_{t-1} + \theta Z_{t-2})] \quad (5.22)$$

$$= \theta \mathbf{E}[Z_{t-1}^2] \quad (5.23)$$

$$= \theta \sigma^2 \equiv c(\theta, \sigma) \quad (5.24)$$

We can solve the system of equations for the parameters, starting with eliminating

¹⁶Nobody will ever say "What? It's snowing in Pittsburgh in *February*? I must call my broker!"

¹⁷Think about trying to recover Z_t , if we knew θ . One might try $X_t - \theta X_{t-1}$, which is almost right, it's $Z_t + \theta Z_{t-1} - \theta Z_{t-1} - \theta^2 Z_{t-2} = Z_t - \theta^2 Z_{t-2}$. Similarly, $X_t - \theta X_{t-1} + \theta^2 X_{t-2} = Z_t + \theta^3 Z_{t-2}$, and so forth. If $|\theta| < 1$, then this sequence of approximations will converge on Z_t ; if not, then not. It turns out that models which are not "invertible" in this way are very strange — see Shumway and Stoffer (2000).

σ^2 :

$$\frac{c(\theta, \sigma)}{v(\theta, \sigma)} = \frac{\sigma^2 \theta}{\sigma^2(1 + \theta^2)} \quad (5.25)$$

$$= \frac{\theta}{1 + \theta^2} \quad (5.26)$$

$$0 = \theta^2 \frac{c}{v} - \theta + \frac{c}{v} \quad (5.27)$$

This is a quadratic in θ ,

$$\theta = \frac{1 \pm \sqrt{1 - 4 \frac{c^2}{v^2}}}{2c/v} \quad (5.28)$$

and it's easy to confirm¹⁸ that this has only one solution in the meaningful range, $-1 \leq \theta \leq 1$. Having found θ , we solve for σ^2 ,

$$\sigma^2 = c/\theta \quad (5.29)$$

The method of moments estimator takes the sample values of these moments, \hat{v} and \hat{c} , and plugs them in to Eqs. 5.28 and 5.29. With the S&P returns, the sample covariance is -1.61×10^{-5} , and the sample variance 1.96×10^{-4} . This leads to $\hat{\theta}_{MM} = -8.28 \times 10^{-2}$, and $\hat{\sigma}_{MM}^2 = 1.95 \times 10^{-4}$. In terms of the model, then, each day's news has a follow-on impact on prices which is about 8% as large as its impact the first day, but with the opposite sign.¹⁹

If we did not know how to solve a quadratic equation, we could use the minimization version of the method of moments estimator:

$$\left[\begin{array}{c} \hat{\theta}_{MM} \\ \hat{\sigma}_{MM}^2 \end{array} \right] = \underset{\theta, \sigma^2}{\operatorname{argmin}} \left\| \begin{array}{c} \sigma^2 \theta - \hat{c} \\ \sigma^2(1 + \theta^2) - \hat{v} \end{array} \right\|^2 \quad (5.30)$$

Computationally, it would go something like Code Example 6.

The parameters estimated by minimization agree with those from direct algebra to four significant figures, which I hope is good enough to reassure you that this works.

Before we can try out the method of simulated moments, we have to figure out how to simulate our model. X_t is a deterministic function of Z_t and Z_{t-1} , so our general strategy says to first generate the Z_t , and then compute X_t from that. But here the Z_t are just a sequence of independent Gaussians, which is a solved problem for us. The one wrinkle is that to get our first value X_1 , we need a previous value Z_0 . Code Example 7 shows the solution.

¹⁸For example, plot c/v as a function of θ , and observe that any horizontal line cuts the graph at only one point.

¹⁹It would be natural to wonder whether $\hat{\theta}_{MM}$ is really significantly different from zero. Assuming Gaussian noise, one could, in principle, calculate the probability that even though $\theta = 0$, by chance \hat{c}/\hat{v} was so far from zero as to give us our estimate. As you will see in the homework, however, Gaussian assumptions are very bad for this data. This sort of thing is why we have bootstrapping.

```

ma.mm.est <- function(c,v) {
  theta.0 <- c/v
  sigma2.0 <- v
  fit <- optim(par=c(theta.0,sigma2.0), fn=ma.mm.objective,
              c=c, v=v)
  return(fit)
}

ma.mm.objective <- function(params,c,v) {
  theta <- params[1]
  sigma2 <- params[2]
  c.pred <- theta*sigma2
  v.pred <- sigma2*(1+theta^2)
  return((c-c.pred)^2 + (v-v.pred)^2)
}

```

Code Example 6: Code for implementing method of moments estimation of a first-order moving average model, as in Eq. 5.30. See Appendix 5.6 for “design notes”, and the online code for comments.

```

rma <- function(n,theta,sigma2,s=1) {
  z <- replicate(s,rnorm(n=n+1,mean=0,sd=sqrt(sigma2)))
  x <- z[-1,] + theta*z[-(n+1),]
  return(x)
}

```

Code Example 7: Function which simulates s independent runs of a first-order moving average model, each of length n , with given noise variance σ^2 and after-effect θ . See online for the version with comments on the code details.

```

sim.var <- function(n,theta,sigma2,s=1) {
  vars <- apply(rma(n,theta,sigma2,s),2,var)
  return(mean(vars))
}

sim.cov <- function(n,theta,sigma2,s=1) {
  x <- rma(n,theta,sigma2,s)
  covs <- colMeans(x[-1,]*x[-n,])
  return(mean(covs))
}

```

Code Example 8: Functions for calculating the variance and covariance for specified parameter values from simulations.

What we need to extract from the simulation are the variance and the covariance. It will be more convenient to have functions which calculate these call `rma()` themselves (Code Example 8).

Figure 5.7 plots the covariance, the variance, and their ratio as functions of θ with $\sigma^2 = 1$, showing both the values obtained from simulation and the theoretical ones.²⁰ The agreement is quite good, though of course not quite perfect.²¹

Conceptually, we could estimate θ by just taking the observed value \hat{c}/\hat{v} , running a horizontal line across Figure 5.7c, and seeing at what θ it hit one of the simulation dots. Of course, there might not be one it hits *exactly*...


The more practical approach is Code Example 9. The code is practically identical to that in Code Example 6, except that the variance and covariance predicted by given parameter settings now come from simulating those settings, not an exact calculation. Also, we have to say how long a simulation to run, and how many simulations to average over per parameter value.

When I run this, with $s=100$, I get $\hat{\theta}_{MSM} = -8.36 \times 10^{-2}$ and $\hat{\sigma}_{MSM}^2 = 1.94 \times 10^{-4}$, which is quite close to the non-simulated method of moments estimate. In fact, in this case there is actually a maximum likelihood estimator (`arima()`, after the more general class of models including MA models), which claims $\hat{\theta}_{ML} = -9.75 \times 10^{-2}$ and $\hat{\sigma}_{ML}^2 = 1.94 \times 10^{-4}$. Since the standard error of the MLE on θ is ± 0.02 , this is working essentially as well as the method of moments, or even the method of simulated moments.

In this case, because there *is* a very tractable maximum likelihood estimator, one generally wouldn't use the method of simulated moments. But we can in this case check whether it works (it does), and so we can use the same technique for other models, where an MLE is unavailable.

²⁰I could also have varied σ^2 and made 3D plots, but that would have been more work. Also, the variance and covariance are both proportional to σ^2 , so the shapes of the figures would all be the same.

²¹If you look at those figures and think "Why not do a nonparametric regression of the simulated moments against the parameters and use the fitted values as \hat{g} , it'll get rid of some of the simulation noise?", congratulations, you've just discovered the smoothed method of simulated moments.



```
theta.grid <- seq(from=-1,to=1,length.out=300)
cov.grid <- sapply(theta.grid,sim.cov,sigma2=1,n=length(sp),s=10)
plot(theta.grid,cov.grid,xlab=expression(theta),ylab="Covariance")
abline(0,1,col="grey",lwd=3)
var.grid <- sapply(theta.grid,sim.var,sigma2=1,n=length(sp),s=10)
plot(theta.grid,var.grid,xlab=expression(theta),ylab="Variance")
curve((1+x^2),col="grey",lwd=3,add=TRUE)
plot(theta.grid,cov.grid/var.grid,xlab=expression(theta),
      ylab="Ratio of covariance to variance")
curve(x/(1+x^2),col="grey",lwd=3,add=TRUE)
```

Figure 5.7: Plots of the covariance, the variance, and their ratio as a function of θ , with $\sigma^2 = 1$. Dots show simulation values (averaging 10 realizations each as long as the data), the grey curves the exact calculations.

```
ma.msm.est <- function(c,v,n,s) {  
  theta.0 <- c/v  
  sigma2.0 <- v  
  fit <- optim(par=c(theta.0,sigma2.0),fn=ma.msm.objective,c=c,v=v,n=n,s=s)  
  return(fit)  
}  
  
ma.msm.objective <- function(params,c,v,n,s) {  
  theta <- params[1]  
  sigma2 <- params[2]  
  c.pred <- sim.cov(n,theta,sigma2,s)  
  v.pred <- sim.var(n,theta,sigma2,s)  
  return((c-c.pred)^2 + (v-v.pred)^2)  
}
```

Code Example 9: Code for implementing the method of simulated moments estimation of a first-order moving average model.

5.5 Exercises

To think through, not to hand in.

Section 5.4 explained the method of simulated moments, where we try to match expectations of various functions of the data. Expectations of functions are summary statistics, but they're not the *only* kind of summary statistics. We could try to estimate our model by matching any set of summary statistics, so long as (i) there's a unique way of mapping back from summaries to parameters, and (ii) estimates of the summary statistics converge as we get more data.

A powerful but somewhat paradoxical version of this is what's called **indirect inference**, where the summary statistics are the parameters of a *different* model. This second or **auxiliary** model does *not* have to be correctly specified, it just has to be easily fit to the data, and satisfy (i) and (ii) above. Say the parameters of the auxiliary model are β , as opposed to the θ of our real model. We calculate $\hat{\beta}$ on the real data. Then we simulate from different values of θ , fit the auxiliary to the simulation outputs, and try to match the auxiliary estimates. Specifically, the indirect inference estimator is

$$\hat{\theta}_{II} = \underset{\theta}{\operatorname{argmin}} \|\tilde{\beta}(\theta) - \hat{\beta}\|^2 \quad (5.31)$$

where $\tilde{\beta}(\theta)$ is the value of β we estimate from a simulation of θ , of the same size as the original data. (We might average together a couple of simulation runs for each θ .) If we have a consistent estimator of β , then

$$\hat{\beta} \rightarrow \beta \quad (5.32)$$

$$\tilde{\beta}(\theta) \rightarrow b(\theta) \quad (5.33)$$

If in addition $b(\theta)$ is invertible, then

$$\hat{\theta}_{II} \rightarrow \theta \quad (5.34)$$

For this to work, the auxiliary model needs to have at least as many parameters as the real model, but we can often arrange this by, say, making the auxiliary model a linear regression with a lot of coefficients.

A specific case, often useful for time series, is to make the auxiliary model an **autoregressive model**, where each observation is linearly regressed on the previous ones. A first-order autoregressive model (or "AR(1)") is

$$X_t = \beta_0 + \beta_1 X_{t-1} + \epsilon_t \quad (5.35)$$

where $\epsilon_t \sim \mathcal{N}(0, \beta_3)$. (So an AR(1) has three parameters.)

1. Convince yourself that if X_t comes from an MA(1) process, it can't also be written as an AR(1) model.
2. Write a function, `ar1.fit`, to fit an AR(1) model to a time series, using `lm`, and to return the three parameters (intercept, slope, noise variance).

3. Apply `ar1.fit` to the S&P 500 data; what are the auxiliary parameter estimates?
4. Combine `ar1.fit` with the simulator `rma`, and plot the three auxiliary parameters as functions of θ , holding σ^2 fixed at 1. (This is analogous to Figure 5.7.)
5. Write functions, analogous to `ma.msm.est` and `ma.msm.objective`, for estimating an MA(1) model, using an AR(1) model as the auxiliary function. Does this recover the right parameter values when given data simulated from an MA(1) model?
6. What values does your estimator give for θ and σ^2 on the S&P 500 data? How do they compare to the other estimates?

5.6 Appendix: Some Design Notes on the Method of Moments Code

Go back to Section 5.4.3 and look at the code for the method of moments. There’ve been a fair amount of questions about writing code, and this is a useful example.

The first function, `ma.mm.est`, estimates the parameters taking as inputs two numbers, representing the covariance and the variance. The real work is done by the built-in `optim` function, which itself takes two major arguments. One, `fn`, is the function to optimize. Another, `par`, is an initial guess about the parameters at which to begin the search for the optimum.²²

The `fn` argument to `optim` must be a function, here `ma.mm.objective`. The first argument to that function has to be a vector, containing all the parameters to be optimized over. (Otherwise, `optim` will quit and complain.) There can be other arguments, not being optimized over, to that function, which `optim` will pass along, as you see here. `optim` will also accept a lot of optional arguments to control the search for the optimum — see `help(optim)`.

All `ma.mm.objective` has to do is calculate the objective function. The first two lines peel out θ and σ^2 from the parameter vector, just to make it more readable. The next two lines calculate what the moments should be. The last line calculates the distance between the model predicted moments and the actual ones, and returns it. The whole thing could be turned into a one-line, like

```
return(t(params-c(c,v)) %*% (params-c(c,v)))
```

or perhaps even more obscure, but that is usually a bad idea.

Notice that I could write these two functions independently of one another, at least to some degree. When writing `ma.mm.est`, I knew I would need the objective function, but all I needed to know about it was its name, and the promise that it would take a parameter vector and give back a real number. When writing `ma.mm.objective`, all I had to remember about the other function was the promise this one needed to fulfill. In my experience, it is usually easiest to do any substantial coding in this “top-down” fashion²³. Start with the high-level goal you are trying to achieve, break it down into a few steps, write something which will put those steps together, presuming other functions or programs can do them. Now go and write the functions to do each of those steps.

The code for the method of simulated moments is entirely parallel to these. Writing it as two separate pairs of functions is therefore somewhat wasteful. If I find a mistake in one pair, or think of a way to improve it, I need to remember to make corresponding changes in the other pair (and not introduce a new mistake). In the long run, when you find yourself writing parallel pieces of code over and over, it is better to try to pull together the common parts and write them *once*. Here, that would mean something like one pair of functions, with the inner one having an argument

²²Here `par` is a very rough guess based on `c` and `v` — it’ll actually be right when `c=0`, but otherwise it’s not much good. Fortunately, it doesn’t have to be! Anyway, let’s return to designing the code

²³What qualifies as “substantial coding” depends on how much experience you have

which controlled whether to calculate the predicted moments by simulation or by a formula. You may try your hand at writing this.