
The Bootstrap

We are now several chapters into a statistics class and have said basically nothing about uncertainty. This should seem odd, and may even be disturbing if you are very attached to your p -values and saying variables have “significant effects”. It is time to remedy this, and talk about how we can quantify uncertainty for complex models. The key technique here is what’s called **bootstrapping**, or **the bootstrap**.

6.1 Stochastic Models, Uncertainty, Sampling Distributions

Statistics is the branch of mathematical engineering which studies ways of drawing inferences from limited and imperfect data. We want to know how a neuron in a rat’s brain responds when one of its whiskers gets tweaked, or how many rats live in Pittsburgh, or how high the water will get under the 16th Street bridge during May, or the typical course of daily temperatures in the city over the year, or the relationship between the number of birds of prey in Schenley Park in the spring and the number of rats the previous fall. We have some data on all of these things. But we know that our data is incomplete, and experience tells us that repeating our experiments or observations, even taking great care to replicate the conditions, gives more or less different answers every time. It is foolish to treat any inference from the data in hand as certain.

If all data sources were totally capricious, there’d be nothing to do beyond piously qualifying every conclusion with “but we could be wrong about this”. A mathematical discipline of statistics is possible because while repeating an experiment gives different results, some kinds of results are more common than others; their relative frequencies are reasonably stable. We thus model the data-generating mechanism through probability distributions and stochastic processes. When and why we can use stochastic models are very deep questions, but ones for another time. If we *can* use them in our problem, quantities like the ones I mentioned above are represented as functions of the stochastic model, i.e., of the underlying probability distribution. Since a function of a function is a “functional”, and these quantities are functions of the true probability distribution function, we’ll call these **functionals** or **statistical functionals**¹. Functionals could be single numbers (like the total rat population), or vectors, or even whole

¹ Most writers in theoretical statistics just call them “parameters” in a generalized sense, but I will

curves (like the expected time-course of temperature over the year, or the regression of hawks now on rats earlier). Statistical inference becomes estimating those functionals, or testing hypotheses about them.

These estimates and other inferences are functions of the data values, which means that they inherit variability from the underlying stochastic process. If we “re-ran the tape” (as the late, great Stephen Jay Gould used to say), we would get different data, with a certain characteristic distribution, and applying a fixed procedure would yield different inferences, again with a certain distribution. Statisticians want to use this distribution to quantify the uncertainty of the inferences. For instance, the standard error is an answer to the question “By how much would our estimate of this functional vary, typically, from one replication of the experiment to another?” (It presumes a particular meaning for “typically vary”, as the root-mean-square deviation around the mean.) A confidence region on a parameter, likewise, is the answer to “What are all the values of the parameter which *could* have produced this data with at least some specified probability?”, i.e., all the parameter values under which our data are not low-probability outliers. The confidence region is a promise that *either* the true parameter point lies in that region, *or* something very unlikely under any circumstances happened — or that our stochastic model is wrong.

To get things like standard errors or confidence intervals, we need to know the distribution of our estimates around the true values of our functionals. These **sampling distributions** follow, remember, from the distribution of the data, since our estimates are functions of the data. Mathematically the problem is well-defined, but actually *computing* anything is another story. Estimates are typically complicated functions of the data, and mathematically-convenient distributions may all be poor approximations to the data source. Saying anything in closed form about the distribution of estimates can be simply hopeless. The two classical responses of statisticians were to focus on tractable special cases, and to appeal to asymptotics.

Your introductory statistics courses mostly drilled you in the special cases. From one side, limit the kind of estimator we use to those with a simple mathematical form — say, means and other linear functions of the data. From the other, assume that the probability distributions featured in the stochastic model take one of a few forms for which exact calculation *is* possible, analytically or via tabulated special functions. Most such distributions have origin myths: the Gaussian arises from averaging many independent variables of equal size (say, the many genes which contribute to height in humans); the Poisson distribution comes from counting how many of a large number of independent and individually-improbable events have occurred (say, radioactive nuclei decaying in a given second), etc. Squeezed from both ends, the sampling distribution of estimators and other functions of the data becomes exactly calculable in terms of the aforementioned special functions.

try to restrict that word to actual parameters specifying statistical models, to minimize confusion. I may slip up.

That these origin myths invoke various limits is no accident. The great results of probability theory — the laws of large numbers, the ergodic theorem, the central limit theorem, etc. — describe limits in which *all* stochastic processes in broad classes of models display the same asymptotic behavior. The central limit theorem, for instance, says that if we average more and more independent random quantities with a common distribution, and that common distribution isn't too pathological, then the average becomes closer and closer to a Gaussian². Typically, as in the CLT, the limits involve taking more and more data from the source, so statisticians use the theorems to find the asymptotic, large-sample distributions of their estimates. We have been especially devoted to re-writing our estimates as averages of independent quantities, so that we can use the CLT to get Gaussian asymptotics.

Up through about the 1960s, statistics was split between developing general ideas about how to draw and evaluate inferences with stochastic models, and working out the properties of inferential procedures in tractable special cases (especially the linear-and-Gaussian case), or under asymptotic approximations. This yoked a very broad and abstract theory of inference to very narrow and concrete practical formulas, an uneasy combination often preserved in basic statistics classes.

The arrival of (comparatively) cheap and fast computers made it feasible for scientists and statisticians to record lots of data and to fit models to it, so they did. Sometimes the models were conventional ones, including the special-case assumptions, which often enough turned out to be detectably, and consequentially, wrong. At other times, scientists wanted more complicated or flexible models, some of which had been proposed long before, but now moved from being theoretical curiosities to stuff that could run overnight³. In principle, asymptotics might handle either kind of problem, but convergence to the limit could be unacceptably slow, especially for more complex models.

By the 1970s, then, statistics faced the problem of quantifying the uncertainty of inferences without using either implausibly-helpful assumptions or asymptotics; all of the solutions turned out to demand *even more* computation. Here we will examine what may be the most successful solution, Bradley Efron's proposal to combine estimation with simulation, which he gave the less-than-clear but persistent name of "the bootstrap" (Efron, 1979).

6.2 The Bootstrap Principle

Remember (from baby stats.) that the key to dealing with uncertainty in parameters and functionals is the sampling distribution of estimators. Knowing what distribution we'd get for our estimates on repeating the experiment would give us things like standard errors. Efron's insight was that we can *simulate* repli-

² The reason is that the non-Gaussian parts of the distribution wash away under averaging, but the average of two Gaussians is another Gaussian.

³ Kernel regression (§1.5.2), kernel density estimation (Ch. 14), and nearest-neighbors prediction (§1.5.1) were all proposed in the 1950s or 1960s, but didn't begin to be widely used until about 1980.




Figure 6.1 Schematic for model-based bootstrapping: simulated values are generated from the fitted model, then treated like the original data, yielding a new estimate of the functional of interest, here called $q_{0.01}$.

cation. After all, we have already fitted a model to the data, which is a guess at the mechanism which generated the data. Running that mechanism generates simulated data which, by hypothesis, has the same distribution as the real data. Feeding the simulated data through our estimator gives us one draw from the sampling distribution; repeating this many times yields the sampling distribution. Since we are using the model to give us its own uncertainty, Efron called this “bootstrapping”; unlike the Baron Munchhausen’s plan for getting himself out of a swamp by pulling on his own bootstraps, it works.

Figure 6.1 sketches the over-all process: fit a model to data, use the model to calculate the functional, then get the sampling distribution by generating new, synthetic data from the model and repeating the estimation on the simulation output.

To fix notation, we’ll say that the original data is x . (In general this is a whole data frame, not a single number.) Our parameter estimate from the data is $\hat{\theta}$. Surrogate data sets simulated from the fitted model will be $\tilde{X}_1, \tilde{X}_2, \dots, \tilde{X}_B$. The corresponding re-estimates of the parameters on the surrogate data are $\tilde{\theta}_1, \tilde{\theta}_2, \dots, \tilde{\theta}_B$.

The functional of interest is estimated by the statistic⁴ T , with sample value $\hat{t} = T(x)$, and values of the surrogates of $\tilde{t}_1 = T(\tilde{X}_1)$, $\tilde{t}_2 = T(\tilde{X}_2)$, \dots , $\tilde{t}_B = T(\tilde{X}_B)$. (The statistic T may be a direct function of the estimated parameters, and only indirectly a function of x .) Everything which follows applies without modification when the functional of interest is the parameter, or some component of the parameter.

In this section, we will assume that the model is correct for *some* value of θ , which we will call θ_0 . This means that we are employing a parametric **model-based bootstrap**. The true (population or ensemble) values of the functional is likewise t_0 .

6.2.1 Variances and Standard Errors

The simplest thing to do is to get the variance or standard error:

$$\widehat{\text{Var}}[\tilde{t}] = \mathbb{V}[\tilde{t}] \quad (6.1)$$

$$\widehat{\text{se}}(\hat{t}) = \text{sd}(\tilde{t}) \quad (6.2)$$

That is, we approximate the variance of our estimate of t_0 under the true but unknown distribution θ_0 by the variance of re-estimates \tilde{t} on surrogate data from the fitted model $\hat{\theta}$. Similarly we approximate the true standard error by the standard deviation of the re-estimates. The logic here is that the simulated \tilde{X} has about the same distribution as the real X that our data, x , was drawn from, so applying the same estimation procedure to the surrogate data gives us the sampling distribution. This assumes, of course, that our model is right, and that $\hat{\theta}$ is not too far from θ_0 .

A code sketch is provided in Code Example 7. Note that this may not work *exactly* as given in some circumstances, depending on the syntax details of, say, just what kind of data structure is needed to store \hat{t} .

6.2.2 Bias Correction

We can use bootstrapping to correct for a biased estimator. Since the sampling distribution of \tilde{t} is close to that of \hat{t} , and \hat{t} itself is close to t_0 ,

$$\mathbb{E}[\tilde{t}] - t_0 \approx \mathbb{E}[\tilde{t}] - \hat{t} \quad (6.3)$$

The left hand side is the bias that we want to know, and the right-hand side the was what we can calculate with the bootstrap.

In fact, Eq. 6.3 remains valid so long as the sampling distribution of $\hat{t} - t_0$ is close to that of $\tilde{t} - \hat{t}$. This is a weaker requirement than asking for \hat{t} and \tilde{t} themselves to have similar distributions, or asking for \hat{t} to be close to t_0 . In statistical theory, a random variable whose distribution does not depend on the parameters is called a **pivot**. (The metaphor is that it stays in one place while

⁴ T is a common symbol in the literature on the bootstrap for a *generic* function of the data. It may or may not have anything to do with Student's t test for difference in means.

```

rboot <- function(statistic, simulator, B) {
  tboots <- replicate(B, statistic(simulator()))
  if (is.null(dim(tboots))) {
    tboots <- array(tboots, dim = c(1, B))
  }
  return(tboots)
}
bootstrap <- function(tboots, summarizer, ...) {
  summaries <- apply(tboots, 1, summarizer, ...)
  return(t(summaries))
}
bootstrap.se <- function(statistic, simulator, B) {
  bootstrap(rboot(statistic, simulator, B), summarizer = sd)
}

```

CODE EXAMPLE 7: Code for calculating bootstrap standard errors. The function `rboot` generates B bootstrap samples (using the `simulator` function) and calculates the statistic on them (using `statistic`). `simulator` needs to be a function which returns a surrogate data set in a form suitable for `statistic`. (How would you modify the code to pass arguments to `simulator` and/or `statistic`?) Because every use of bootstrapping is going to need to do this, it makes sense to break it out as a separate function, rather than writing the same code many times (with many chances of getting it wrong). The `bootstrap` function takes the output of `rboot` and applies a summarizing function. `bootstrap.se` just calls `rboot` and makes the summarizing function `sd`, which takes a standard deviation. IMPORTANT NOTE: This is just a code sketch, because depending on the data structure which the statistic returns, it may not (e.g.) be feasible to just run `sd` on it, and so it might need some modification. See detailed examples below.

```

bootstrap.bias <- function(simulator, statistic, B, t.hat) {
  expect <- bootstrap(rboot(statistic, simulator, B), summarizer = mean)
  return(expect - t.hat)
}

```

CODE EXAMPLE 8: Sketch of code for bootstrap bias correction. Arguments are as in Code Example 7, except that `t.hat` is the estimate on the original data. IMPORTANT NOTE: As with Code Example 7, this is just a code sketch, because it won't work with all data types that might be returned by `statistic`, and so might require modification.

the parameters turn around it.) A sufficient (but not necessary) condition for Eq. 6.3 to hold is that $\hat{t} - t_0$ be a pivot, or approximately pivotal.

6.2.3 Confidence Intervals

A confidence interval is a random interval which contains the truth with high probability (the confidence level). If the confidence interval for g is C , and the confidence level is $1 - \alpha$, then we want

$$\Pr(t_0 \in C) = 1 - \alpha \quad (6.4)$$

no matter what the true value of t_0 . When we calculate a confidence interval, our inability to deal with distributions exactly means that the true confidence level, or **coverage** of the interval, is not quite the desired confidence level $1 - \alpha$; the

closer it is, the better the approximation, and the more accurate the confidence interval.⁵

When we simulate, we get samples of \tilde{t} , but what we really care about is the distribution of \hat{t} . When we have enough data to start with, those two distributions will be approximately the same. But at any given amount of data, the distribution of $\tilde{t} - \hat{t}$ will usually be closer to that of $\tilde{t} - t_0$ than the distribution of \tilde{t} is to that of \hat{t} . That is, the distribution of fluctuations around the true value usually converges quickly. (Think of the central limit theorem.) We can use this to turn information about the distribution of \tilde{t} into accurate confidence intervals for t_0 , essentially by re-centering \tilde{t} around \hat{t} .

Specifically, let $q_{\alpha/2}$ and $q_{1-\alpha/2}$ be the $\alpha/2$ and $1 - \alpha/2$ quantiles of \tilde{t} . Then

$$1 - \alpha = \Pr(q_{\alpha/2} \leq \tilde{T} \leq q_{1-\alpha/2}) \quad (6.5)$$

$$= \Pr(q_{\alpha/2} - \hat{T} \leq \tilde{T} - \hat{T} \leq q_{1-\alpha/2} - \hat{T}) \quad (6.6)$$

$$\approx \Pr(q_{\alpha/2} - \hat{T} \leq \hat{T} - t_0 \leq q_{1-\alpha/2} - \hat{T}) \quad (6.7)$$

$$= \Pr(q_{\alpha/2} - 2\hat{T} \leq -t_0 \leq q_{1-\alpha/2} - 2\hat{T}) \quad (6.8)$$

$$= \Pr(2\hat{T} - q_{1-\alpha/2} \leq t_0 \leq 2\hat{T} - q_{\alpha/2}) \quad (6.9)$$

The interval $C = [2\hat{T} - q_{\alpha/2}, 2\hat{T} - q_{1-\alpha/2}]$ is random, because \hat{T} is a random quantity, so it makes sense to talk about the probability that it contains the true value t_0 . Also, notice that the upper and lower quantiles of \tilde{T} have, as it were, swapped roles in determining the upper and lower confidence limits. Finally, notice that we do not actually know those quantiles exactly, but they're what we approximate by bootstrapping.

This is the **basic bootstrap confidence interval**, or the **pivotal CI**. It is simple and reasonably accurate, and makes a very good default choice for finding confidence intervals.

6.2.3.1 Other Bootstrap Confidence Intervals

The basic bootstrap CI relies on the distribution of $\tilde{t} - \hat{t}$ being approximately the same as that of $\hat{t} - t_0$. Even when this is false, however, it can be that the distribution of

$$\tau = \frac{\hat{t} - t_0}{\widehat{\text{se}}(\hat{t})} \quad (6.10)$$

is close to that of

$$\tilde{\tau} = \frac{\tilde{t} - \hat{t}}{\widehat{\text{se}}(\tilde{t})} \quad (6.11)$$

⁵ You might wonder why we'd be unhappy if the coverage level was *greater* than $1 - \alpha$. This is certainly better than if it's *less* than the nominal confidence level, but it usually means we could have used a smaller set, and so been more precise about t_0 , without any more real risk. Confidence intervals whose coverage is greater than the nominal level are called **conservative**; those with less than nominal coverage are **anti-conservative** (and not, say, "liberal").

```

equitails <- function(x, alpha) {
  lower <- quantile(x, alpha/2)
  upper <- quantile(x, 1 - alpha/2)
  return(c(lower, upper))
}
bootstrap.ci <- function(statistic = NULL, simulator = NULL, tboots = NULL,
  B = if (!is.null(tboots)) {
    ncol(tboots)
  }, t.hat, level) {
  if (is.null(tboots)) {
    stopifnot(!is.null(statistic))
    stopifnot(!is.null(simulator))
    stopifnot(!is.null(B))
    tboots <- rboot(statistic, simulator, B)
  }
  alpha <- 1 - level
  intervals <- bootstrap(tboots, summarizer = equitails, alpha = alpha)
  upper <- t.hat + (t.hat - intervals[, 1])
  lower <- t.hat + (t.hat - intervals[, 2])
  CIs <- cbind(lower = lower, upper = upper)
  return(CIs)
}

```

CODE EXAMPLE 9: Sketch of code for calculating the basic bootstrap confidence interval. See Code Example 7 for *rboot* and *bootstrap*, and cautions about blindly applying this to arbitrary data-types.

This is like what we calculate in a *t*-test, and since the *t*-test was invented by “Student”, these are called **studentized** quantities. If τ and $\tilde{\tau}$ have the same distribution, then we can reason as above and get a confidence interval

$$(\hat{t} - \widehat{\text{se}}(\hat{t})Q_{\tilde{\tau}}(1 - \alpha/2), \hat{t} - \widehat{\text{se}}(\hat{t})Q_{\tilde{\tau}}(\alpha/2)) \quad (6.12)$$

This is the same as the basic interval when $\widehat{\text{se}}(\hat{t}) = \text{se}(\tilde{t})$, but different otherwise. To find $\text{se}(\tilde{t})$, we need to actually do a *second* level of bootstrapping, as follows.

1. Fit the model with $\hat{\theta}$, find \hat{t} .
2. For $i \in 1 : B_1$
 1. Generate \tilde{X}_i from $\hat{\theta}$
 2. Estimate $\tilde{\theta}_i, \tilde{t}_i$
 3. For $j \in 1 : B_2$
 1. Generate X_{ij}^\dagger from $\tilde{\theta}_i$
 2. Calculate t_{ij}^\dagger
 4. Set $\tilde{\sigma}_i =$ standard deviation of the t_{ij}^\dagger
 5. Set $\tilde{\tau}_{ij} = \frac{t_{ij}^\dagger - \tilde{t}_i}{\tilde{\sigma}_i}$ for all j
3. Set $\widehat{\text{se}}(\hat{t}) =$ standard deviation of the \tilde{t}_i
4. Find the $\alpha/2$ and $1 - \alpha/2$ quantiles of the distribution of the $\tilde{\tau}$
5. Plug into Eq. 6.12.


```
boot.pvalue <- function(test, simulator, B, testthat) {
  testboot <- rboot(B = B, statistic = test, simulator = simulator)
  p <- (sum(testboot >= testthat) + 1)/(B + 1)
  return(p)
}
```

CODE EXAMPLE 10: *Bootstrap p-value calculation.* `testthat` should be the value of the test statistic on the actual data. `test` is a function which takes in a data set and calculates the test statistic, presuming that large values indicate departure from the null hypothesis. Note the +1 in the numerator and denominator of the p-value — it would be more straightforward to leave them off, but this is a little more stable when B is comparatively small. (Also, it keeps us from ever reporting a p-value of exactly 0.)

The advantage of the studentized intervals is that they are more accurate than the basic ones; the disadvantage is that they are more work! At the other extreme, the **percentile method** simply sets the confidence interval to

$$(Q_i(\alpha/2), Q_i(1 - \alpha/2)) \quad (6.13)$$

This is definitely easier to calculate, but not as accurate as the basic, pivotal CI.

All of these methods have many variations, described in the monographs referred to at the end of this chapter (§6.9).

6.2.4 Hypothesis Testing

For hypothesis tests, we may want to calculate two sets of sampling distributions: the distribution of the test statistic under the null tells us about the size of the test and significance levels, and the distribution under the alternative tells us about power and realized power. We can find either with bootstrapping, by simulating from either the null or the alternative. In such cases, the statistic of interest, which I've been calling T , is the test statistic. Code Example 10 illustrates how to find a p -value by simulating under the null hypothesis. The same procedure would work to calculate power, only we'd need to simulate from the alternative hypothesis, and `testthat` would be set to the critical value of T separating acceptance from rejection, not the observed value.

6.2.4.1 Double bootstrap hypothesis testing

When the hypothesis we are testing involves estimated parameters, we may need to correct for this. Suppose, for instance, that we are doing a goodness-of-fit test. If we estimate our parameters on the data set, we adjust our distribution so that it matches the data. It is thus not surprising if it seems to fit the data well! (Essentially, it's the problem of evaluating performance by looking at in-sample fit, which gave us so much trouble in Chapter 3.)

Some test statistics have distributions which are not affected by estimating parameters, at least not asymptotically. In other cases, one can analytically come up with correction terms. When these routes are blocked, one uses a **double bootstrap**, where a second level of bootstrapping checks how much estimation

```

doubleboot.pvalue <- function(test, simulator, B1, B2, estimator, thetahat,
  testthat, ...) {
  for (i in 1:B1) {
    xboot <- simulator(theta = thetahat, ...)
    thetaboot <- estimator(xboot)
    testboot[i] <- test(xboot)
    pboot[i] <- boot.pvalue(test, simulator, B2, testthat = testboot[i],
      theta = thetaboot)
  }
  p <- (sum(testboot >= testthat) + 1)/(B1 + 1)
  p.adj <- (sum(pboot <= p) + 1)/(B1 + 1)
  return(p.adj)
}

```

CODE EXAMPLE 11: Code sketch for “double bootstrap” significance testing. The inner or second bootstrap is used to calculate the distribution of nominal bootstrap p -values. For this to work, we need to draw our second-level bootstrap samples from $\tilde{\theta}$, the bootstrap re-estimate, not from $\hat{\theta}$, the data estimate. The code presumes the `simulator` function takes a `theta` argument allowing this. Exercise: replace the `for` loop with `replicate`.

improves the apparent fit of the model. This is perhaps most easily explained in pseudo-code (Code Example 11).

6.2.5 Model-Based Bootstrapping Example: Pareto’s Law of Wealth Inequality

The **Pareto** or **power-law** distribution⁶, is a popular model for data with “heavy tails”, i.e. where the probability density $f(x)$ goes to zero only very slowly as $x \rightarrow \infty$. The probability density is

$$f(x) = \frac{\theta - 1}{x_0} \left(\frac{x}{x_0} \right)^{-\theta} \quad (6.14)$$

where x_0 is the minimum scale of the distribution, and θ is the **scaling exponent** (Exercise 6.1). The Pareto is highly right-skewed, with the mean being much larger than the median.

If we know x_0 , one can show that the maximum likelihood estimator of the exponent θ is

$$\hat{\theta} = 1 + \frac{n}{\sum_{i=1}^n \log \frac{x_i}{x_0}} \quad (6.15)$$

and that this is consistent (Exercise 6.3), and efficient. Picking x_0 is a harder problem (see Clauset *et al.* 2009) — for the present purposes, pretend that the Oracle tells us. The file `pareto.R`, on the book website, contains a number of functions related to the Pareto distribution, including a function `pareto.fit` for estimating it. (There’s an example of its use below.)

Pareto came up with this density when he attempted to model the distribution

⁶ Named after Vilfredo Pareto (1848–1923), the highly influential economist, political scientist, and proto-Fascist.

```

sim.wealth <- function() {
  rpareto(n = n.tail, threshold = wealth.pareto$xmin, exponent = wealth.pareto$exponent)
}
est.pareto <- function(data) {
  pareto.fit(data, threshold = x0)$exponent
}

```

CODE EXAMPLE 12: *Simulator and estimator for model-based bootstrapping of the Pareto distribution.*

of personal wealth. Approximately, but quite robustly across countries and time-periods, the upper tail of the distribution of income and wealth follows a power law, with the exponent varying as money is more or less concentrated among the very richest individuals and households⁷. Figure 6.2 shows the distribution of net worth for the 400 richest Americans in 2003⁸.

```

source("http://www.stat.cmu.edu/~cshalizi/ADAfaEPoV/code/pareto.R")
wealth <- scan("http://www.stat.cmu.edu/~cshalizi/ADAfaEPoV/data/wealth.dat")
x0 <- 9e+08
n.tail <- sum(wealth >= x0)
wealth.pareto <- pareto.fit(wealth, threshold = x0)

```

Taking $x_0 = 9 \times 10^8$ (again, see Clauset *et al.* 2009), the number of individuals in the tail is 302, and the estimated exponent is $\hat{\theta} = 2.34$.

How much uncertainty is there in this estimate of the exponent? Naturally, we'll bootstrap. We need a function to generate Pareto-distributed random variables; this, along with some related functions, is part of the file `pareto.R` on the course website. With that tool, model-based bootstrapping proceeds as in Code Example 12.

Using these functions, we can now calculate the bootstrap standard error, bias and 95% confidence interval for $\hat{\theta}$, setting $B = 10^4$:

```

pareto.se <- bootstrap.se(statistic = est.pareto, simulator = sim.wealth, B = 10000)
pareto.bias <- bootstrap.bias(statistic = est.pareto, simulator = sim.wealth,
  t.hat = wealth.pareto$exponent, B = 10000)
pareto.ci <- bootstrap.ci(statistic = est.pareto, simulator = sim.wealth, B = 10000,
  t.hat = wealth.pareto$exponent, level = 0.95)


```

This gives a standard error of ± 0.078 , matching the asymptotic approximation reasonably well⁹, but not needing asymptotic assumptions.

⁷ Most of the distribution, for ordinary people, roughly conforms to a log-normal.

⁸ For the data source and a fuller analysis, see Clauset *et al.* (2009).

⁹ “In Asymptopia”, the variance of the MLE should be $\frac{(\theta-1)^2}{n}$, in this case 0.076. The intuition is that this variance depends on how sharp the maximum of the likelihood function is — if it's sharply peaked, we can find the maximum very precisely, but a broad maximum is hard to pin down. Variance is thus inversely proportional to the second derivative of the negative log-likelihood. (The minus sign is because the second derivative has to be negative at a maximum, while variance has to be positive.) For one sample, the expected second derivative of the negative log-likelihood is $(\theta - 1)^{-2}$. (This is called the **Fisher information** of the model.) Log-likelihood adds across



```
plot.survival.loglog(wealth, xlab = "Net worth (dollars)", ylab = "Fraction of top 400 above that we",
  rug(wealth, side = 1, col = "grey")
  curve((n.tail/400) * ppareto(x, threshold = x0, exponent = wealth.pareto$exponent,
    lower.tail = FALSE), add = TRUE, lty = "dashed", from = x0, to = 2 * max(wealth))
```

Figure 6.2 Upper cumulative distribution function (or “survival function”) of net worth for the 400 richest individuals in the US (2000 data). The solid line shows the fraction of the 400 individuals whose net worth W equaled or exceeded a given value w , $\Pr(W \geq w)$. (Note the logarithmic scale for both axes.) The dashed line is a maximum-likelihood estimate of the Pareto distribution, taking $x_0 = \$9 \times 10^8$. (This threshold was picked using the method of Clauset *et al.* 2009.) Since there are 302 individuals at or above the threshold, the cumulative distribution function of the Pareto has to be reduced by a factor of $(302/400)$.

```

ks.stat.pareto <- function(x, exponent, x0) {
  x <- x[x >= x0]
  ks <- ks.test(x, ppareto, exponent = exponent, threshold = x0)
  return(ks$statistic)
}
ks.pvalue.pareto <- function(B, x, exponent, x0) {
  testthat <- ks.stat.pareto(x, exponent, x0)
  testboot <- vector(length = B)
  for (i in 1:B) {
    xboot <- rpareto(length(x), exponent = exponent, threshold = x0)
    exp.boot <- pareto.fit(xboot, threshold = x0)$exponent
    testboot[i] <- ks.stat.pareto(xboot, exp.boot, x0)
  }
  p <- (sum(testboot >= testthat) + 1)/(B + 1)
  return(p)
}

```

CODE EXAMPLE 13: *Calculating a p-value for the Pareto distribution, using the Kolmogorov-Smirnov test and adjusting for the way estimating the scaling exponent moves the fitted distribution closer to the data.*

Asymptotically, the bias is known to go to zero; at this size, bootstrapping gives a bias of 0.0051, which is effectively negligible.

We can also get the confidence interval; with the same 10^4 replications, the 95% CI is 2.17, 2.48. In theory, the confidence interval could be calculated exactly, but it involves the inverse gamma distribution (Arnold, 1983), and it is quite literally faster to write and do the bootstrap than go to look it up.

A more challenging problem is goodness-of-fit; we'll use the Kolmogorov-Smirnov statistic.¹⁰ Code Example 13 calculates the p -value. With ten thousand bootstrap replications,

```

signif(ks.pvalue.pareto(10000, wealth, wealth.pareto$exponent, x0), 4)
## [1] 0.0101

```

Ten thousand replicates is enough that we should be able to accurately estimate probabilities of around 0.01 (since the binomial standard error will be $\sqrt{\frac{(0.01)(0.99)}{10^4}} \approx 9.9 \times 10^{-4}$); if it weren't, we might want to increase B .

Simply plugging in to the standard formulas, and thereby ignoring the effects of estimating the scaling exponent, gives a p -value of 0.171, which is not outstanding but not awful either. Properly accounting for the flexibility of the model, however, the discrepancy between what it predicts and what the data shows is so large that it would take a big (one-in-a-hundred) coincidence to produce it. We have,

independent samples, giving us an over-all factor of n . In the large-sample limit, the actual log-likelihood will converge on the expected log-likelihood, so this gives us the asymptotic variance. (See also §H.5.5.)

¹⁰ The `pareto.R` file contains a function, `pareto.tail.ks.test`, which does a goodness-of-fit test for fitting a power-law to the tail of the distribution. That differs somewhat from what follows, because it takes into account the extra uncertainty which comes from having to estimate x_0 . Here, I am pretending that an Oracle told us $x_0 = 9 \times 10^8$.

therefore, detected that the Pareto distribution makes systematic errors for this data, but we don't know much about what they are. In Chapter 15, we'll look at techniques which can begin to tell us something about *how* it fails.

6.3 Bootstrapping by Resampling

The bootstrap approximates the sampling distribution, with three sources of approximation error. First, **simulation error**: using finitely many replications to stand for the full sampling distribution. Clever simulation design can shrink this, but brute force — just using enough replicates — can also make it arbitrarily small. Second, **statistical error**: the sampling distribution of the bootstrap re-estimates under our estimated model is not exactly the same as the sampling distribution of estimates under the true data-generating process. The sampling distribution changes with the parameters, and our initial estimate is not completely accurate. But it often turns out that distribution of estimates *around* the truth is more nearly invariant than the distribution of estimates themselves, so subtracting the initial estimate from the bootstrapped values helps reduce the statistical error; there are many subtler tricks to the same end. Third, **specification error**: the data source doesn't exactly follow our model at all. Simulating the model then never quite matches the actual sampling distribution.

Efron had a second brilliant idea, which is to address specification error by replacing simulation from the model with re-sampling from the data. After all, our initial collection of data gives us a lot of information about the relative probabilities of different values. In a sense the empirical distribution is the least prejudiced estimate possible of the underlying distribution — anything else imposes biases or pre-conceptions, possibly accurate but also potentially misleading¹¹. Lots of quantities can be estimated directly from the empirical distribution, without the mediation of a model. Efron's **resampling bootstrap** (a.k.a. the **non-parametric bootstrap**) treats the original data set as a complete population and draws a new, simulated sample from it, picking each observation with equal probability (allowing repeated values) and then re-running the estimation (Figure 6.3, Code Example 14). In fact, this is usually what people mean when they talk about “the bootstrap” without any modifier.

Everything we did with model-based bootstrapping can also be done with re-sampling bootstrapping — the only thing that's changing is the distribution the surrogate data is coming from.

The resampling bootstrap should remind you of k -fold cross-validation. The analog of leave-one-out CV is a procedure called the **jack-knife**, where we repeat the estimate n times on $n - 1$ of the data points, holding each one out in turn. It's historically important (it dates back to the 1940s), but generally doesn't work as well as resampling.

An important variant is the **smoothed bootstrap**, where we re-sample the

¹¹ See §14.6 in Chapter 14.




Figure 6.3 Schematic for the resampling bootstrapping. New data is simulated by re-sampling from the original data (with replacement), and functionals are calculated either directly from the empirical distribution, or by estimating a model on this surrogate data.

```
resample <- function(x) {
  sample(x, size = length(x), replace = TRUE)
}
resample.data.frame <- function(data) {
  sample.rows <- resample(1:nrow(data))
  return(data[sample.rows, ])
}
```

CODE EXAMPLE 14: A utility function to resample from a vector, and another which resamples from a data frame. Can you write a single function which determines whether its argument is a vector or a data frame, and does the right thing in each case/

data points and then perturb each by a small amount of noise, generally Gaussian¹².

¹² We will see in Chapter 14 that this corresponds to sampling from a kernel density estimate.

Back to the Pareto example

Let's see how to use re-sampling to get a 95% confidence interval for the Pareto exponent¹³.

```
wealth.resample <- function() {
  resample(wealth[wealth >= x0])
}
pareto.CI.resamp <- bootstrap.ci(statistic = est.pareto, simulator = wealth.resample,
  t.hat = wealth.pareto$exponent, level = 0.95, B = 10000)
```

The interval is 2.17, 2.48; this is very close to the interval we got from the model-based bootstrap, which should actually reassure us about the latter's validity.

6.3.1 Model-Based vs. Resampling Bootstraps

When we have a properly specified model, simulating from the model gives more accurate results (at the same n) than does re-sampling the empirical distribution — parametric estimates of the distribution converge faster than the empirical distribution does. If on the other hand the model is mis-specified, then it is rapidly converging to the *wrong* distribution. This is of course just another bias-variance trade-off, like those we've seen in regression.

Since I am suspicious of most parametric modeling assumptions, I prefer re-sampling, when I can figure out how to do it, or at least until I have convinced myself that a parametric model is a good approximation to reality.

6.4 Bootstrapping Regression Models

Let's recap what we're doing estimating regression models. We want to learn the regression function $\mu(x) = \mathbb{E}[Y|X = x]$. We estimate the model on a set of predictor-response pairs, $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$, resulting in an estimated curve (or surface) $\hat{\mu}(x)$, fitted values $\hat{\mu}_i = \hat{\mu}(x_i)$, and residuals, $\epsilon_i = y_i - \hat{\mu}_i$. For any such model, we have a choice of several ways of bootstrapping, in decreasing order of reliance on the model.

- Simulate new X values from the model's distribution of X , and then draw Y from the specified conditional distribution $Y|X$.
- Hold the x fixed, but draw $Y|X$ from the specified distribution.
- Hold the x fixed, but make Y equal to $\hat{\mu}(x)$ plus a randomly re-sampled ϵ_j .
- Re-sample (x, y) pairs.

¹³ Even if the Pareto model is wrong, the estimator of the exponent will converge on the value which gives, in a certain sense, the best approximation to the true distribution from among all power laws. Econometricians call such parameter values the **pseudo-truth**; we are getting a confidence interval for the pseudo-truth. In this case, the pseudo-true scaling exponent can still be a useful way of summarizing *how* heavy tailed the income distribution is, despite the fact that the power law makes systematic errors.

The first case is pure model-based bootstrapping. (So is the second, sometimes, when the regression model is agnostic about X .) The last case is just re-sampling from the joint distribution of (X, Y) . The next-to-last case is called **re-sampling the residuals** or **re-sampling the errors**. When we do that, we rely on the regression model to get the conditional expectation function right, but we don't count on it getting the distribution of the noise around the expectations.

The specific procedure of re-sampling the residuals is to re-sample the ϵ_i , with replacement, to get $\tilde{\epsilon}_1, \tilde{\epsilon}_2, \dots, \tilde{\epsilon}_n$, and then set $\tilde{x}_i = x_i$, $\tilde{y}_i = \hat{\mu}(\tilde{x}_i) + \tilde{\epsilon}_i$. This surrogate data set is then re-analyzed like new data.

6.4.1 Re-sampling Points: Parametric Model Example

A classic data set contains the time between 299 eruptions of the Old Faithful geyser in Yellowstone, and the length of the subsequent eruptions; these variables are called **waiting** and **duration**. (We saw this data set already in §5.4.2.1, and will see it again in §10.3.2.) We'll look at the linear regression of **waiting** on **duration**. We'll re-sample (**duration**, **waiting**) pairs, and would like confidence intervals for the regression coefficients. This is a confidence interval for the coefficients of *the best linear predictor*, a functional of the distribution, which, as we saw in Chapters 1 and 2, exists no matter how nonlinear the process really is. It's only a confidence interval for the *true regression parameters* if the real regression function is linear.

Before anything else, look at the model:

```
library(MASS)
data(geyser)
geyser.lm <- lm(waiting ~ duration, data = geyser)
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	99.3	1.960	50.7	0
duration	-7.8	0.537	-14.5	0

The first step in bootstrapping this is to build our simulator, which just means sampling rows from the data frame:

```
resample.geyser <- function() {
  resample.data.frame(geyser)
}
```

We can check this by running `summary(geyser.resample())`, and seeing that it gives about the same quartiles and mean for both variables as `summary(geyser)`¹⁴, but that the former gives different numbers each time it's run.

Next, we define the estimator:

¹⁴ The minimum and maximum won't match up well — why not?

```
est.geyser.lm <- function(data) {
  fit <- lm(waiting ~ duration, data = data)
  return(coefficients(fit))
}
```

We can check that this function works by seeing that `coefficients(geyser.lm)` matches `est.geyser.lm(geyser)`, but that `est.geyser.lm(resample.geyser())` is different every time we run it.

Put the pieces together:

```
geyser.lm.ci <- bootstrap.ci(statistic=est.geyser.lm,
  simulator=resample.geyser,
  level=0.95,
  t.hat=coefficients(geyser.lm),
  B=1e4)
```

	lower	upper
(Intercept)	96.60	102.00
duration	-8.72	-6.95

Notice that we do not have to assume homoskedastic Gaussian noise — fortunately, because that’s a very bad assumption here¹⁵.

6.4.2 Re-sampling Points: Non-parametric Model Example

Nothing in the logic of re-sampling data points for regression requires us to use a parametric model. Here we’ll provide 95% confidence bounds for the kernel smoothing of the geyser data. Since the functional is a whole curve, the confidence set is often called a **confidence band**.

We use the same simulator, but start with a different regression curve, and need a different estimator.

```
evaluation.points <- data.frame(duration = seq(from = 0.8, to = 5.5, length.out = 200))
library(np)
npr.geyser <- function(data, tol = 0.1, ftol = 0.1, plot.df = evaluation.points) {
  bw <- npregbw(waiting ~ duration, data = data, tol = tol, ftol = ftol)
  mdl <- npreg(bw)
  return(predict(mdl, newdata = plot.df))
}
```

Now we construct pointwise 95% confidence bands for the regression curve.

¹⁵ We have calculated 95% confidence intervals for the intercept β_0 and the slope β_1 separately. These intervals cover their coefficients all but 5% of the time. Taken together, they give us a rectangle in (β_0, β_1) space, but the coverage probability of *this* rectangle could be anywhere from 95% all the way down to 90%. To get a confidence *region* which simultaneously covers both coefficients 95% of the time, we have two big options. One is to stick to a box-shaped region and just increase the confidence level on each coordinate (to 97.5%). The other is to define some suitable metric of how far apart coefficient vectors are (e.g., ordinary Euclidean distance), find the 95% percentile of the distribution of this metric, and trace the appropriate contour around $\hat{\beta}_0, \hat{\beta}_1$.

```

main.curve <- npr.geyser(geyser)

# We already defined this in a previous example, but it doesn't hurt
resample.geyser <- function() { resample.data.frame(geyser) }

geyser.resampled.curves <- rboot(statistic=npr.geyser,
                                simulator=resample.geyser,
                                B=800)

```

CODE EXAMPLE 15: *Generating multiple kernel-regression curves for the geyser data, by resampling that data frame and re-estimating the model on each simulation. `geyser.resampled.curves` stores the predictions of those 800 models, evaluated at a common set of values for the predictor variable. The vector `main.curve`, which we'll use presently to get confidence intervals, stores predictions of the model fit to the whole data, evaluated at that same set of points.*

For this end, we don't really need to keep around the whole kernel regression object — we'll just use its predicted values on a uniform grid of points, extending slightly beyond the range of the data (Code Example 15). Observe that this will go through bandwidth selection again for each bootstrap sample. This is slow, but it is the most secure way of getting good confidence bands. Applying the bandwidth we found on the data to each re-sample would be faster, but would introduce an extra level of approximation, since we wouldn't be treating each simulation run the same as the original data.

Figure 6.4 shows the curve fit to the data, the 95% confidence limits, and (faintly) all of the bootstrapped curves. Doing the 800 bootstrap replicates took 4 minutes on my laptop¹⁶.

6.4.3 Re-sampling Residuals: Example

As an example of re-sampling the residuals, rather than data points, let's take a linear regression, based on the data-analysis assignment in §A.14. We will regress `gdp.growth` on `log(gdp)`, `pop.growth`, `invest` and `trade`:


```

penn <- read.csv("http://www.stat.cmu.edu/~cshalizi/uADA/13/hw/02/penn-select.csv")
penn.formula <- "gdp.growth ~ log(gdp) + pop.growth + invest + trade"
penn.lm <- lm(penn.formula, data = penn)

```

(Why make the formula a separate object here?) The estimated parameters are

¹⁶ Specifically, I ran `system.time(geyser.resampled.curves <- rboot(statistic=npr.geyser, simulator=resample.geyser, B=800))`, which not only did the calculations and stored them in `geyser.resampled.curves`, but told me how much time it took R to do all that.



```
plot(0, type = "n", xlim = c(0.8, 5.5), ylim = c(0, 100), xlab = "Duration (min)",
     ylab = "Waiting (min)")
for (i in 1:ncol(geyser.resampled.curves)) {
  lines(evaluation.points$duration, geyser.resampled.curves[, i], lwd = 0.1,
        col = "grey")
}
geyser.npr.cis <- bootstrap.ci(tboots = geyser.resampled.curves, t.hat = main.curve,
                              level = 0.95)
lines(evaluation.points$duration, geyser.npr.cis[, "lower"])
lines(evaluation.points$duration, geyser.npr.cis[, "upper"])
lines(evaluation.points$duration, main.curve)
rug(geyser$duration, side = 1)
points(geyser$duration, geyser$waiting)
```

Figure 6.4 Kernel regression curve for Old Faithful (central black line), with 95% confidence bands (other black lines), the 800 bootstrapped curves (thin, grey lines), and the data points. Notice that the confidence bands get wider where there is less data. *Caution:* doing the bootstrap took 4 minutes to run on my computer.

```

resample.residuals.penn <- function() {
  new.frame <- penn
  new.growths <- fitted(penn.lm) + resample(residuals(penn.lm))
  new.frame$gdp.growth <- new.growths
  return(new.frame)
}
penn.estimator <- function(data) {
  mdl <- lm(penn.formula, data = data)
  return(coefficients(mdl))
}
penn.lm.cis <- bootstrap.ci(statistic = penn.estimator, simulator = resample.residuals.penn,
  B = 10000, t.hat = coefficients(penn.lm), level = 0.95)

```

CODE EXAMPLE 16: *Re-sampling the residuals to get confidence intervals in a linear model.*

	x
(Intercept)	5.71e-04
log(gdp)	5.07e-04
pop.growth	-1.87e-01
invest	7.15e-04
trade	3.11e-05

Code Example 16 shows the new simulator for this set-up (`resample.residuals.penn`)¹⁷, the new estimation function (`penn.estimator`)¹⁸, and the confidence interval calculation (`penn.lm.cis`):

	lower	upper
(Intercept)	-1.62e-02	1.71e-02
log(gdp)	-1.46e-03	2.49e-03
pop.growth	-3.58e-01	-1.75e-02
invest	4.94e-04	9.37e-04
trade	-1.94e-05	8.21e-05

Doing ten thousand linear regressions took 45 seconds on my computer, as opposed to 4 minutes for eight hundred kernel regressions.

6.5 Bootstrap with Dependent Data

If the data points we are looking at are vectors (or more complicated structures) with dependence between components, but each data point is independently generated from the same distribution, then dependence isn't really an issue. We

¹⁷ How would you check that this worked?

¹⁸ How would you check that this worked?

re-sample vectors, or generate vectors from our model, and proceed as usual. In fact, that's what we've done so far in several cases.

If there is dependence *across* data points, things are more tricky. If our model incorporates this dependence, then we can just simulate whole data sets from it. An appropriate re-sampling method is trickier — just re-sampling individual data points destroys the dependence, so it won't do. We will revisit this question when we look at time series in Chapter 25.

6.6 Confidence Bands for Nonparametric Regression

Many of the examples in this chapter use bootstrapping to get confidence bands for nonparametric regression. It is worth mentioning that there is a subtle issue with doing so, but one which I do not think really matters, usually, for practice.

The issue is that when we do nonparametric regression, we accept some bias in our estimate of the regression function. In fact, we saw in Chapter 4 that minimizing the total MSE means accepting *matching* amounts of bias and variance. So our nonparametric estimate of μ is biased. If we simulate from it, we're simulating from something biased; if we simulate from the residuals, those residuals contain bias; and even if we do a pure resampling bootstrap, we're comparing the bootstrap replicates to a biased estimate. This means that we are really looking at sampling intervals around the biased estimate, rather than confidence intervals around μ .

The two questions this raises are (1) how much this matters, and (2) whether there is any alternative. As for the size of the bias, we know from Chapter 4 that the squared bias, in 1D, goes like $n^{-4/5}$, so the bias itself goes like $n^{-2/5}$. This does go to zero, but slowly.

[[Living with it vs. Hall and Horowitz (2013) paper, which gives $1 - \alpha$ coverage at $1 - \eta$ fraction of points. Essentially, construct naive bands, and then work out by how much they need to be expanded to achieve desired coverage]]

6.7 Things Bootstrapping Does Poorly

The principle behind bootstrapping is that sampling distributions under the true process should be close to sampling distributions under good estimates of the truth. If small perturbations to the data-generating process produce huge swings in the sampling distribution, bootstrapping will not work well, and may fail spectacularly. For model-based bootstrapping, this means that small changes to the underlying parameters must produce small changes to the functionals of interest. Similarly, for resampling, it means that adding or removing a few data points must change the functionals only a little¹⁹.

Re-sampling in particular has trouble with extreme values. Here is a simple

¹⁹ More generally, moving from one distribution function f to another $(1 - \epsilon)f + \epsilon g$ mustn't change the functional very much when ϵ is small, no matter in what "direction" g we perturb it. Making this idea precise calls for some fairly deep mathematics, about differential calculus on spaces of functions (see, e.g., van der Vaart 1998, ch. 20).

example: Our data points X_i are IID, with $X_i \sim \text{Unif}(0, \theta_0)$, and we want to estimate θ_0 . The maximum likelihood estimate $\hat{\theta}$ is just the sample maximum of the x_i . We'll use resampling to get a confidence interval for this, as above — but I will fix the true $\theta_0 = 1$, and see how often the 95% confidence interval covers the truth.

```
max.boot.ci <- function(x, B) {
  max.boot <- replicate(B, max(resample(x)))
  return(2 * max(x) - quantile(max.boot, c(0.975, 0.025)))
}
boot.cis <- replicate(1000, max.boot.ci(x = runif(100), B = 1000))
(true.coverage <- mean((1 >= boot.cis[, 1] & (1 <= boot.cis[, 2])))
## [1] 0.87
```

That is, the actual coverage probability is not 95% but about 87%.

If you suspect that your use of the bootstrap may be setting yourself up for a similar epic fail, your two options are (1) learn some of the theory of the bootstrap from the references in the “Further Reading” section below, or (2) set up a simulation experiment like this one.

6.8 Which Bootstrap When?

This chapter has introduced a bunch of different bootstraps, and before it closes it's worth reviewing the general principles, and some of the considerations which go into choosing among them in a particular problem.

When we bootstrap, we try to approximate the sampling distribution of some statistic (mean, median, correlation coefficient, regression coefficients, smoothing curve, difference in MSEs . . .) by running simulations, and calculating the statistic on the simulation. We've seen three major ways of doing this:

- The model-based bootstrap: we estimate the model, and then simulate from x the estimated model;
- Resampling residuals: we estimate the model, and then simulate by resampling residuals to that estimate and adding them back to the fitted values;
- Resampling cases or whole data points: we ignore the estimated model completely in our simulation, and just re-sample whole rows from the data frame.

Which kind of bootstrap is appropriate depends on how much trust we have in our model.

The model-based bootstrap trusts the model to be completely correct for *some* parameter value. In, e.g., regression, it trusts that we have the right shape for the regression function *and* that we have the right distribution for the noise. When we trust our model this much, we could in principle work out sampling distributions analytically; the model-based bootstrap replaces hard math with simulation.

Resampling residuals doesn't trust the model as much. In regression problems, it assumes that the model gets the *shape* of the regression function right, and that the noise around the regression function is independent of the predictor

variables, but doesn't make any further assumption about how the fluctuations are distributed. It is therefore more secure than model-based bootstrap.²⁰

Finally, resampling cases assumes nothing at all about either the shape of the regression function or the distribution of the noise, it just assumes that each data point (row in the data frame) is an independent observation. Because it assumes so little, and doesn't depend on any particular model being correct, it is very safe.

The reason we do not always use the safest bootstrap, which is resampling cases, is that there is, as usual, a bias-variance trade-off. Generally speaking, if we compare three sets of bootstrap confidence intervals on the same data for the same statistic, the model-based bootstrap will give the narrowest intervals, followed by resampling residuals, and resampling cases will give the loosest bounds. If the model really *is* correct about the shape of the curve, we can get more precise results, without any loss of accuracy, by resampling residuals rather than resampling cases. If the model is also correct about the distribution of noise, we can do even better with a model-based bootstrap.

To sum up: resampling cases is safer than resampling residuals, but gives wider, weaker bounds. If you have good reason to trust a model's guess at the shape of the regression function, then resampling residuals is preferable. If you don't, or it's not a regression problem so there are no residuals, then you prefer to resample cases. The model-based bootstrap works best when the over-all model is correct, and we're just uncertain about the exact parameter values we need.

6.9 Further Reading

Davison and Hinkley (1997) is both a good textbook, and the reference I consult most often. Efron and Tibshirani (1993), while also very good, is more theoretical. Canty *et al.* (2006) has useful advice for serious applications.

All the bootstraps discussed in this chapter presume IID observations. For bootstraps for time series, see §25.5.

Software

For professional purposes, I strongly recommend using the R package `boot` (Canty and Ripley, 2013), based on Davison and Hinkley (1997). I deliberately do *not* use it in this chapter, or later in the book, for pedagogical reasons; I have found that forcing students to write their own bootstrapping code helps build character, or at least understanding.

The bootstrap vs. robust standard errors

For linear regression coefficients, econometricians have developed a variety of “robust” standard errors which are valid under weaker conditions than the usual

²⁰ You could also imagine simulations where we presume that the noise takes a very particular form (e.g., a t -distribution with 10 degrees of freedom), but are agnostic about the shape of the regression function, and learn that non-parametrically. It's harder to think of situations where this is really plausible, however, except *maybe* Gaussian noise arising from central-limit-theorem considerations.

assumptions. Buja *et al.* (2014) shows their equivalence to resampling cases. (See also King and Roberts 2015.)

Historical notes

The original paper on the bootstrap, Efron (1979), is extremely clear, and for the most part presented in the simplest possible terms; it's worth reading. His later small book (Efron, 1982), while often cited, is not in my opinion so useful nowadays²¹.

As the title of that last reference suggests, the bootstrap is in some ways a successor to an older method, apparently dating back to the 1940s if not before, called the “jackknife”, in which each data point is successively held back and the estimate is re-calculated; the variance of these re-estimates, appropriately scaled, is then taken as the variance of estimation, and similarly for the bias²². The jackknife is appealing in its simplicity, but is only valid under much stronger conditions than the bootstrap.

Exercises

- 6.1 Show that x_0 is the mode of the Pareto distribution.
- 6.2 Derive the maximum likelihood estimator for the Pareto distribution (Eq. 6.15) from the density (Eq. 6.14).
- 6.3 Show that the MLE of the Pareto distribution is consistent.
 1. Using the law of large numbers, show that $\hat{\theta}$ (Eq. 6.15) converges to a limit which depends on $\mathbb{E}[\log X/x_0]$.
 2. Find an expression for $\mathbb{E}[\log X/x_0]$ in terms of θ and from the density (Eq. 6.14).
Hint: Write $\mathbb{E}[\log X/x_0]$ as an integral, change the variable of integration from x to $z = \log(x/x_0)$, and remember that the mean of an exponential random variable with rate λ is $1/\lambda$.
- 6.4 Find confidence bands for the linear regression model of §6.4.1 using
 1. The usual Gaussian assumptions (*hint:* try the `intervals="confidence"` option to `predict`);
 2. Resampling of residuals; and
 3. Resampling of cases.
- 6.5 (Computational) Writing new functions to simulate every particular linear model is somewhat tedious.
 1. Write a function which takes, as inputs, an `lm` model and a data frame, and returns a new data frame where the response variable is replaced by the model's predictions plus Gaussian noise, but all other columns are left alone.
 2. Write a function which takes, as inputs, an `lm` model and a data frame, and returns a new data frame where the response variable is replaced by the model's predictions plus resampled residuals.

²¹ It seems to have done a good job of explaining things to people who were already professional statisticians in 1982.

²² A “jackknife” is a knife with a blade which folds into the handle; think of the held-back data point as the folded-away blade.

3. Will your functions work with `npreg` models, as well as `lm` models? If not, what do you have to modify?

Hint: See Code Example 3 in Chapter 3 for some R tricks to extract the name of the response variable from the estimated model.