

Convexity II: Optimization Basics

Lecturer: Ryan Tibshirani
Convex Optimization 10-725/36-725




See supplements for reviews of

- *basic multivariate calculus*
- *basic linear algebra*

Last time: convex sets and functions

“Convex calculus” makes it easy to check convexity. Tools:

- Definitions of **convex sets and functions**, classic examples



- Key properties (e.g., first- and second-order characterizations for functions)
- Operations that preserve convexity (e.g., affine composition)

E.g., is $\max \left\{ \log \left(\frac{1}{(a^T x + b)^7} \right), \|Ax + b\|_1^5 \right\}$ convex?

Outline

Today:

- Optimization terminology
- Properties and first-order optimality
- Equivalent transformations

Optimization terminology

Reminder: a convex optimization problem (or **program**) is

$$\begin{array}{ll} \min_{x \in D} & f(x) \\ \text{subject to} & g_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{array}$$

where f and g_i , $i = 1, \dots, m$ are all convex, and the optimization domain is $D = \text{dom}(f) \cap \bigcap_{i=1}^m \text{dom}(g_i)$ (often we do not write D)

- f is called **criterion** or **objective** function
- g_i is called **inequality constraint** function
- If $x \in D$, $g_i(x) \leq 0$, $i = 1, \dots, m$, and $Ax = b$ then x is called a **feasible point**
- The minimum of $f(x)$ over all feasible points x is called the **optimal value**, written f^*

- If x is feasible and $f(x) = f^*$, then x is called **optimal**; also called a **solution**, or a **minimizer**¹
- If x is feasible and $f(x) \leq f^* + \epsilon$, then x is called **ϵ -suboptimal**
- If x is feasible and $g_i(x) = 0$, then we say g_i is **active** at x
- Convex minimization can be reposed as concave maximization

$$\begin{array}{ll}
 \min_x & f(x) \\
 \text{subject to} & g_i(x) \leq 0, \\
 & i = 1, \dots, m \\
 & Ax = b
 \end{array}
 \iff
 \begin{array}{ll}
 \max_x & -f(x) \\
 \text{subject to} & g_i(x) \leq 0, \\
 & i = 1, \dots, m \\
 & Ax = b
 \end{array}$$

Both are called convex optimization problems

¹Note: a convex optimization problem need not have solutions, i.e., need not attain its minimum, but we will not be careful about this

Convex solution sets

Let X_{opt} be the set of all solutions of convex problem, written

$$\begin{aligned} X_{\text{opt}} = \operatorname{argmin} \quad & f(x) \\ \text{subject to} \quad & g_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{aligned}$$

Key property: X_{opt} is a **convex set**

Proof: use definitions. If x, y are solutions, then for $0 \leq t \leq 1$,

- $tx + (1 - t)y \in D$
- $g_i(tx + (1 - t)y) \leq tg_i(x) + (1 - t)g_i(y) \leq 0$
- $A(tx + (1 - t)y) = tAx + (1 - t)Ay = b$
- $f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y) = f^*$

Therefore $tx + (1 - t)y$ is also a solution

Another key property: if f is strictly convex, then the **solution is unique**, i.e., X_{opt} contains one element

Example: lasso

Given $y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n \times p}$, consider the **lasso** problem:

$$\begin{aligned} \min_{\beta} \quad & \|y - X\beta\|_2^2 \\ \text{subject to} \quad & \|\beta\|_1 \leq s \end{aligned}$$

Is this convex? What is the criterion function? The inequality and equality constraints? Feasible set? Is the solution unique, when:

- $n \geq p$ and X has full column rank?
- $p > n$ (“high-dimensional” case)?

How do our answers change if we changed criterion to **Huber loss**:

$$\sum_{i=1}^n \rho(y_i - x_i^T \beta), \quad \rho(z) = \begin{cases} \frac{1}{2}z^2 & |z| \leq \delta \\ \delta|z| - \frac{1}{2}\delta^2 & \text{else} \end{cases} \quad ?$$

Example: support vector machines

Given $y \in \{-1, 1\}^n$, $X \in \mathbb{R}^{n \times p}$ with rows x_1, \dots, x_n , consider the **support vector machine** or SVM problem:

$$\begin{aligned} \min_{\beta, \beta_0, \xi} \quad & \frac{1}{2} \|\beta\|_2^2 + C \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & \xi_i \geq 0, \quad i = 1, \dots, n \\ & y_i(x_i^T \beta + \beta_0) \geq 1 - \xi_i, \quad i = 1, \dots, n \end{aligned}$$

Is this convex? What is the criterion, constraints, feasible set? Is the solution (β, β_0, ξ) unique? What if changed the criterion to

$$\frac{1}{2} \|\beta\|_2^2 + \frac{1}{2} \beta_0^2 + C \sum_{i=1}^n \xi_i^{1.01}?$$

For original criterion, what about β component, at the solution?


Local minima are global minima

For a convex problem, a feasible point x is called **locally optimal** if there is some $R > 0$ such that


$$f(x) \leq f(y) \quad \text{for all feasible } y \text{ such that } \|x - y\|_2 \leq R$$

Reminder: for convex optimization problems, **local optima are global optima**

Proof simply follows
from definitions



Convex



Nonconvex

Rewriting constraints

The optimization problem

$$\begin{aligned} \min_x \quad & f(x) \\ \text{subject to} \quad & g_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{aligned}$$

can be rewritten as

$$\min_x f(x) \quad \text{subject to} \quad x \in C$$

where $C = \{x : g_i(x) \leq 0, i = 1, \dots, m, Ax = b\}$, the feasible set.
Hence the above formulation is **completely general**

With I_C the indicator of C , we can write this in unconstrained form

$$\min_x f(x) + I_C(x)$$


First-order optimality condition

For a convex problem

$$\min_x f(x) \quad \text{subject to } x \in C$$

and differentiable f , a feasible point x is optimal if and only if

$$\nabla f(x)^T (y - x) \geq 0 \quad \text{for all } y \in C$$



This is called the **first-order condition for optimality**

In words: all feasible directions from x are aligned with gradient $\nabla f(x)$

Important special case: if $C = \mathbb{R}^n$ (unconstrained optimization), then optimality condition reduces to familiar $\nabla f(x) = 0$

Example: quadratic minimization

Consider minimizing the **quadratic function**

$$f(x) = \frac{1}{2}x^T Qx + b^T x + c$$

where $Q \succeq 0$. The first-order condition says that solution satisfies

$$\nabla f(x) = Qx + b = 0$$

Cases:

- if $Q \succ 0$, then there is a unique solution $x = -Q^{-1}b$
- if Q is singular and $b \notin \text{col}(Q)$, then there is no solution (i.e., $\min_x f(x) = -\infty$)
- if Q is singular and $b \in \text{col}(Q)$, then there are infinitely many solutions

$$x = -Q^+b + z, \quad z \in \text{null}(Q)$$

where Q^+ is the **pseudoinverse** of Q

Example: equality-constrained minimization

Consider the equality-constrained convex problem:

$$\min_x f(x) \quad \text{subject to} \quad Ax = b$$

with f differentiable. Let's prove **Lagrange multiplier** optimality condition

$$\nabla f(x) + A^T u = 0 \quad \text{for some } u$$

According to first-order optimality, solution x satisfies $Ax = b$ and

$$\nabla f(x)^T (y - x) \geq 0 \quad \text{for all } y \text{ such that } Ay = b$$

This is equivalent to

$$\nabla f(x)^T v = 0 \quad \text{for all } v \in \text{null}(A)$$

Result follows because $\text{null}(A)^\perp = \text{row}(A)$

Example: projection onto a convex set

Consider **projection onto convex set** C :

$$\min_x \|a - x\|_2^2 \quad \text{subject to } x \in C$$


First-order optimality condition says that the solution x satisfies

$$\nabla f(x)^T (y - x) = (x - a)^T (y - x) \geq 0 \quad \text{for all } y \in C$$

Equivalently, this says that

$$a - x \in \mathcal{N}_C(x)$$

where recall $\mathcal{N}_C(x)$ is the normal cone to C at x



Partial optimization

Reminder: $g(x) = \min_{y \in C} f(x, y)$ is convex in x , provided that f is convex in (x, y) and C is a convex set

Therefore we can always **partially optimize** a convex problem and retain convexity

E.g., if we decompose $x = (x_1, x_2) \in \mathbb{R}^{n_1+n_2}$, then

$$\begin{array}{ll} \min_{x_1, x_2} & f(x_1, x_2) \\ \text{subject to} & g_1(x_1) \leq 0 \\ & g_2(x_2) \leq 0 \end{array} \iff \begin{array}{ll} \min_{x_1} & \tilde{f}(x_1) \\ \text{subject to} & g_1(x_1) \leq 0 \end{array}$$

where $\tilde{f}(x_1) = \min\{f(x_1, x_2) : g_2(x_2) \leq 0\}$. The right problem is convex if the left problem is

Example: hinge form of SVMs

Recall the SVM problem

$$\begin{aligned} \min_{\beta, \beta_0, \xi} \quad & \frac{1}{2} \|\beta\|_2^2 + C \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & \xi_i \geq 0, y_i(x_i^T \beta + \beta_0) \geq 1 - \xi_i, i = 1, \dots, n \end{aligned}$$

Rewrite the constraints as $\xi_i \geq \max\{0, 1 - y_i(x_i^T \beta + \beta_0)\}$. Indeed we can argue that we have = at solution

Therefore plugging in for optimal ξ gives the **hinge form** of SVMs:

$$\min_{\beta, \beta_0} \frac{1}{2} \|\beta\|_2^2 + C \sum_{i=1}^n [1 - y_i(x_i^T \beta + \beta_0)]_+$$

where $a_+ = \max\{0, a\}$ is called the hinge function

Transformations and change of variables

If $h : \mathbb{R} \rightarrow \mathbb{R}$ is a **monotone increasing transformation**, then

$$\begin{aligned} & \min_x f(x) \quad \text{subject to } x \in C \\ \iff & \min_x h(f(x)) \quad \text{subject to } x \in C \end{aligned}$$

Similarly, inequality or equality constraints can be transformed and yield equivalent optimization problems. Can use this to reveal the “hidden convexity” of a problem

If $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, and its image covers feasible set C , then we can **change variables** in an optimization problem:

$$\begin{aligned} & \min_x f(x) \quad \text{subject to } x \in C \\ \iff & \min_y f(\phi(y)) \quad \text{subject to } \phi(y) \in C \end{aligned}$$

Example: geometric programming

A **monomial** is a function $f : \mathbb{R}_{++}^n \rightarrow \mathbb{R}$ of the form

$$f(x) = \gamma x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$$

for $\gamma > 0$, $a_1, \dots, a_n \in \mathbb{R}$. A **posynomial** is a sum of monomials,

$$f(x) = \sum_{k=1}^p \gamma_k x_1^{a_{k1}} x_2^{a_{k2}} \cdots x_n^{a_{kn}}$$

A **geometric program** is of the form

$$\begin{aligned} \min_x \quad & f(x) \\ \text{subject to} \quad & g_i(x) \leq 1, \quad i = 1, \dots, m \\ & h_j(x) = 1, \quad j = 1, \dots, r \end{aligned}$$

where f , g_i , $i = 1, \dots, m$ are posynomials and h_j , $j = 1, \dots, r$ are monomials. This is nonconvex

Let's prove that a geometric program is equivalent to a convex one. Given $f(x) = \gamma x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$, let $y_i = \log x_i$ and rewrite this as


$$\gamma (e^{y_1})^{a_1} (e^{y_2})^{a_2} \cdots (e^{y_n})^{a_n} = e^{a^T y + b}$$

for $b = \log \gamma$. Also, a posynomial can be written as $\sum_{k=1}^p e^{a_k^T y + b_k}$. With this variable substitution, and after taking logs, a geometric program is equivalent to

$$\begin{aligned} \min_x \quad & \log \left(\sum_{k=1}^{p_0} e^{a_{0k}^T y + b_{0k}} \right) \\ \text{subject to} \quad & \log \left(\sum_{k=1}^{p_i} e^{a_{ik}^T y + b_{ik}} \right) \leq 0, \quad i = 1, \dots, m \\ & c_j^T y + d_j = 0, \quad j = 1, \dots, r \end{aligned}$$

This is convex, recalling the convexity of soft max functions

Many interesting problems are geometric programs, e.g., floor planning:



See Boyd et al. (2007), “A tutorial on geometric programming”, and also Chapter 8.8 of B & V book

Example floor planning program:

$$\begin{aligned} & \min_{\substack{W, H, \\ x, y, w, h}} && WH \\ & \text{subject to} && 0 \leq x_i \leq W, \quad i = 1, \dots, n \\ & && 0 \leq y_i \leq H, \quad i = 1, \dots, n \\ & && x_i + w_i \leq x_j, \quad (i, j) \in \mathcal{L} \\ & && y_i + h_i \leq y_j, \quad (i, j) \in \mathcal{B} \\ & && w_i h_i = C_i, \quad i = 1, \dots, n. \end{aligned}$$

Check: why is this a geometric program?

Eliminating equality constraints

Important special case of change of variables: **eliminating equality constraints**. Given the problem

$$\begin{aligned} \min_x \quad & f(x) \\ \text{subject to} \quad & g_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{aligned}$$

we can always express any feasible point as $x = My + x_0$, where $Ax_0 = b$ and $\text{col}(M) = \text{null}(A)$. Hence the above is equivalent to

$$\begin{aligned} \min_y \quad & f(My + x_0) \\ \text{subject to} \quad & g_i(My + x_0) \leq 0, \quad i = 1, \dots, m \end{aligned}$$

Note: this is fully general but not always a good idea (practically)

Introducing slack variables

Essentially opposite to eliminating equality constraints: **introducing slack variables**. Given the problem

$$\begin{aligned} \min_x \quad & f(x) \\ \text{subject to} \quad & g_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b \end{aligned}$$

we can transform the inequality constraints via

$$\begin{aligned} \min_{x,s} \quad & f(x) \\ \text{subject to} \quad & s_i \geq 0, \quad i = 1, \dots, m \\ & g_i(x) + s_i = 0, \quad i = 1, \dots, m \\ & Ax = b \end{aligned}$$

Note: this is no longer convex unless $g_i, i = 1, \dots, m$ are affine

Relaxing nonaffine equality constraints

Given an optimization problem

$$\min_x f(x) \quad \text{subject to } x \in C$$

we can always take an enlarged constraint set $\tilde{C} \supseteq C$ and consider

$$\min_x f(x) \quad \text{subject to } x \in \tilde{C}$$

This is called a **relaxation** and its optimal value is always smaller or equal to that of the original problem

Important special case: **relaxing nonaffine equality constraints**, i.e.,

$$h_j(x) = 0, \quad j = 1, \dots, r$$

where $h_j, j = 1, \dots, r$ are convex but nonaffine, are replaced with

$$h_j(x) \leq 0, \quad j = 1, \dots, r$$

Example: maximum utility problem

The **maximum utility problem** models investment/consumption:

$$\begin{aligned} \max_{x,b} \quad & \sum_{t=1}^T \alpha_t u(x_t) \\ \text{subject to} \quad & b_{t+1} = b_t + f(b_t) - x_t, \quad t = 1, \dots, T \\ & 0 \leq x_t \leq b_t, \quad t = 1, \dots, T \end{aligned}$$

Here b_t is the budget and x_t is the amount consumed at time t ; f is an investment return function, u utility function, both concave and increasing

Is this a convex problem? What if we replace equality constraints with inequalities:

$$b_{t+1} \leq b_t + f(b_t) - x_t, \quad t = 1, \dots, T?$$

Example: principal components analysis

Given $X \in \mathbb{R}^{n \times p}$, consider the low rank approximation problem:

$$\min_R \|X - R\|_F^2 \quad \text{subject to} \quad \text{rank}(R) = k$$

Here $\|A\|_F^2 = \sum_{i=1}^n \sum_{j=1}^p A_{ij}^2$, the entrywise squared ℓ_2 norm, and $\text{rank}(A)$ denotes the rank of A

Also called principal components analysis or PCA problem. Given $X = UDV^T$, singular value decomposition or SVD, the solution is

$$R = U_k D_k V_k^T$$

where U_k, V_k are the first k columns of U, V and D_k is the first k diagonal elements of D . I.e., R is reconstruction of X from its **first k principal components**

This problem is not convex. Why?

We can recast the PCA problem in a convex form. First rewrite as

$$\begin{aligned} \min_{Z \in \mathbb{S}^p} \|X - XZ\|_F^2 \quad \text{subject to } \text{rank}(Z) = k, \quad Z \text{ is a projection} \\ \iff \max_{Z \in \mathbb{S}^p} \text{tr}(SZ) \quad \text{subject to } \text{rank}(Z) = k, \quad Z \text{ is a projection} \end{aligned}$$

where $S = X^T X$. Hence constraint set is the nonconvex set

$$C = \left\{ Z \in \mathbb{S}^p : \lambda_i(Z) \in \{0, 1\}, \quad i = 1, \dots, p, \quad \text{tr}(Z) = k \right\}$$

where $\lambda_i(Z)$, $i = 1, \dots, n$ are the eigenvalues of Z . Solution in this formulation is

$$Z = V_k V_k^T$$

where V_k gives first k columns of V

Now consider relaxing constraint set to $\mathcal{F}_k = \text{conv}(C)$, its convex hull. Note

$$\begin{aligned}\mathcal{F}_k &= \{Z \in \mathbb{S}^p : \lambda_i(Z) \in [0, 1], i = 1, \dots, p, \text{tr}(Z) = k\} \\ &= \{Z \in \mathbb{S}^p : 0 \preceq Z \preceq I, \text{tr}(Z) = k\}\end{aligned}$$

Recall this is called the **Fantope** of order k

Hence, the linear maximization over the Fantope, namely

$$\max_{Z \in \mathcal{F}_k} \text{tr}(SZ)$$

is convex. Remarkably, this is equivalent to the nonconvex PCA problem (admits the same solution)!

(Famous result: Fan (1949), "On a theorem of Weyl concerning eigenvalues of linear transformations")

References and further reading

- S. Boyd and L. Vandenberghe (2004), “Convex optimization”, Chapter 4
- O. Guler (2010), “Foundations of optimization”, Chapter 4