

Appendix A

Properties of positive (semi)definite matrices

In this appendix we list some well-known properties of positive (semi)definite matrices which are used in this monograph. The proofs which are omitted here may be found in [85]. A more detailed review of the matrix analysis which is relevant for SDP is given by Jarre in [94].

A.1 CHARACTERIZATIONS OF POSITIVE (SEMI)DEFINITENESS

Theorem A.1 *Let $X \in \mathcal{S}_n$. The following statements are equivalent:*

- $X \in \mathcal{S}_n^+$ or $X \succeq 0$ (X is positive semidefinite);
- $z^T X z \geq 0 \quad \forall z \in \mathbf{R}^n$;
- $\lambda_{\min}(X) \geq 0$;
- All principal minors of X are nonnegative;
- $X = LL^T$ for some $L \in \mathbf{R}^{n \times n}$.

We can replace ‘positive semidefinite’ by ‘positive definite’ in the statement of the theorem by changing the respective nonnegativity requirements to positivity, and by requiring that the matrix L in the last item be nonsingular. If X is positive definite ($X \in \mathcal{S}_n^{++}$ or $X \succ 0$), the matrix L can be chosen to be lower triangular, in which case we call $X = LL^T$ the *Choleski factorization* of X .

NB: In this monograph positive (semi)definite matrices are necessarily symmetric, *i.e.* we will use ‘positive (semi)definite’ instead of ‘symmetric positive (semi)definite’.¹

¹ In the literature a matrix $X \in \mathbf{R}^{n \times n}$ is sometimes called positive (semi)definite if its symmetric part $\frac{1}{2}(X + X^T)$ is positive (semi)definite.

As an immediate consequence of the second item in Theorem A.1 we have that

$$X \in \mathcal{S}_n^+ \Leftrightarrow AXA^T \in \mathcal{S}_n^+$$

for any given, nonsingular $A \in \mathbf{R}^{n \times n}$.

Another implication is that a block diagonal matrix is positive (semi)definite if and only if each of its diagonal blocks is positive (semi)definite.

A.2 SPECTRAL PROPERTIES

The characterization of positive (semi)definiteness in terms of nonnegative eigenvalues follows from the Raleigh–Ritz theorem.

Theorem A.2 (Raleigh–Ritz) *Let $A \in \mathcal{S}_n$. Then*

$$\lambda_{\min}(A) = \min_{z \in \mathbf{R}^n} \{z^T A z \mid \|z\| = 1\}. \quad (\text{A.1})$$

It is well known that a symmetric matrix has an orthonormal set of eigenvectors, which implies the following result.

Theorem A.3 (Spectral decomposition) *Let $A \in \mathcal{S}_n$. Now A can be decomposed as*

$$A = Q^T \Lambda Q = \sum_{i=1}^n \lambda_i(A) q_i q_i^T$$

where Λ is a diagonal matrix with the eigenvalues $\lambda_i(A)$ ($i = 1, \dots, n$) of A on the diagonal, and Q is an orthogonal matrix with a corresponding set of orthonormal eigenvectors q_1, \dots, q_n of A as columns.

Since $\lambda_i(X) \geq 0$ ($i = 1, \dots, n$) if $X \in \mathcal{S}_n^+$, we can define the symmetric square root factorization of $X \in \mathcal{S}_n^+$:

$$X^{\frac{1}{2}} = \sum_{i=1}^n \sqrt{\lambda_i(X)} q_i q_i^T \quad (X \in \mathcal{S}_n^+).$$

Note that $X^{\frac{1}{2}} X^{\frac{1}{2}} = X$; $X^{\frac{1}{2}}$ is the only matrix with this property.

Theorem A.4 *Let $X \in \mathcal{S}_n^{++}$ and $S \in \mathcal{S}_n^{++}$. Then all the eigenvalues of XS are real and positive.*

Proof:

The proof is immediate by noting that

$$XS \sim \left(X^{\frac{1}{2}}\right)^{-1} XS \left(X^{\frac{1}{2}}\right) = X^{\frac{1}{2}} S X^{\frac{1}{2}} \succ 0.$$

□

We will often use the notation $X^{-\frac{1}{2}} := \left(X^{\frac{1}{2}}\right)^{-1}$.

The eigenvalues of a symmetric matrix can be viewed as smooth functions on \mathcal{S}_n in a sense made precise by the following theorem.

Theorem A.5 (Rellich) *Let an interval $(a, b) \subset \mathbf{R}$ be given. If $A : (a, b) \mapsto \mathcal{S}_n$ is a continuously differentiable function, then there exist n continuously differentiable functions $\lambda_i : (a, b) \mapsto \mathbf{R}$ ($i = 1, \dots, n$) such that $\lambda_1(t), \dots, \lambda_n(t)$ give the values of the eigenvalues of $A(t)$ for each $t \in (a, b)$.*

The next lemma shows what happens to the spectrum of a positive semidefinite matrix if a skew symmetric matrix is added to it, in the case where the eigenvalues of the sum of the two matrices remain real numbers.

Lemma A.1 *Let $Q \in \mathcal{S}_n^{++}$, and let $M \in \mathbf{R}^{n \times n}$ be skew-symmetric ($M = -M^T$). One has $\det(Q + M) > 0$. Moreover, if $\lambda_i(Q + M) \in \mathbf{R}$ ($i = 1, \dots, n$), then*

$$0 < \lambda_{\min}(Q) \leq \lambda_{\min}(Q + M) \leq \lambda_{\max}(Q + M) \leq \lambda_{\max}(Q).$$

Proof:

First note that $Q + M$ is nonsingular since for all nonzero $x \in \mathbf{R}^n$:

$$x^T(Q + M)x = x^T Q x > 0,$$

using the skew symmetry of M . We therefore know that

$$\psi(t) := \det[Q + tM] \neq 0 \quad \forall t \in \mathbf{R},$$

since tM remains skew-symmetric. One now has that ψ is a continuous function of t which is nowhere zero and strictly positive for $t = 0$ as $\det(Q) > 0$. This shows $\det(Q + M) > 0$.

To prove the second part of the lemma, assume $\lambda > 0$ is such that $\lambda > \lambda_{\max}(Q)$. It then follows that $Q - \lambda I \prec 0$. By the same argument as above we then have $(Q + M) - \lambda I$ nonsingular, or

$$\det((Q + M) - \lambda I) \neq 0.$$

This implies that λ cannot be an eigenvalue of $Q + M$. Similarly, $Q + M$ cannot have an eigenvalue smaller than $\lambda_{\min}(Q)$. This gives the required result. □

The *spectral norm* $\|\cdot\|_2$ on $\mathbf{R}^{n \times n}$ is the norm induced by the Euclidean vector norm:

$$\|A\|_2 := \max_{x \in \mathbf{R}^n} \frac{\|Ax\|}{\|x\|} \quad (A \in \mathbf{R}^{n \times n}).$$

By the Raleigh–Ritz theorem, the spectral norm and spectral radius $\rho(\cdot)$ coincide for symmetric matrices:

$$\|A\|_2 = \rho(A) := \max_i |\lambda_i(A)| \quad \forall A \in \mathcal{S}_n.$$

The location of the eigenvalues of a matrix is bounded by the famous Gerschgorin theorem. For symmetric matrices the theorem states that

$$\lambda_k(A) \in \bigcup_{i=1}^n \left\{ t \mid |t - a_{ii}| \leq \sum_{j \neq i} |a_{ij}| \right\}, \quad k = 1, \dots, n, \quad A \in \mathcal{S}_n.$$

As a consequence we find that the so-called diagonally dominant matrices are positive semi-definite.

Theorem A.6 (Diagonally dominant matrix is PSD) *A matrix $A \in \mathcal{S}_n$ is called diagonally dominant if*

$$a_{ii} \geq \sum_{j \neq i} |a_{ij}|, \quad i = 1, \dots, n.$$

If A is diagonally dominant, then $A \in \mathcal{S}_n^+$.

A.3 THE TRACE OPERATOR AND THE FROBENIUS NORM

The trace of an $n \times n$ matrix A is defined as

$$\text{Tr}(A) = \sum_{i=1}^n a_{ii}.$$

The trace is clearly a linear operator and has the following properties.

Theorem A.7 *Let $A \in \mathbf{R}^{n \times n}$ and $B \in \mathbf{R}^{n \times n}$. Then the following holds:*

- $\text{Tr}(A) = \sum_{i=1}^n \lambda_i(A)$;
- $\text{Tr}(A) = \text{Tr}(A^T)$;
- $\text{Tr}(AB) = \text{Tr}(BA)$
- $\text{Tr}(AB^T) = \text{vec}(A)^T \text{vec}(B) = \sum_{i,j=1}^n a_{ij} b_{ij}$.

The last item shows that we can view the usual Euclidean inner product on \mathbf{R}^{n^2} as an inner product on $\mathbf{R}^{n \times n}$:

$$\langle A, B \rangle := \text{Tr}(AB^T) = \text{Tr}(B^T A) = \text{Tr}(A^T B) = \text{Tr}(B^T A). \quad (\text{A.2})$$

The inner product in (A.2) induces the so-called *Frobenius* (or Euclidean) norm on $\mathbf{R}^{n \times n}$:

$$\|A\|^2 := \langle A, A \rangle = \text{Tr} (AA^T) = \sum_{i,j=1}^n a_{ij}^2.$$

It now follows from the first item in Theorem (A.7) that

$$\|C\|^2 = \sum_{i=1}^n \lambda_i(C)^2 \text{ if } C \in \mathcal{S}_n.$$

The Frobenius and spectral norms are sub-multiplicative, *i.e.*

$$\|AB\| \leq \|A\| \|B\|, \quad \|AB\|_2 \leq \|A\|_2 \|B\|_2 \quad \forall A \in \mathbf{R}^{n \times n}, B \in \mathbf{R}^{n \times n},$$

and $\|A\|_2 \leq \|A\|$ for all $A \in \mathbf{R}^{n \times n}$.

One can easily prove the useful inequality:

$$\text{Tr} (AB) \leq \lambda_{\max}(A) \text{Tr} (B), \text{ for } A, B \succeq 0, \quad (\text{A.3})$$

that is equivalent to

$$\|AB\| \leq \|A\|_2 \|B\|, \text{ for } A, B \in \mathbf{R}^{n \times n}. \quad (\text{A.4})$$

Inequality (A.4) follows from (A.3) by replacing A by AA^T and B by BB^T in (A.3). Conversely, (A.3) follows from (A.4) by replacing A by $A^{\frac{1}{2}}$ and B by $B^{\frac{1}{2}}$ in (A.4).

Theorem A.8 (Fejer) *A matrix $A \in \mathcal{S}_n$ is positive semidefinite if and only if $\langle A, B \rangle \geq 0$ for all $B \in \mathcal{S}_n^+$. In other words, the cone \mathcal{S}_n^+ is self-dual.*

Proof:

Let $A \in \mathcal{S}_n^+$ and $B \in \mathcal{S}_n^+$; then

$$\langle A, B \rangle = \text{Tr} \left(A^{\frac{1}{2}} A^{\frac{1}{2}} B^{\frac{1}{2}} B^{\frac{1}{2}} \right) = \text{Tr} \left(A^{\frac{1}{2}} B^{\frac{1}{2}} B^{\frac{1}{2}} A^{\frac{1}{2}} \right) = \left\| A^{\frac{1}{2}} B^{\frac{1}{2}} \right\|^2 \geq 0.$$

Conversely, if $A \in \mathcal{S}_n$ and $\langle A, B \rangle \geq 0$ for all $B \in \mathcal{S}_n^+$, then let $x \in \mathbf{R}^n$ be given and set $B = xx^T \in \mathcal{S}_n^+$. Now

$$0 \leq \langle A, B \rangle = \text{Tr} (Axx^T) = \sum_{i,j=1}^n a_{ij} x_i x_j = x^T A x.$$

□

For positive semidefinite matrices, the trace dominates the Frobenius norm, *i.e.*

$$\text{Tr} (X) \geq \|X\| \quad \forall X \in \mathcal{S}_n^+.$$

This follows by applying the inequality

$$\sum_{i=1}^n x_i \geq \sqrt{\sum_{i=1}^n x_i^2} \quad \forall x \in \mathbf{R}_n^+$$

to the nonnegative eigenvalues of X . Similarly, one can apply the arithmetic-geometric mean inequality

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i \quad \forall x \in \mathbf{R}_n^+$$

to the eigenvalues of $X \in \mathcal{S}_n^+$ to obtain the inequality

$$\frac{1}{n} \text{Tr}(X) \geq (\det(X))^{1/n} \quad \forall X \in \mathcal{S}_n^+,$$

where we have used the fact that $\det(A) = \prod_i \lambda_i(A)$ for any $A \in \mathbf{R}^{n \times n}$.

Lemma A.2 *If $X \in \mathcal{S}_n^+$ and $S \in \mathcal{S}_n^+$ and $\text{Tr}(XS) = 0$, then $XS = SX = 0$.*

Proof:

By the properties of the trace operator

$$\text{Tr}(XS) = \text{Tr}\left(X^{\frac{1}{2}} X^{\frac{1}{2}} S^{\frac{1}{2}} S^{\frac{1}{2}}\right) = \text{Tr}\left(S^{\frac{1}{2}} X^{\frac{1}{2}} X^{\frac{1}{2}} S^{\frac{1}{2}}\right) = \left\| S^{\frac{1}{2}} X^{\frac{1}{2}} \right\|^2.$$

Thus if $\text{Tr}(XS) = 0$, it follows that $S^{\frac{1}{2}} X^{\frac{1}{2}} = 0$. Pre-multiplying by $S^{\frac{1}{2}}$ and post-multiplying by $X^{\frac{1}{2}}$ yields $SX = 0$, which in turn implies $(SX)^T = XS = 0$. \square

The following lemma is used to prove that the search directions for the interior point methods described in this monograph are well defined. The proof given here is based on a proof given by Faybusovich [54].

Lemma A.3 *Let $A_i \in \mathcal{S}_n$ ($i = 1, \dots, m$) be linearly independent, and let $0 \neq Y \in \mathcal{S}_n^+$, $Z \in \mathcal{S}_n^{++}$. The matrix $M \in \mathcal{S}_m$ with entries*

$$m_{ij} := \text{Tr}(A_i Z A_j Y), \quad i, j = 1, \dots, m$$

is positive definite.

Proof:

We prove that the quadratic form

$$q(x) = x^T M x = \sum_{i,j=1}^m m_{ij} x_i x_j$$

is strictly positive for all nonzero $x \in \mathbf{R}^n$. To this end, note that for given $x \neq 0$,

$$q(x) = \text{Tr} \left(\left(\sum_{i=1}^m x_i A_i \right) Z \left(\sum_{j=1}^m x_j A_j \right) Y \right).$$

Denoting $A(x) := \sum_{i=1}^m x_i A_i$ (which is nonzero by the linear independence of the A'_i 's), one has:

$$q(x) = \text{Tr} (A(x) Z A(x) Y) > 0,$$

where the inequality follows from $0 \neq A(x) Z A(x) \succeq 0$ and $Y \succ 0$. \square

A.4 THE LÖWNER PARTIAL ORDER AND THE SCHUR COMPLEMENT THEOREM

We define a partial ordering on \mathcal{S}_n via:

$$A \succeq B \iff A - B \in \mathcal{S}_n^+.$$

This partial ordering is called the *Löwner partial order* on \mathcal{S}^n . (It motivates the alternative notation $X \succeq 0$ instead of $X \in \mathcal{S}_n^+$.) It follows immediately that

$$A \succeq B \iff C^T A C \succeq C^T B C \quad \forall C \in \mathbf{R}^{n \times n}.$$

One also has

$$A \succeq B \iff B^{-1} \succeq A^{-1} \quad (A \in \mathcal{S}_n^{++}, B \in \mathcal{S}_n^{++}).$$

The Schur complement theorem gives us useful ways to express positive semidefiniteness of matrices with a block structure.

Theorem A.9 (Schur complement) *If*

$$M = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}$$

where A is positive definite and C is symmetric, then the matrix

$$C - B^T A^{-1} B$$

is called the *Schur complement* of A in X . The following are equivalent:

- M is positive (semi)definite;
- $C - B^T A^{-1} B$ is positive (semi)definite.

Proof:

The result follows by setting $D = -A^{-1}B$, and noting that

$$\begin{bmatrix} I & 0 \\ D^T & I \end{bmatrix} \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \begin{bmatrix} I & D \\ 0 & I \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & C - B^T A^{-1} B \end{bmatrix}.$$

Since a block diagonal matrix is positive (semi)definite if and only if its diagonal blocks are positive (semi)definite, the proof is complete. \square

Appendix B

Background material on convex optimization

In this Appendix we give some background material on convex analysis, convex optimization and nonlinear programming. All proofs are omitted here but may be found in the books by Rockafellar [160] (convex analysis) and Bazaraa *et al.* [16] (nonlinear programming).

B.1 CONVEX ANALYSIS

Definition B.1 (Convex set) Let two points $x^1, x^2 \in \mathbf{R}^n$ and $0 \leq \lambda \leq 1$ be given. Then the point

$$x = \lambda x^1 + (1 - \lambda)x^2$$

is a convex combination of the two points x^1, x^2 .

The set $\mathcal{C} \subset \mathbf{R}^n$ is called convex, if all convex combinations of any two points $x^1, x^2 \in \mathcal{C}$ are again in \mathcal{C} .

Definition B.2 (Convex function) A function $f : \mathcal{C} \rightarrow \mathbf{R}$ defined on a convex set \mathcal{C} is called convex if for all $x^1, x^2 \in \mathcal{C}$ and $0 \leq \lambda \leq 1$ one has

$$f(\lambda x^1 + (1 - \lambda)x^2) \leq \lambda f(x^1) + (1 - \lambda)f(x^2).$$

The function is called strictly convex if the last inequality is strict.

A function is convex if and only if its epigraph is convex.

Definition B.3 (Epigraph) The epigraph of a function $f : \mathcal{C} \rightarrow \mathbf{R}$ is the $(n + 1)$ -dimensional set

$$\{(x, \tau) : f(x) \leq \tau, x \in \mathcal{C}, \tau \in \mathbf{R}\}.$$

Theorem B.1 A twice differentiable function f is convex (resp. strictly convex) on an open set \mathcal{C} if and only if its Hessian $\nabla^2 f$ is positive semidefinite (resp. positive definite) on \mathcal{C} .

Example B.1 The function $f : \mathcal{S}_n^{++} \mapsto \mathbf{R}$ defined by $f(X) = -\log(\det(X))$ has a positive definite Hessian and is therefore strictly convex. (This is proven in Appendix C.) \square

Strictly convex functions are useful for proving ‘uniqueness properties’, due to the following result.

Theorem B.2 If a strictly convex function has a minimizer over a convex set, then this minimizer is unique.

Definition B.4 (Convex cone) The set $\mathcal{K} \subset \mathbf{R}^n$ is a convex cone if it is a convex set and for all $x \in \mathcal{K}$ and $0 < \lambda$ one has $\lambda x \in \mathcal{K}$.

Example B.2 Four examples of convex cones in \mathcal{S}_n are:

- the symmetric positive semidefinite cone:

$$\mathcal{S}_n^+ := \{A \in \mathcal{S}_n \mid x^T A x \geq 0 \text{ for all } x \in \mathbf{R}^n\};$$

- the copositive cone:

$$\mathcal{C}_n := \{A \in \mathcal{S}_n \mid x^T A x \geq 0 \text{ for all } x \in \mathbf{R}_+^n\};$$

- the cone of completely positive matrices:

$$\mathcal{C}_n^* = \left\{ A \in \mathcal{S}_n \mid A = \sum_{i=1}^k x_i x_i^T, x_i \in \mathbf{R}_+^n \text{ } (i = 1, \dots, k) \text{ for any } k \right\};$$

- the cone of nonnegative matrices:

$$\mathcal{N}_n = \{A \in \mathcal{S}_n \mid a_{ij} \geq 0 \text{ } (i, j = 1, \dots, n)\}.$$

\square

Definition B.5 (Face (of a cone)) A subset \mathcal{F} of a convex cone \mathcal{K} is called a face of \mathcal{K} if for all $x \in \mathcal{F}$ and $y, z \in \mathcal{K}$ one has $x = y + z$ if and only if $y \in \mathcal{F}$ and $z \in \mathcal{F}$.

Example B.3 An example of a face of the cone of positive semidefinite matrices \mathcal{S}_n^+ is

$$\mathcal{F} := \left\{ \begin{bmatrix} U & 0_{r \times (n-r)} \\ 0_{(n-r) \times r} & 0_{(n-r) \times (n-r)} \end{bmatrix} \mid U \in \mathcal{S}_r^+ \right\}.$$

Note that if $A \in \mathcal{S}_n^+$ and $B \in \mathcal{S}_n^+$, then $A + B \in \mathcal{F}$ if and only if $A \in \mathcal{F}$ and $B \in \mathcal{F}$. \square

Definition B.6 (Extreme ray (of a cone)) A subset of a convex cone \mathcal{K} is called an extreme ray of \mathcal{K} if it is a one dimensional face of \mathcal{K} , i.e. a face that is a half line emanating from the origin.

Example B.4 Any $0 \neq x \in \mathbf{R}^n$ defines an extreme ray of the cone of positive semidefinite matrices \mathcal{S}_n^+ via

$$\{U \mid U = cxx^T, c > 0\}.$$

Similarly, any $y \in \mathbf{R}_+^n$ defines an extreme ray of the cone of completely positive semidefinite matrices \mathcal{C}_n^* via

$$\{U \mid U = cyy^T, c > 0\}.$$

\square

Definition B.7 Let a convex set \mathcal{C} be given. The point $x \in \mathcal{C}$ is in the relative interior of \mathcal{C} if for all $\hat{x} \in \mathcal{C}$ there exists $\tilde{x} \in \mathcal{C}$ and $0 < \lambda < 1$ such that $x = \lambda \hat{x} + (1 - \lambda)\tilde{x}$. The set of relative interior points of the set \mathcal{C} will be denoted by $\text{ri}(\mathcal{C})$.

Theorem B.3 Assume that \mathcal{C}_1 and \mathcal{C}_2 are nonempty convex sets and $\text{ri}(\mathcal{C}_1) \cap \text{ri}(\mathcal{C}_2) \neq \emptyset$. Then

$$\text{ri}(\mathcal{C}_1 \cap \mathcal{C}_2) = \text{ri}(\mathcal{C}_1) \cap \text{ri}(\mathcal{C}_2).$$

Example B.5 Note that the interior of the cone of positive semidefinite matrices \mathcal{S}_n^+ is the cone of positive definite matrices \mathcal{S}_n^{++} . Let $\mathcal{A} \subset \mathcal{S}_n$ denote an affine space. Assume that there exists an $X \in \mathcal{A}$ that is also positive definite. Then, by the last theorem,

$$\text{ri}(\mathcal{A} \cap \mathcal{S}_n^+) = \text{ri}(\mathcal{A}) \cap \text{ri}(\mathcal{S}_n^+) = \mathcal{A} \cap \mathcal{S}_n^{++},$$

since $\text{ri}(\mathcal{A}) = \mathcal{A}$. \square

Theorem B.4 (Separation theorem for convex sets) *Let \mathcal{C}_1 and \mathcal{C}_2 be nonempty convex sets in \mathbf{R}^k . There exists a $r \in \mathbf{R}^k$ such that*

$$\sup_{x \in \mathcal{C}_1} r^T x \leq \inf_{y \in \mathcal{C}_2} r^T y$$

and

$$\inf_{x \in \mathcal{C}_1} r^T x < \sup_{y \in \mathcal{C}_2} r^T y$$

if and only if the relative interiors of \mathcal{C}_1 and \mathcal{C}_2 are disjoint.

The second inequality merely excludes the uninteresting case where the separating hyperplane contains both \mathcal{C}_1 and \mathcal{C}_2 , i.e. it ensures so-called *proper separation*.

B.2 DUALITY IN CONVEX OPTIMIZATION

We consider the generic convex optimization problem:

$$\begin{aligned} (CO) \quad & p^* = \inf_x f(x) \\ \text{s.t.} \quad & g_j(x) \leq 0, \quad j = 1, \dots, m \\ & x \in \mathcal{C}, \end{aligned} \tag{B.1}$$

where $\mathcal{C} \subseteq \mathbf{R}^n$ is a convex set and f, g_1, \dots, g_m are differentiable convex functions on \mathcal{C} (or on an open set that contains the set \mathcal{C}).

For the convex optimization problem (CO) one defines the Lagrange function (or Lagrangian)

$$L(x, y) := f(x) + \sum_{j=1}^m y_j g_j(x) \tag{B.2}$$

where $x \in \mathcal{C}$ and $y \geq 0$.

Definition B.8 *A vector pair $(\bar{x}, \bar{y}) \in \mathbf{R}^{n+m}$, $\bar{x} \in \mathcal{C}$ and $\bar{y} \geq 0$ is called a saddle point of the Lagrange function L if*

$$L(\bar{x}, y) \leq L(\bar{x}, \bar{y}) \leq L(x, \bar{y}) \tag{B.3}$$

for all $x \in \mathcal{C}$ and $y \geq 0$.

One easily sees that (B.3) is equivalent with

$$L(\bar{x}, y) \leq L(x, \bar{y}) \quad \text{for all } x \in \mathcal{C}, \quad y \geq 0.$$

Lemma B.1 *The vector $(\bar{x}, \bar{y}) \in \mathbf{R}^{n+m}$, $\bar{x} \in \mathcal{C}$ and $\bar{y} \geq 0$ is a saddle point of $L(x, y)$ if and only if*

$$\inf_{x \in \mathcal{C}} \sup_{y \geq 0} L(x, y) = L(\bar{x}, \bar{y}) = \sup_{y \geq 0} \inf_{x \in \mathcal{C}} L(x, y). \tag{B.4}$$

Since we can reformulate (CO) as

$$p^* = \inf_{x \in \mathcal{C}} \sup_{y \geq 0} \left\{ f(x) + \sum_{j=1}^m y_j g_j(x) \right\},$$

it follows that \bar{x} is an optimal solution of problem (CO) if there exists a $\bar{y} \geq 0$ such that (\bar{x}, \bar{y}) is a saddle point of the Lagrangian.

To ensure the existence of a saddle point of the Lagrangian, it is sufficient to require the so-called Slater regularity condition (Slater constraint qualification).

Assumption B.1 (Slater regularity) *There exists an $x \in \text{ri}(\mathcal{C})$ such that*

- $g_j(x) < 0$ if g_j not linear or affine;
- $g_j(x) \leq 0$ if g_j is linear or affine.

Under the Slater regularity assumption, we therefore have a one-to-one correspondence between a saddle point of the Lagrangian and an optimal solution of (CO).

Theorem B.5 (Karush–Kuhn–Tucker) *The convex optimization problem (CO) is given. Assume that the Slater regularity condition is satisfied. The vector \bar{x} is an optimal solution of (CO) if and only if there is a vector \bar{y} such that (\bar{x}, \bar{y}) is a saddle point of the Lagrange function L .*

The formulation of the saddle point condition in Lemma (B.1) motivates the concept of the Lagrangian dual.

Definition B.9 (Lagrangian dual) *Denote*

$$\psi(y) = \inf_{x \in \mathcal{C}} \left\{ f(x) + \sum_{j=1}^m y_j g_j(x) \right\}.$$

The problem

$$\begin{aligned} & \psi(y) \\ & y \geq 0 \end{aligned}$$

is called the Lagrangian dual of the convex optimization problem (CO).

It is straightforward to show the so-called weak duality property.

Theorem B.6 (Weak duality) *If \bar{x} is a feasible solution of (CO) and $\bar{y} \geq 0$, then*

$$\psi(\bar{y}) \leq f(\bar{x})$$

and equality holds if and only if $\inf_{x \in \mathcal{C}} \{f(x) + \sum_{j=1}^m \bar{y}_j g_j(x)\} = f(\bar{x})$.

Under the Slater regularity assumption we have a stronger duality result, by the Karush–Kuhn–Tucker theorem and Lemma (B.1).

Theorem B.7 (Strong duality) Assume that (CO) satisfies the Slater regularity condition, and let \bar{x} be a feasible solution of (CO). Now the vector \bar{x} is an optimal solution of (CO) if and only if there exists a $\bar{y} \geq 0$ such that \bar{y} is an optimal solution of the Lagrangian dual problem and

$$\psi(\bar{y}) = f(\bar{x}).$$

B.3 THE KKT OPTIMALITY CONDITIONS

We now state the Karush–Kuhn–Tucker (KKT) necessary and sufficient optimality conditions for problem (CO). First we define the notion of a KKT point.

Definition B.10 (KKT point) The vector $(\hat{x}, \bar{y}) \in \mathcal{C} \times \mathbf{R}^m$ is called a Karush–Kuhn–Tucker (KKT) point of (CO) if

- (i) $g_j(\hat{x}) \leq 0$, for all $j = 1, \dots, m$
- (ii) $0 = \nabla f(\hat{x}) + \sum_{j=1}^m \bar{y}_j \nabla g_j(\hat{x})$
- (iii) $\sum_{j=1}^m \bar{y}_j g_j(\hat{x}) = 0$
- (iv) $\bar{y} \geq 0$.

A KKT point is a saddle point of the Lagrangian L of (CO). Conversely, a saddle point of L , is a KKT point of (CO). This leads us to the following result.

Theorem B.8 (KKT conditions) If (\bar{x}, \bar{y}) is a KKT point, then \bar{x} is an optimal solution of (CO). Conversely — under the Slater regularity assumption — a feasible solution \bar{x} of (CO) is optimal if there exists a $\bar{y} \in \mathbf{R}^m$ such that (\bar{x}, \bar{y}) is a KKT point.

We say that $x \in \mathcal{C}$ meets the KKT conditions if there exists a $y \geq 0$ such that (x, y) is a KKT point of (CO).

If we drop the convexity requirements on f and g_i in the statement of (CO), then the KKT conditions remain *necessary* optimality conditions under the Slater regularity assumption.

Appendix C

The function $\log \det(X)$

In this appendix we develop the matrix calculus needed to derive the gradient and Hessian of the function $\log \det(X)$, and show that it is a strictly concave function.

Lemma C.1 *Let $f : \text{int}(\mathcal{S}_n^+) \mapsto \mathbf{R}$ be given by*

$$f(X) = \log \det X,$$

Denoting

$$\nabla f(X) := \begin{bmatrix} \frac{\partial f(X)}{\partial x_{11}} & \cdots & \frac{\partial f(X)}{\partial x_{1n}} \\ \vdots & \ddots & \vdots \\ \frac{\partial f(X)}{\partial x_{n1}} & \cdots & \frac{\partial f(X)}{\partial x_{nn}} \end{bmatrix},$$

one has $\nabla f(X) = X^{-1}$.

Proof:

Let $X \in \text{int}(\mathcal{S}_n^+)$ be given and let $H \in \mathcal{S}_n$ be such that $X + H \in \text{int}(\mathcal{S}_n^+)$. One has

$$\begin{aligned} f(X + H) - f(X) &= \log \det(X + H) - \log \det(X) \\ &= \log \det(X^{-1}(X + H)) \\ &= \log \det\left(I + X^{-\frac{1}{2}}HX^{-\frac{1}{2}}\right). \end{aligned}$$

By the arithmetic-geometric inequality applied to the eigenvalues of $X^{-\frac{1}{2}}HX^{-\frac{1}{2}}$ one has

$$\begin{aligned} \log \det\left(I + X^{-\frac{1}{2}}HX^{-\frac{1}{2}}\right) &\leq \log\left(\frac{1}{n}\text{Tr}\left(I + X^{-\frac{1}{2}}HX^{-\frac{1}{2}}\right)\right)^n \\ &= n \log\left(\frac{1}{n}\text{Tr}\left(I + X^{-\frac{1}{2}}HX^{-\frac{1}{2}}\right)\right) \end{aligned}$$

$$= n \log \left(1 + \frac{1}{n} \text{Tr} \left(X^{-\frac{1}{2}} H X^{-\frac{1}{2}} \right) \right).$$

Using the well-known inequality $\log(1+t) \leq t$ we arrive at

$$f(X+H) - f(X) \leq \text{Tr} \left(X^{-\frac{1}{2}} H X^{-\frac{1}{2}} \right) = \langle X^{-1}, H \rangle.$$

This shows that X^{-1} is a subgradient of f at X . Since f is assumed differentiable, the subgradient is unique and equals the gradient $\nabla f(X)$. \square

The proof of the next result is trivial.

Lemma C.2 *Let $f : \text{int}(\mathcal{S}_n^+) \mapsto \mathbf{R}$ be given by*

$$f(X) = \text{Tr}(CX),$$

where $C \in \mathcal{S}_n$. One has $\nabla f(X) = C$.

The following result is used to derive the Hessian of the log-barrier function $f_{\text{bar}}(X) = -\log \det(X)$.

Lemma C.3 *Let $f : \mathcal{S}_n^{++} \mapsto \mathbf{R}$ be given by*

$$f(X) = \log \det X.$$

If $\nabla^2 f$ denotes the derivative of $\nabla f : X \mapsto X^{-1}$ with respect to X , then $\nabla^2 f(X)$ is the linear operator which satisfies

$$\nabla^2 f(X)H = -X^{-1}HX^{-1}, \quad \forall H \in \mathcal{S}_n,$$

for a given invertible X .

Proof:

Let $\mathbf{L}(\mathcal{S}_n, \mathcal{S}_n)$ denote the space of linear operators which map \mathcal{S}_n to \mathcal{S}_n . The Frechet derivative of ∇f is defined as the (unique) function $\nabla^2 f : \mathcal{S}_n \mapsto \mathbf{L}(\mathcal{S}_n, \mathcal{S}_n)$ such that

$$\lim_{\|H\| \rightarrow 0} \frac{\|\nabla f(X+H) - \nabla f(X) - \nabla^2 f(X)H\|}{\|H\|} = 0. \quad (\text{C.1})$$

We show that $\nabla^2 f(X)H := -X^{-1}HX^{-1}$ satisfies (C.1). To this end, let $H \in \mathcal{S}_n$ be such that $(X+H)$ is invertible, and consider

$$\begin{aligned} & \|\nabla f(X+H) - \nabla f(X) - \nabla^2 f(X)H\| \\ &= \|(X+H)^{-1} - X^{-1} + X^{-1}HX^{-1}\| \\ &= \|(X+H)^{-1}(I - (X+H)X^{-1} + (X+H)X^{-1}HX^{-1})\| \\ &= \|(X+H)^{-1}(HX^{-1}HX^{-1})\| \\ &\leq \|(X+H)^{-1}\| \|H\| \|X^{-1}HX^{-1}\|, \end{aligned}$$

which shows that (C.1) indeed holds. \square

By Lemma A.3, the Hessian of the function $f(X) = -\log \det(X)$ is a positive definite operator which implies that f is strictly convex on \mathcal{S}_n^{++} . We state this observation as a theorem.

Theorem C.1 *The function $f : \mathcal{S}_n^{++} \mapsto \mathbf{R}$ defined by*

$$f(X) = -\log \det(X)$$

is strictly convex.

An alternative proof of this theorem is given in [85] (Theorem 7.6.6).

Appendix D

Real analytic functions

This appendix gives some elementary properties of analytic functions which are used in this monograph. It is based on notes by M. Halická [76].

Definition D.1 A function $f(x) : \mathbf{R} \rightarrow \mathbf{R}$ is said to be analytic at $x = a$ if there exist $r > 0$ and $\{a_n\}$ such that

$$f(x) = \sum_{n=0}^{\infty} a_n (x - a)^n, \quad \forall x : |x - a| < r. \quad (D.1)$$

Remark D.1 Taking the n th derivative in (D.1), it is easy to see that $a_n = \frac{f^{(n)}(a)}{n!}$. Hence the series in (D.1) is the Taylor series of $f(x)$ at $x = a$.

Remark D.2 A function $f(x)$ that is analytic at $x = a$ is infinitely differentiable in some neighborhood of a and the corresponding Taylor series converges to $f(x)$ in some neighborhood of a (sometimes this is used as the definition of an analytic function). If a function is infinitely differentiable, then it is not necessarily analytic. A well-known example is the Cauchy function $f(x) = e^{-1/x^2}$ for $x \neq 0$ and $f(x) = 0$ for $x = 0$. At $x = 0$ all derivatives are zero and hence the corresponding Taylor series converges to the zero function.

Definition D.2 Let $I \subset \mathbf{R}$ be an open interval. Then f is analytic on I if it is analytic at any point $a \in I$.

We can extend this definition to closed intervals by making the following changes to the above.

- (a) if $f(x)$ is defined for all $x \geq a$, then $f(x)$ is defined to be analytic at a if there exist an $r > 0$ and a series of coefficients $\{a_n\}$ such that the equality in (D.1) holds for all $x \geq a$.

- (b) if $f(x)$ is defined for all $x > a$, then we say that $f(x)$ can be analytically extended to a if there exist an $r > 0$ and a series of coefficients $\{a_n\}$ such that the equality in (D.1) holds for all $x \geq a$.

The following result is used in the proof of Theorem 3.6 (that the central path has a unique limit point in the optimal set).

Theorem D.1 *Let $f : [0, 1) \rightarrow \mathbf{R}$ be analytic on $[0, 1)$. Let $f(0) = 0$ and $x = 0$ be an accumulation point of all x such that $f(x) = 0$. Then $f(x) = 0$ for all $x \in [0, 1)$.*

Proof:

Since f is analytic at $x = 0$, there exist $r > 0$ and a_n such that

$$f(x) = \sum_{n=0}^{\infty} a_n x^n, \quad \forall x : 0 \leq x < r. \quad (\text{D.2})$$

We show that $a_n = 0, \forall n$. Let a_k be the first non-zero coefficient in (D.2). Hence

$$f(x) = x^k(a_k + a_{k+1}x + \dots), \quad \forall x : 0 \leq x < r. \quad (\text{D.3})$$

Let $0 < \rho < r$. The series in (D.3) converges at $x = \rho$ and hence the numbers $a_n \rho^n$ are bounded. Let $|a_n| \rho^n < K$, i.e., $|a_n| < K/\rho^n, \forall n$. Then

$$|f(x)| \geq |x^k| \left(|a_k| - \frac{K|x|}{\rho^{k+1}} - \frac{K|x|^2}{\rho^{k+2}} - \dots \right) = |x^k| \left(|a_k| - \frac{K|x|}{\rho^k(\rho - |x|)} \right), \quad (\text{D.4})$$

for all $0 < x < \rho$. The last expression in (D.4) is positive for all sufficiently small x . This is in contradiction with the assumption that f has roots arbitrarily close to $x = 0$. Hence $a_n = 0, \forall n$ and $f(x) = 0$ in some right neighborhood of 0. Using the analyticity in $(0, 1)$ it can be shown that it is zero on the whole domain. \square

Appendix E

The (symmetric) Kronecker product

This appendix contains various results about the Kronecker and symmetric Kronecker product. The part on the symmetric Kronecker product is based on Appendix D in Aarts[1].

We will use the **vec** and **svec** notation frequently in this appendix, and restate the definitions here for convenience.

Definition E.1 For any symmetric $n \times n$ matrix U , the vector **svec**(U) $\in \mathbf{R}^{\frac{1}{2}n(n+1)}$ is defined as

$$\mathbf{svec}(U) = \left(u_{11}, \sqrt{2}u_{21}, \dots, \sqrt{2}u_{n1}, u_{22}, \sqrt{2}u_{32}, \dots, \sqrt{2}u_{n2}, \dots, u_{nn} \right)^T,$$

such that

$$\mathbf{svec}(U)^T \mathbf{svec}(U) = \text{Tr}(U^T U) = \mathbf{vec}(U)^T \mathbf{vec}(U),$$

where

$$\mathbf{vec}(U) = (u_{11}, u_{21}, \dots, u_{n1}, u_{12}, \dots, u_{nn})^T.$$

The inverse map of **svec** is denoted by **smat**.

E.1 THE KRONECKER PRODUCT

Definition E.2 Let $G \in \mathbf{R}^{m \times n}$ and $K \in \mathbf{R}^{r \times s}$. Then $G \otimes K$ is defined as the $mr \times ns$ matrix with block structure

$$G \otimes K = [g_{ij}K] \quad i = 1, \dots, m, \quad j = 1, \dots, n,$$

i.e. the block in position ij is given by $g_{ij}K$.

The following identities are proven in Horn and Johnson [86]. (We assume that the sizes of the matrices are such that the relations are defined and that the inverses exist where referenced.)

Theorem E.1 *Let K, L, G and H be real matrices.*

- $G \otimes K \text{vec}(H) = \text{vec}(KHG^T)$;
- $(G \otimes K)^T = G^T \otimes K^T$;
- $(G \otimes K)^{-1} = G^{-1} \otimes K^{-1}$;
- $(G \otimes K)(H \otimes L) = (GH) \otimes (KL)$;
- *The eigenvalues of $G \otimes K$ are given by $\lambda_i(G)\lambda_j(K) \forall i, j = 1, \dots, n$. As a consequence, if G and K are positive (semi)definite, then so is $G \otimes K$;*
- *If $Gx_i = \lambda_i(G)x_i$ and $Ky_j = \lambda_j(K)y_j$, then $\text{vec}(y_j x_i^T)$ is an eigenvector of $G \otimes K$ with corresponding eigenvalue $\lambda_i(G)\lambda_j(K)$.*

As an application of the Kronecker product we can analyse the solutions of so-called Lyapunov equations.

Theorem E.2 *Let $A \in \mathbf{R}^{n \times n}$ and $B \in \mathbf{S}_n$. The Lyapunov equation*

$$AX + XA^T = B \tag{E.1}$$

has a unique symmetric solution if A and $-A$ have no eigenvalues in common.

Proof:

Using the first item in Theorem E.1, we can rewrite (E.1) as

$$\text{vec}(AX + XA^T) = \text{vec}(AXI + IXA^T) = (I \otimes A + A \otimes I) \text{vec}(X) = \text{vec}(B).$$

By using the fourth item of Theorem E.1 we have

$$(I \otimes A)(A \otimes I) = (IA) \otimes (AI) = (AI) \otimes (IA) = (A \otimes I)(I \otimes A).$$

In other words, the matrices $A \otimes I$ and $I \otimes A$ commute and therefore share a set of eigenvectors. Moreover, by the fifth item of Theorem E.1 we know that the eigenvalues of $A \otimes I$ and $I \otimes A$ are obtained by taking n copies of the spectrum of A . Therefore each eigenvalue of $(I \otimes A + A \otimes I)$ is given by $\lambda_i(A) + \lambda_j(A)$ for some i, j . It follows that the matrix $(I \otimes A + A \otimes I)$ is nonsingular if A and $-A$ have no eigenvalues in common. In this case equation (E.1) has a unique solution. We must still verify that this solution is symmetric. To this end, note that if X is a solution of (E.1), then so is X^T . This completes the proof. \square

E.2 THE SYMMETRIC KRONECKER PRODUCT

Definition E.3 The symmetric Kronecker product of any two $n \times n$ matrices G and K (not necessarily symmetric), is a mapping on a vector $u = \text{svec}(U)$ where U is a symmetric $n \times n$ matrix and is defined as

$$(G \otimes_s K)u = \frac{1}{2} \text{svec}(KUG^T + GUK^T). \quad (\text{E.2})$$

Note that the linear operator $G \otimes_s K$ is defined implicitly in (E.2). We can give a matrix representation of $G \otimes_s K$ by introducing the orthogonal $\frac{1}{2}n(n+1) \times n$ matrix Q (i.e. $QQ^T = I_{\frac{1}{2}n(n+1)}$), with the property that

$$Q \text{vec}(U) = \text{svec}(U) \text{ and } Q^T \text{svec}(U) = \text{vec}(U) \quad \forall U \in \mathcal{S}_n. \quad (\text{E.3})$$

Theorem E.3 Let Q be the unique orthogonal $\frac{1}{2}n(n+1) \times n$ matrix that satisfies (E.3). For any $G \in \mathbb{R}^{n \times n}$ and $K \in \mathbb{R}^{n \times n}$ one has

$$G \otimes_s K = \frac{1}{2} Q (G \otimes K + K \otimes G) Q^T.$$

Proof:

Let $U \in \mathcal{S}_n$ be given and note that

$$\begin{aligned} \frac{1}{2} Q (G \otimes K + K \otimes G) Q^T \text{svec}(U) &= \frac{1}{2} Q (G \otimes K + K \otimes G) \text{vec}(U) \\ &= \frac{1}{2} Q ((G \otimes K) \text{vec}(U) + (K \otimes G) \text{vec}(U)) \\ &= \frac{1}{2} Q (\text{vec}(KUG^T) + \text{vec}(GUK^T)) \\ &= \frac{1}{2} Q (\text{vec}(KUG^T + (KUG^T)^T)) \\ &= \frac{1}{2} \text{svec}(KUG^T + GUK^T) \\ &= (G \otimes_s K) \text{svec}(U), \end{aligned}$$

where we have used the first identity in Theorem E. 1 to obtain the third equality. \square

Definition E.4 If for every vector $u = \text{svec}(U)$ where U is a symmetric nonzero matrix,

$$u^T (G \otimes_s K) u > 0,$$

then $(G \otimes_s K)$ is called positive definite.

Lemma E.1 *The symmetric Kronecker product has the following properties.*

1. $(G \otimes_s K) = (K \otimes_s G)$;
2. $(G \otimes_s K)(H \otimes_s L) = \frac{1}{2}(GH \otimes_s KL + GL \otimes_s KH)$;
3. $\mathbf{svec}(U)^T \mathbf{svec}(V) = \mathbf{Tr}(UV) = \mathbf{Tr}(VU)$ for two symmetric matrices U and V ;
4. If G and K are symmetric positive definite, then $(G \otimes_s K)$ is positive definite;
5. $(G \otimes_s K)^T = (G^T \otimes_s K^T)$.

Proof:

1. This directly follows from Definition E.3.
2. Let U be a symmetric matrix, then

$$\begin{aligned}
 & (G \otimes_s K)(H \otimes_s L) \mathbf{svec}(U) \\
 &= \frac{1}{2}(G \otimes_s K) \mathbf{svec}(HUL^T + LUH^T) \\
 &= \frac{1}{4} \mathbf{svec}(GHUL^T K^T + KHUL^T G^T + GLUH^T K^T + KLUH^T G^T) \\
 &= \frac{1}{4} \mathbf{svec}(GHU(KL)^T + KLU(GH)^T + KHU(GL)^T + GLU(KH)^T) \\
 &= \frac{1}{2}(GH \otimes_s KL + GL \otimes_s KH) \mathbf{svec}(U).
 \end{aligned}$$

3. See Definition E.1.
4. For every symmetric nonzero matrix U we have to prove that

$$u^T (G \otimes_s K) u > 0,$$

where $u = \mathbf{svec}(U)$. Now

$$u^T (G \otimes_s K) u = \frac{1}{2} u^T \mathbf{svec}(GUK + KUG) = \frac{1}{2} \mathbf{svec}(U)^T \mathbf{svec}(GUK + KUG),$$

since G and K are symmetric. By using Property 3 we derive

$$\frac{1}{2} (\mathbf{svec}(U))^T \mathbf{svec}(GUK + KUG) = \frac{1}{2} (\mathbf{Tr}(UGUK) + \mathbf{Tr}(UKUG)).$$

Since U is nonzero and K and G are symmetric positive definite and therefore nonsingular, we obtain that $K^{\frac{1}{2}} U G^{\frac{1}{2}} \neq 0$ and thus

$$\frac{1}{2} (\mathbf{Tr}(UGUK) + \mathbf{Tr}(UKUG)) = \mathbf{Tr}(UKUG) = \left\| K^{\frac{1}{2}} U G^{\frac{1}{2}} \right\|^2 > 0.$$

5. Define $u = \mathbf{svec}(U)$ and $\ell = \mathbf{svec}(L)$ for arbitrary symmetric matrices U and L and $m = (G \otimes_s K)u$. Now

$$\ell^T m = (\mathbf{svec}(L))^T (G \otimes_s K)u = \frac{1}{2}(\mathbf{svec}(L))^T \mathbf{svec}(GUK^T + KUG^T)$$

and by using Property 3 it follows that

$$\begin{aligned} & \frac{1}{2}(\mathbf{svec}(L))^T \mathbf{svec}(GUK^T + KUG^T) \\ &= \frac{1}{2} \mathbf{Tr}(LGUK^T + LKUG^T) \\ &= \frac{1}{2} \mathbf{Tr}(UK^T LG + UG^T LK) \\ &= \frac{1}{2}(\mathbf{svec}(U))^T \mathbf{svec}(K^T LG + G^T LK) \\ &= (\mathbf{svec}(U))^T (G^T \otimes_s K^T) \mathbf{svec}(L) \\ &= (\mathbf{svec}(U))^T (G^T \otimes_s K^T) \ell. \end{aligned}$$

Since $\ell^T m = m^T \ell$, we obtain

$$m^T = (\mathbf{svec}(U))^T (G^T \otimes_s K^T)$$

and from the definition of m we derive

$$m^T = ((G \otimes_s K) \mathbf{svec}(U))^T = \mathbf{svec}(U)^T (G \otimes_s K)^T$$

and thus

$$(G \otimes_s K)^T = (G^T \otimes_s K^T).$$

□

Definition E.5 For every nonsingular matrix P we define the operators \mathcal{E} and \mathcal{F} as

$$\mathcal{E} = (P \otimes_s P^{-T} S), \quad \mathcal{F} = (PX \otimes_s P^{-T}).$$

By using Property 2 in Lemma E.1 we obtain

$$\mathcal{E} = (P \otimes_s P^{-T} S) = (I \otimes_s P^{-T} S P^{-1})(P \otimes_s P)$$

and

$$\mathcal{F} = (PX \otimes_s P^{-T}) = (PX P^T \otimes_s I)(P^{-T} \otimes_s P^{-T}).$$

We define the operator H_P for any nonsingular $n \times n$ matrix P as

$$H_P(Q) = \frac{1}{2}(PQP^{-1} + P^{-T}Q^T P^T).$$

The next theorems are based on Theorem 3.1 and Theorem 3.2 in Todd *et al.* [173].

Theorem E.4 *If the matrices X and S are positive definite and the matrix $H_P(XS)$ is symmetric positive semidefinite, then $\mathcal{E}^{-1}\mathcal{F}$ is positive definite.*

Proof:

Consider an arbitrary nonzero vector $g \in \mathbf{R}^{\frac{n(n+1)}{2}}$. We now prove that $g^T \mathcal{E}^{-1} \mathcal{F} g > 0$. Defining $k := \mathcal{E}^{-T} g$ and $K = \mathbf{smat}(k)$, where k is also nonzero since \mathcal{E}^{-T} exists, it follows by using Property 5 in Lemma E.1 that

$$g^T \mathcal{E}^{-1} \mathcal{F} g = k^T \mathcal{F} \mathcal{E}^T k = k^T (PX \otimes_s P^{-T}) (P^T \otimes_s SP^{-1}) k.$$

By using Property 2 in Lemma E.1 we derive

$$\begin{aligned} k^T (PX \otimes_s P^{-T}) (P^T \otimes_s SP^{-1}) k &= \frac{1}{2} k^T (PXP^T \otimes_s P^{-T} SP^{-1}) k \\ &\quad + \frac{1}{2} k^T (PXS P^{-1} \otimes_s I) k. \end{aligned}$$

Since X and S are symmetric positive definite, PXP^T and $P^{-T} SP^{-1}$ are symmetric positive definite and by using Property 4 in Lemma E.1 it follows that $(PXP^T \otimes_s P^{-T} SP^{-1})$ is positive definite. Therefore

$$\begin{aligned} &\frac{1}{2} k^T (PXP^T \otimes_s P^{-T} SP^{-1}) k + \frac{1}{2} k^T (PXS P^{-1} \otimes_s I) k \\ &> \frac{1}{2} k^T (PXS P^{-1} \otimes_s I) k \\ &= \frac{1}{4} (\mathbf{svec}(K))^T \mathbf{svec}(PXS P^{-1} K + KP^{-T} SXP^T). \end{aligned}$$

By using Property 3 in Lemma E.1 we now obtain

$$\begin{aligned} &\frac{1}{4} \mathbf{svec}(K)^T \mathbf{svec}(PXS P^{-1} K + KP^{-T} SXP^T) \\ &= \frac{1}{4} \mathbf{Tr} (K (PXS P^{-1} + P^{-T} SXP^T) K) \\ &= \frac{1}{2} \mathbf{Tr} (KH_P(XS)K). \end{aligned}$$

Since we assumed that $H_P(XS)$ is positive semidefinite and the matrix K is symmetric, it follows that the matrix $KH_P(XS)K$ is positive semidefinite. Thus

$$\frac{1}{2} \mathbf{Tr} (KH_P(XS)K) \geq 0.$$

□

Theorem E.5 *If the matrices X and S are symmetric positive definite and the matrices PXP^T and $P^{-T}SP^{-1}$ commute, then $H_P(XS)$ is symmetric positive definite.*

Proof:

If PXP^T and $P^{-T}SP^{-1}$ commute, it follows that

$$PXS P^{-1} = (PXP^T)(P^{-T}SP^{-1}) = (P^{-T}SP^{-1})(PXP^T) = P^{-T}SXP^T.$$

Therefore the matrix $PXS P^{-1}$ is symmetric and

$$H_P(XS) = \frac{1}{2}(PXS P^{-1} + P^{-T}SXP^T) = PXS P^{-1}.$$

From Theorem A.4 we know that XS — and therefore $PXS P^{-1}$ — have positive eigenvalues. Since $H_P(XS)$ is symmetric it is therefore positive definite. \square

Appendix F

Search directions for the embedding problem

Conditions for the existence and uniqueness of several search directions for the self-dual embedding problem of Chapter 4 (Section 4.2) are derived here.

The feasible directions of interior point methods for the embedding problem (4.4) can be computed from the following generic linear system:

$$\begin{array}{ccccccc}
 & \text{Tr}(A_i \Delta X) & -\Delta \tau b_i & +\Delta \theta \bar{b}_i & & & = 0 \\
 -\sum_{i=1}^m \Delta y_i A_i & & +\Delta \tau C & -\Delta \theta \bar{C} & -\Delta S & & = 0 \\
 b^T \Delta y & -\text{Tr}(C \Delta X) & & +\Delta \theta \alpha & & -\Delta \rho & = 0 \\
 -\bar{b}^T \Delta y & +\text{Tr}(\bar{C} \Delta X) & -\Delta \tau \alpha & & & -\Delta \nu & = 0
 \end{array} \tag{F.1}$$

where $i = 1, \dots, m$, and

$$\left. \begin{array}{lcl}
 H_P(\Delta X S + X \Delta S) & = & \mu I - H_P(XS), \\
 \rho \Delta \tau + \tau \Delta \rho & = & \mu - \tau \rho \\
 \nu \Delta \theta + \theta \Delta \nu & = & \mu - \theta \nu,
 \end{array} \right\} \tag{F.2}$$

where H_P is the linear transformation given by

$$H_P(M) := \frac{1}{2} [P M P^{-1} + P^{-T} M^T P^T],$$

for any matrix M , and where the *scaling matrix* P determines the symmetrization strategy. The best-known choices of P from the literature are listed in Table F. 1. We will now prove (or derive sufficient conditions) for existence and uniqueness of the search directions corresponding to each of the choices of P in Table F. 1. To this end,

P	Reference
$\left[X^{\frac{1}{2}} \left(X^{\frac{1}{2}} S X^{\frac{1}{2}} \right)^{-\frac{1}{2}} X^{\frac{1}{2}} \right]^{\frac{1}{2}}$	Nesterov and Todd [138];
$X^{-\frac{1}{2}}$	Monteiro [124], Kojima <i>et al.</i> [108];
$S^{\frac{1}{2}}$	Monteiro [124], Helmberg <i>et al.</i> [84], Kojima <i>et al.</i> [108];
I	Alizadeh, Haeberley and Overton [6];

Table F.1. Choices for the scaling matrix P .

we will write the equations (F.1) and (F.2) as a single linear system and show that the coefficient matrix of this system is nonsingular.¹

We will use the notation

$$\tilde{A}_i := \begin{bmatrix} A_i & & \\ & -b_i & \\ & & \bar{b}_i \end{bmatrix} \quad (i = 1, \dots, m),$$

and define \tilde{P} by replacing X by \tilde{X} and S by \tilde{S} in Table F.1. We will rewrite (F.2) by using the *symmetric Kronecker product*. For a detailed review of the Kronecker product see Appendix E; we only restate the relevant definitions here for convenience.

- $\mathbf{svec}(X) := [X_{11}, \sqrt{2}X_{12}, \dots, \sqrt{2}X_{1n}, X_{22}, \sqrt{2}X_{23}, \dots, X_{nn}]^T$;
- The *symmetric Kronecker product* $G \otimes_s K$ of $G, K \in \mathbf{R}^{n \times n}$ is implicitly defined via

$$(G \otimes_s K) \mathbf{svec}(H) := \frac{1}{2} \mathbf{svec}(KHG^T + GHK^T) \quad (\forall H \in S^n).$$

Using the symmetric Kronecker notation, we can combine (F.1) and (F.2) as

$$\begin{bmatrix} 0 & \tilde{A} & 0 \\ -\tilde{A}^T & S_{\text{kew}} & I \\ 0 & E & F \end{bmatrix} \begin{bmatrix} \Delta y \\ \mathbf{svec}(\Delta \tilde{X}) \\ \mathbf{svec}(\Delta \tilde{S}) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \mathbf{svec}(\mu I - H_{\tilde{P}}(\tilde{X}\tilde{S})) \end{bmatrix} \quad (\text{F.3})$$

¹The approach used here is a straightforward extension of the analysis by Todd *et al.* in [173], where this result was proved for SDP problems in the standard form (P) and (D).

where

$$\begin{aligned}\tilde{\mathcal{A}} &:= [\text{svec}(\tilde{A}_1) \dots \text{svec}(\tilde{A}_m)]^T \\ \mathcal{S}_{kew} &:= \begin{bmatrix} 0 & \text{svec}(C) & -\text{svec}(\tilde{C}) \\ -\text{svec}(C)^T & 0 & \alpha \\ \text{svec}(\tilde{C})^T & -\alpha & 0 \end{bmatrix} \\ E &:= \tilde{P} \otimes_s (\tilde{P}^{-T} \tilde{S}), \quad F := (\tilde{P} \tilde{X}) \otimes_s \tilde{P}^{-T}.\end{aligned}$$

By Theorem E.4 in Appendix E we have the following result.

Lemma F.1 (Toh et al. [175]) *Let \tilde{P} be invertible and \tilde{X} and \tilde{S} symmetric positive definite. Then the matrices E and F are invertible. If one also has $H_{\tilde{P}}(\tilde{X} \tilde{S}) \succ 0$, then the symmetric part of $E^{-1}F$ is also positive definite.*

We are now in a position to prove a sufficient condition for uniqueness of the search direction.

Theorem F.1 *The linear system (F.3) has a unique solution if $H_{\tilde{P}}(\tilde{X} \tilde{S}) \succ 0$.*

Proof:

We consider the homogeneous system

$$\begin{bmatrix} 0 & \tilde{\mathcal{A}} & 0 \\ -\tilde{\mathcal{A}}^T & \mathcal{S}_{kew} & I \\ 0 & E & F \end{bmatrix} \begin{bmatrix} \Delta y \\ \text{svec}(\Delta \tilde{X}) \\ \text{svec}(\Delta \tilde{S}) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \quad (\text{F.4})$$

and prove that it has only the zero vector as solution.

From (F.4) we have

$$\text{svec}(\Delta \tilde{S}) = \tilde{\mathcal{A}}^T \Delta y - \mathcal{S}_{kew} \text{svec}(\Delta \tilde{X})$$

and

$$\text{svec}(\Delta \tilde{S}) = -(F^{-1}E) \text{svec}(\Delta \tilde{X}). \quad (\text{F.5})$$

Eliminating $\text{svec}(\Delta \tilde{S})$ from the last two equations gives

$$\tilde{\mathcal{A}}^T \Delta y - \mathcal{S}_{kew} \text{svec}(\Delta \tilde{X}) + (F^{-1}E) \text{svec}(\Delta \tilde{X}) = 0. \quad (\text{F.6})$$

System (F.4) also implies

$$\tilde{\mathcal{A}} \text{svec}(\Delta \tilde{X}) = 0. \quad (\text{F.7})$$

From (F.6) we have

$$\begin{aligned} \mathbf{svec} \left(\Delta \tilde{X} \right)^T \tilde{A}^T \Delta y - \mathbf{svec} \left(\Delta \tilde{X} \right)^T \mathcal{S}_{\text{keew}} \mathbf{svec} \left(\Delta \tilde{X} \right) \\ + \mathbf{svec} \left(\Delta \tilde{X} \right)^T (F^{-1}E) \mathbf{svec} \left(\Delta \tilde{X} \right) = 0. \end{aligned}$$

The first term on the left-hand side is zero, by (F.7), and the second term is zero by the skew-symmetry of $\mathcal{S}_{\text{keew}}$. We therefore have

$$\mathbf{svec} \left(\Delta \tilde{X} \right)^T (F^{-1}E) \mathbf{svec} \left(\Delta \tilde{X} \right) = 0,$$

which shows that $\Delta \tilde{X} = 0$, since EF^{-1} is assumed to be (non-symmetric) positive definite. It follows that $\Delta \tilde{S} = 0$ by (F.5). Furthermore, $\Delta y = 0$ by (F.6), since \tilde{A} has full rank (the matrices A_i ($i = 1, \dots, m$) are linearly independent). \square

All that remains is to analyze the condition

$$H_{\tilde{P}}(\tilde{X}\tilde{S}) \succ 0 \tag{F.8}$$

in the theorem. For the first three choices of \tilde{P} in Table F.1, condition (F.8) always holds (by Theorem E.5 in Appendix E). For $\tilde{P} = I$ (the so-called AHO direction), (F.8) becomes the condition $\tilde{X}\tilde{S} + \tilde{S}\tilde{X} \succ 0$.

An alternative sufficient condition for existence of the AHO direction was derived by Monteiro and Zanjácomo [126], namely

$$\left\| \frac{1}{\mu} \tilde{X}^{\frac{1}{2}} \tilde{S} \tilde{X}^{\frac{1}{2}} - I \right\| \leq \frac{1}{2},$$

where $\mu = \text{Tr} \left(\tilde{X}\tilde{S} \right) / \tilde{n}$.

Appendix G

Regularized duals

In this appendix we review some strong duality results for SDP problems that do not satisfy the Slater condition.

The duals in question are obtained through a procedure called regularization. Although a detailed treatment of regularization is beyond the scope of this monograph, the underlying idea is quite simple:¹ if the problem

$$(D): \quad d^* := \sup_{y, S} \left\{ b^T y \mid \sum_{i=1}^m y_i A_i + S = C, S \succeq 0, y \in \mathbf{R}^m \right\}$$

is feasible, but not strictly feasible, we can obtain a ‘strictly feasible reformulation’ by replacing the semidefinite cone by a suitable lower dimensional face (say $\mathcal{F} \subset \mathcal{S}_n^+$) of it, such that the new problem

$$(\bar{D}): \quad d^* := \sup_{y, S} \left\{ b^T y \mid \sum_{i=1}^m y_i A_i + S = C, S \in \mathcal{F}, y \in \mathbf{R}^m \right\}$$

is strictly feasible in the sense that there exists a pair (y, S) such that $S \in \text{ri}(\mathcal{F})$ and $\sum_{i=1}^m y_i A_i + S = C$. Problem (\bar{D}) will have a perfect dual, by the conic duality theorem (Theorem 2.3). The main point is to find an explicit expression of the dual of the face \mathcal{F} . The resulting dual problem now takes the form:

$$(P') \quad \inf_X \{ \text{Tr}(CX) \mid \text{Tr}(A_i X) = b_i \ (i = 1, \dots, m), X \in \mathcal{F}^* \},$$

where \mathcal{F}^* denotes the dual cone of \mathcal{F} . In the SDP case \mathcal{F}^* can be described by a system of linear matrix inequalities. There is more than one way to do this and one can obtain different dual problems within this framework (see Pataki [144] for details).

¹The idea of regularization was introduced by Borwein and Wolkowicz [30]. For a more recent (and simplified) treatment the reader is referred to the excellent exposition by Pataki [144].

Ramana [153] first obtained a regularized dual for (D) ;² the so-called gap-free (or extended Lagrange–Slater) primal dual (P_{gf}) of (D) takes the form:

$$p_{gf}^* := \inf \text{Tr} (C(U_0 + W_L))$$

subject to

$$\begin{aligned} \text{Tr} (A_k(U_0 + W_L)) &= b_k, \quad k = 1, \dots, m \\ \text{Tr} (C(U_i + W_{i-1})) &= 0, \quad i = 1, \dots, L \\ \text{Tr} (A_k(U_i + W_{i-1})) &= 0, \quad i = 1, \dots, L, \quad k = 1, \dots, m \\ W_0 &= 0 \\ \begin{bmatrix} I & W_i^T \\ W_i & U_i \end{bmatrix} &\succeq 0, \quad i = 1, \dots, L \\ U_0 &\succeq 0, \end{aligned}$$

where the variables are $U_i \succeq 0$ and $W_i \in \mathbf{R}^{n \times n}$, $i = 0, \dots, L$, and

$$L = \min \left\{ n, \frac{1}{2}n(n+1) - m - 1 \right\}.$$

Note that the gap-free primal problem is easily cast in the standard primal form. Moreover, its size is polynomial in the size of (D) . Unlike the Lagrangian dual (P) of (D) , (P_{gf}) has the following desirable features:

- (Weak duality) If $(y, S) \in \mathcal{D}$ and (U_i, W_i) ($i = 0, \dots, m$) is feasible for (P_{gf}) , then

$$b^T y \leq \text{Tr} (C(U_0 + W_m)).$$
- (Dual boundedness) If (D) is feasible, its optimal value is finite if and only if (P_{gf}) is feasible.
- (Zero duality gap) The optimal value p_{gf}^* of (P_{gf}) equals the optimal value of (D) if and only if both (P_{gf}) and (D) are feasible.
- (Attainment) If the optimal value of (D) is finite, then it is attained by (P_{gf}) .

The standard (Lagrangian) dual problem associated with (P_{gf}) is called the *corrected dual* (D_{cor}) . The pair (P_{gf}) and (D_{cor}) are now in perfect duality; see Ramana and Freund[154].

Moreover, a feasible solution to (D) can be extracted from a feasible solution to (D_{cor}) . The only problem is that (D_{cor}) does not necessarily attain its supremum, even if (D) does.

²In fact, Ramana did not derive this dual via regularization initially; it was only shown subsequently that it can be derived in this way in Ramana *et al.* [155].

Example G.1 *It is readily verified that the weakly infeasible problem (D) in Example 2.2 has a weakly infeasible corrected problem (D_{cor}) .*

The possible duality relations are listed in Table G.1. The optimal value of D_{cor} is denoted by d_{cor}^* .

Status of (D)	Status of (P_{gf})	Status of (D_{cor})
$d^* < \infty$	$p_{gf}^* = d^*$	$d_{cor}^* = d^*$
unbounded	infeasible	unbounded
infeasible	unbounded	infeasible

Table G.1. Duality relations for a given problem (D) , its gap-free dual (P_{gf}) and its corrected problem (D_{cor}) .

References

- [1] L.P. Aarts. Primal-dual search directions in semidefinite optimization. Master's thesis, Delft University of Technology, Faculty of Information Technology and Systems, Delft, The Netherlands, 1999.
- [2] D.V. Alekseevskij, E.B. Vinberg, and A.S. Solodovnikov. Geometry of spaces of constant curvature. In E.B. Vinberg, editor, *Geometry II*, volume 29 of *Encyclopedia of Mathematical Sciences*. Springer-Verlag, 1993.
- [3] F. Alizadeh. *Combinatorial optimization with interior point methods and semi-definite matrices*. PhD thesis, University of Minnesota, Minneapolis, USA, 1991.
- [4] F. Alizadeh. Interior point methods in semidefinite programming with applications to combinatorial optimization. *SIAM Journal on Optimization*, 5:13–51, 1995.
- [5] F. Alizadeh, J.-P.A. Haeberley, and M.L. Overton. Complementarity and non-degeneracy in semidefinite programming. *Mathematical Programming, Series B*, 77(2): 129–162, 1997.
- [6] F. Alizadeh, J.-P.A. Haeberley, and M.L. Overton. Primal-dual methods for semidefinite programming: convergence rates, stability and numerical results. *SIAM Journal on Optimization*, 8(3):746–768, 1998.
- [7] F. Alizadeh and S. Schmieta. Symmetric cones, potential reduction methods. In H. Wolkowicz, R. Saigal, and L. Vandenbergh, editors, *Handbook of semidefinite programming*, pages 195–233. Kluwer Academic Publishers, Norwell, MA, 2000.
- [8] E.D. Andersen. Finding all linearly dependent rows in large-scale linear programming. *Optimization Methods and Software*, 6(3):219–227, 1995.
- [9] E.D. Andersen and K.D. Andersen. The MOSEK interior point optimizer for linear programming: an implementation of the homogeneous algorithm. In H. Frenk, K. Roos, T. Terlaky, and S. Zhang, editors, *High performance optimization*, pages 197–232. Kluwer Academic Publishers, 2000.

- [10] E.D. Andersen, J. Gondzio, C. Mészáros, and X. Xu. Implementation of interior-point methods for large scale linear programs. In T. Terlaky, editor, *Interior point methods of mathematical programming*, pages 189–252. Kluwer, Dordrecht, The Netherlands, 1996.
- [11] M.F. Anjos and H. Wolkowicz. A strengthened SDP relaxation via a second lifting for the Max-Cut problem. *Special Issue of Discrete Applied Mathematics devoted to Foundations of Heuristics in Combinatorial Optimization*, to appear.
- [12] K.M. Anstreicher and M. Fampa. A long-step path following algorithm for semidefinite programming problems. In P. M. Pardalos and H. Wolkowicz, editors, *Topics in Semidefinite and Interior-Point Methods*, volume 18 of *Fields Institute Communications Series*, pages 181–196. American Mathematical Society, 1998.
- [13] V. Balakrishnan and F. Wang. Sdp in systems and control theory. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 421–442. Kluwer Academic Publishers, Norwell, MA, 2000.
- [14] G.P. Barker and D. Carlson. Cones of diagonally dominant matrices. *Pacific Journal of Mathematics*, 57:15–32, 1975.
- [15] E.R. Barnes. A variation on Karmarkar’s algorithm for solving linear programming problems. *Mathematical Programming*, 36:174–182, 1986.
- [16] M.S. Bazaraa, H.D. Sherali, and C.M. Shetty. *Nonlinear Programming: Theory and Algorithms*. John Wiley and Sons, New York, 1993.
- [17] R. Beigel and D. Eppstein. 3-Coloring in time $O(1.3446^n)$: a no-MIS algorithm. In *Proc. 36th IEEE Symp. Foundations of Comp. Sci.*, pages 444–453, 1995.
- [18] M. Bellare and P. Rogaway. The complexity of approximating a nonlinear program. *Mathematical Programming*, 69:429–441, 1995.
- [19] R. Bellman and K. Fan. On systems of linear inequalities in hermitian matrix variables. In V.L. Klee, editor, *Convexity*, Vol. 7 Proc. Symposia in Pure Mathematics, pages 1–11. Amer. Math. Soc., Providence, RI, 1963.
- [20] A. Ben-Tal, L. El Ghaoui, and A.S. Nemirovski. Robustness. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 139–162. Kluwer Academic Publishers, Norwell, MA, 2000.
- [21] A. Ben-Tal and A.S. Nemirovski. Structural design. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 443–467. Kluwer Academic Publishers, Norwell, MA, 2000.
- [22] S.J. Benson and Y. Ye. DSDP3: Dual scaling algorithm for general positive semidefinite programming. Working paper, Computational Optimization Lab, Dept. of Management Science, University of Iowa, Iowa City, USA, 2001.
- [23] S.J. Benson, Y. Ye, and X. Zhang. Solving large-scale sparse semidefinite programs for combinatorial optimization. *SIAM Journal on Optimization*, 10:443–461, 2000.

- [24] Lenore Blum, Mike Shub, and Steve Smale. On a theory of computation and complexity over the real numbers: NP-completeness, recursive functions and universal machines. *Bull. Amer. Math. Soc. (N.S.)*, 21(1): 1–46, 1989.
- [25] I.M. Bomze. On standard quadratic optimization problems. *Journal of Global Optimization*, 13:369–387, 1998.
- [26] I.M. Bomze, M. Budinich, P.M. Pardalos, and M. Pelillo. The maximum clique problem. In D.-Z. Du and P.M. Pardalos, editors, *Handbook of Combinatorial Optimization*, volume suppl. Vol. A, pages 1–74. Kluwer, Dordrecht, 1999.
- [27] I.M. Bomze and E. de Klerk. Solving standard quadratic optimization problems via linear, semidefinite and copositive programming. Technical Report TR 2001–03, Institut für Statistik und Decision Support Systems, Universität Wien, 2001.
- [28] I.M. Bomze, M. Dür, E. de Klerk, C. Roos, A. Quist, and T. Terlaky. On copositive programming and standard quadratic optimization problems. *Journal of Global Optimization*, 18:301–320, 2000.
- [29] R. Boppana and M.M. Halldórsson. Approximating maximum independent sets by excluding subgraphs. *Bit*, 32:180–196, 1992.
- [30] J.M. Borwein and H. Wolkowicz. Regularizing the abstract convex program. *J. Math. Anal. Appl.*, 83(2):495–530, 1981.
- [31] S.E. Boyd, L. El Ghaoui, E. Feron, and V. Balakrishnan. *Linear matrix inequalities in system and control theory*. SIAM Studies in Applied Mathematics, Vol. 15. SIAM, Philadelphia, USA, 1994.
- [32] S. Burer and R.D.C. Monteiro. A Projected Gradient Algorithm for Solving the Maxcut SDP Relaxation. *Optimization Methods and Software*, 15:175–200, 2001.
- [33] J.-D. Cho and M. Sarrafzadeh. Fast approximation algorithms on Maxcut, k -Coloring and k -Color ordering for VLSI applications. *IEEE Transactions on Computers*, 47(11):1253–1266, 1998.
- [34] C. Choi and Y. Ye. Solving sparse semidefinite programs using the dual scaling algorithm with an iterative solver. Working paper, Computational Optimization Lab, Dept. of Management Science, University of Iowa, Iowa City, USA, 2000.
- [35] S.A. Cook. The complexity of theorem proving procedures. In *Proceedings of the 3rd annual ACM symposium on the Theory of Computing*, pages 151–158, 1971.
- [36] W. Cook, C.R. Coullard, and G. Turan. On the complexity of cutting plane proofs. *Discrete Applied Mathematics*, 18:25–38, 1987.
- [37] B. Craven and B. Mond. Linear programming with matrix variables. *Linear Algebra Appl.*, 38:73–80, 1981.
- [38] D. Cvetković, M. Cangalović, and V. Kovačević-Vujčić. Semidefinite programming methods for the traveling salesman problem. In *Proceedings of the 7th International IPCO conference*, pages 126–136, 1999.

- [39] E. de Klerk. *Interior Point Methods for Semidefinite Programming*. PhD thesis, Delft University of Technology, Delft, The Netherlands, 1997.
- [40] E. de Klerk and D.V. Pasechnik. On the performance guarantee of MAX-3-CUT approximation algorithms. Manuscript, March, 2001.
- [41] E. de Klerk and D.V. Pasechnik. Approximation of the stability number of a graph via copositive programming. *SIAM Journal on Optimization*, to appear.
- [42] E. de Klerk, D.V. Pasechnik, and J.P. Warners. On approximate graph colouring and MAX- k -CUT algorithms based on the ϑ -function. Submitted to *Journal of Combinatorial Optimization*, 2000.
- [43] E. de Klerk, J. Peng, C. Roos, and T. Terlaky. A scaled Gauss-Newton primal-dual search direction for semidefinite optimization. *SIAM Journal on Optimization*, 11:870–888, 2001.
- [44] E. de Klerk, C. Roos, and T. Terlaky. Initialization in semidefinite programming via a self-dual, skew-symmetric embedding. *OR Letters*, 20:213–221, 1997.
- [45] E. de Klerk, C. Roos, and T. Terlaky. A short survey on semidefinite programming. In W.K. Klein Haneveld, O.J. Vrieze, and L.C.M. Kallenberg, editors, *Ten years LNMB*. Stichting Mathematisch Centrum, Amsterdam, The Netherlands, 1997.
- [46] E. de Klerk, C. Roos, and T. Terlaky. Infeasible-start semidefinite programming algorithms via self-dual embeddings. In P. M. Pardalos and H. Wolkowicz, editors, *Topics in Semidefinite and Interior-Point Methods*, volume 18 of *Fields Institute Communications Series*, pages 215–236. American Mathematical Society, 1998.
- [47] E. de Klerk, C. Roos, and T. Terlaky. On primal-dual path-following algorithms in semidefinite programming. In F. Giannessi, S. Komlósi, and T. Rapcsák, editors, *New trends in mathematical programming*, pages 137–157. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1998.
- [48] E. de Klerk, C. Roos, and T. Terlaky. Polynomial primal-dual affine scaling algorithms in semidefinite programming. *Journal of Combinatorial Optimization*, 2:51–69, 1998.
- [49] E. de Klerk and H. van Maaren. On semidefinite programming relaxations of $(2 + p)$ -SAT. Submitted to *Annals of Mathematics of Artificial Intelligence*, 2000.
- [50] E. de Klerk, H. van Maaren, and J.P. Warners. Relaxations of the satisfiability problem using semidefinite programming. *Journal of automated reasoning*, 24:37–65, 2000.
- [51] D. den Hertog, E. de Klerk, and C. Roos. On convex quadratic approximation. CentER Discussion Paper 2000-47, CentER for Economic Research, Tilburg University, Tilburg, The Netherlands, 2000.
- [52] J. Dieudonné. *Foundations of Modern Analysis*. Academic Press, New York, 1960.

- [53] I.I. Dikin. Iterative solution of problems of linear and quadratic programming. *Doklady Akademii Nauk SSSR*, 174:747–748, 1967. (Translated in: *Soviet Mathematics Doklady*, 8:674–675, 1967).
- [54] L. Faybusovich. On a matrix generalization of affine-scaling vector fields. *SIAM J. Matrix Anal. Appl.*, 16:886–897, 1995.
- [55] L. Faybusovich. Semi-definite programming: a path-following algorithm for a linear-quadratic functional. *SIAM Journal on Optimization*, 6(4): 1007–1024, 1996.
- [56] U. Feige and M. Goemans. Approximating the value of two prover proof systems with applications to MAX 2SAT and MAX DICUT. In *Proc. Third Israel Symposium on Theory of Computing and Systems*, pages 182–189, 1995.
- [57] U. Feige and J. Kilian. Zero knowledge and the chromatic number. *J. Comput. System Sci.*, 57:187–199, 1998.
- [58] A.V. Fiacco and G.P. McCormick. *Nonlinear programming: sequential unconstrained minimization techniques*. John Wiley & Sons, New York, 1968. (Reprint: Volume 4 of *SIAM Classics in Applied Mathematics*, SIAM Publications, Philadelphia, USA, 1990).
- [59] A. Frieze and M. Jerrum. Improved approximation algorithms for MAX k -cut and MAX BISECTION. In *Proc. 4th IPCO conference*, pages 1–13, 1995.
- [60] K. Fujisawa, M. Kojima, and K. Nakata. Exploiting sparsity in primal-dual interior-point methods for semidefinite programming. *Mathematical Programming*, 79:235–253, 1997.
- [61] M.R. Garey and D.S. Johnson. *Computers and intractability: a guide to the theory of NP-completeness*. W.H. Freeman and Company, Publishers, San Francisco, USA, 1979.
- [62] A. Genz. Numerical computation of multivariate normal probabilities. *J. Comp. Graph. Stat.*, 1:141–149, 1992.
- [63] P.E. Gill, W. Murray, M.A. Saunders, J.A. Tomlin, and M.H. Wright. On projected Newton barrier methods for linear programming and an equivalence to Karmarkar’s projective method. *Mathematical Programming*, 36:183–209, 1986.
- [64] M. Goemans and F. Rendl. Combinatorial optimization. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 343–360. Kluwer Academic Publishers, Norwell, MA, 2000.
- [65] M.X. Goemans. Semidefinite programming in combinatorial optimization. *Math. Programming*, 79(1-3, Ser. B):143–161, 1997. Lectures on mathematical programming (ismpr97).
- [66] M.X. Goemans and D.P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the ACM*, 42(6): 1115–1145, 1995.

- [67] M.X. Goemans and D.P. Williamson. Approximation algorithms for MAX 3-CUT and other problems via complex semidefinite programming. In *Proceedings of the 33rd annual symposium on theory of computing (STOC '01)*, pages 443–452. ACM, 2001.
- [68] D. Goldfarb and K. Scheinberg. Interior point trajectories in semidefinite programming. *SIAM Journal on Optimization*, 8(4):871–886, 1998.
- [69] A.J. Goldman and A.W. Tucker. Theory of linear programming. In H.W. Kuhn and A.W. Tucker, editors, *Linear inequalities and related systems, Annals of Mathematical Studies, No. 38*, pages 53–97. Princeton University Press, Princeton, New Jersey, 1956.
- [70] C.C. Gonzaga. Path following methods for linear programming. *SIAM Review*, 34:167–227, 1992.
- [71] L.M. Graña Drummond and Y. Peterzil. The central path in smooth convex semidefinite programs. *Optimization*, to appear.
- [72] M. Grötschel, L. Lovász, and A. Schrijver. *Geometric Algorithms and Combinatorial Optimization*. Springer-Verlag, Berlin, 1988.
- [73] M. Grötschel, L.A. Lovász, and A. Schrijver. *Geometric algorithms and combinatorial optimization*. Springer Verlag, Berlin, 1988.
- [74] J. Gu, P.W. Purdom, J. Franco, and B.W. Wah. Algorithms for the satisfiability (SAT) problem: a survey. In D. Du, J. Gu, and P.M. Pardalos, editors, *Satisfiability problem: Theory and applications*, volume 35 of *DIMACS series in Discrete Mathematics and Computer Science*. American Mathematical Society, 1997.
- [75] A. Haken. The intractability of resolution. *Theoretical Computer Science*, 39:297–308, 1985.
- [76] M. Halická. Private communication, 2001.
- [77] M. Halická. Analyticity of the central path at the boundary point in semidefinite programming. *European Journal of Operations Research*, to appear.
- [78] M. Halická, E. de Klerk, and C. Roos. On the convergence of the central path in semidefinite optimization. *SIAM Journal on Optimization*, to appear.
- [79] E. Halperin and U. Zwick. Approximation algorithms for MAX 4-SAT and rounding procedures for semidefinite programs. In *Proc. 7th IPCO*, pages 202–217, 1999.
- [80] J. Håstad. Some optimal inapproximability results. In *Proc. 29th Ann. ACM Symp. on Theory of Comp.*, pages 1–10, 1997.
- [81] J. Håstad. Clique is hard to approximate within $|V|^{1-\epsilon}$. *Acta Mathematica*, 182:105–142, 1999.
- [82] B. He, E. de Klerk, C. Roos, and T. Terlaky. Method of approximate centers for semi-definite programming. *Optimization Methods and Software*, 7:291–309, 1997.

- [83] C. Helmberg and F. Oustry. Bundle methods and eigenvalue functions. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 307–337. Kluwer Academic Publishers, Norwell, MA, 2000.
- [84] C. Helmberg, F. Rendl, R.J. Vanderbei, and H. Wolkowicz. An interior-point method for semidefinite programming. *SIAM Journal on Optimization*, 6:342–361, 1996.
- [85] R.A. Horn and C.R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985.
- [86] R.A. Horn and C.R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1991.
- [87] W.-Y. Hsiang. On infinitesimal symmetrization and volume formula for spherical or hyperbolic tetrahedrons. *Quart. J. Math. Oxford*, 39(2):463–468, 1988.
- [88] B. Jansen. *Interior Point Techniques in Optimization*. PhD thesis, Delft University of Technology, Delft, The Netherlands, 1995.
- [89] B. Jansen, C. Roos, and T. Terlaky. The theory of linear programming : Skew symmetric self-dual problems and the central path. *Optimization*, 29:225–233, 1994.
- [90] B. Jansen, C. Roos, and T. Terlaky. Interior point methods: a decade after Karmarkar. A survey, with application to the smallest eigenvalue problem. *Statistica Neerlandica*, 50, 1995.
- [91] B. Jansen, C. Roos, and T. Terlaky. A polynomial primal–dual Dikin–type algorithm for linear programming. *Mathematics of Operations Research*, 21:431–353, 1996.
- [92] B. Jansen, C. Roos, and T. Terlaky. A family of polynomial affine scaling algorithms for positive semi-definite linear complementarity problems. *SIAM Journal on Optimization*, 7(1):126–140, 1997.
- [93] B. Jansen, C. Roos, T. Terlaky, and J.-Ph. Vial. Primal–dual algorithms for linear programming based on the logarithmic barrier method. *Journal of Optimization Theory and Applications*, 83:1–26, 1994.
- [94] F. Jarre. Convex analysis on symmetric matrices. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 13–27. Kluwer Academic Publishers, Norwell, MA, 2000.
- [95] Jr. J.E. Dennis and H. Wolkowicz. Sizing and least change secant methods. *SIGNUM*, 10:1291–1314, 1993.
- [96] J. Jiang. A long step primal–dual path following method for semidefinite programming. *OR Letters*, 23(1,2):53–62, 1998.
- [97] V. Kann, S. Khanna, J. Lagergren, and A. Panconesi. On the hardness of approximating Max k-Cut and its dual. *Chicago Journal of Theoretical Computer Science*, 1997(2), June 1997.

- [98] D. Karger, R. Motwani, and M. Sudan. Approximate graph coloring by semidefinite programming. In *35th Symposium on Foundations of Computer Science*, pages 2–13. IEEE Computer Society Press, 1994.
- [99] H. Karloff and U. Zwick. A 7/8-approximation algorithm for MAX 3SAT? In *Proc. 38th FOCS*, pages 406–415, 1997.
- [100] N.K. Karmarkar. A new polynomial-time algorithm for linear programming. *Combinatorica*, 4:373–395, 1984.
- [101] L. Khachiyan. A polynomial time algorithm in linear programming. *Soviet Mathematics Doklady*, 20:191–194, 1979.
- [102] S. Khanna, N. Linial, and S. Safra. On the hardness of approximating the chromatic number. *Combinatorica*, 20:393–415, 2000.
- [103] D.E. Knuth. The sandwich theorem. *The Electronic Journal of Combinatorics*, 1:1–48, 1994.
- [104] M. Kojima, N. Megiddo, T. Noma, and A. Yoshise. *A unified approach to interior point algorithms for linear complementarity problems*, volume 538 of *Lecture Notes in Computer Science*. Springer Verlag, Berlin, Germany, 1991.
- [105] M. Kojima, M. Shida, and S. Shindoh. Local Convergence of Predictor-Corrector Infeasible-Interior-Point Algorithms for SDP's and SDLCP's. *Mathematical Programming*, 80:129–160, 1998.
- [106] M. Kojima, M. Shida, and S. Shindoh. A note on the Nesterov-Todd and the Kojima-Shindoh-Hara search directions in semidefinite programming. *Optimization Methods and Software*, 11-12:47–52, 1999.
- [107] M. Kojima, M. Shidah, and S. Shindoh. A predictor-corrector interior point algorithm for the semidefinite linear complementarity problem using the Alizadeh-Haeberley-Overton search direction. *SIAM Journal on Optimization*, 9(2):444–465, 1999.
- [108] M. Kojima, S. Shindoh, and S. Hara. Interior point methods for the monotone semidefinite linear complementarity problem in symmetric matrices. *SIAM Journal on Optimization*, 7(1):88–125, 1997.
- [109] S. Kruk, M. Muramatsu, F. Rendl, R.J. Vanderbei, and H. Wolkowicz. The Gauss-Newton direction in linear and semidefinite programming. *Optimization Methods and Software*, 15(1): 1–27, 2001.
- [110] J.B. Lasserre. An explicit exact SDP relaxation for nonlinear 0-1 programs. In *Proc. IPCO VIII*, pages 293–303, 2001.
- [111] J.B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM J. Optimization*, 11:796–817, 2001.
- [112] M. Laurent. A comparison of the Sherali-Adams, Lovász-Schrijver and Lasserre relaxations for 0-1 programming. Technical report pna-r0108, CWI, Amsterdam, The Netherlands, 2001.
- [113] M. Laurent. Tighter linear and semidefinite relaxations for max-cut based on the Lovász-Schrijver lift-and-project procedure. *SIAM Journal on Optimization*, to appear.

- [114] A.S. Lewis and M.L. Overton. Eigenvalue optimization. *Acta Numerica*, 5:149–190, 1996.
- [115] L. Lovász. On the Shannon capacity of a graph. *IEEE Trans. on Information Theory*, 25:1–7, 1979.
- [116] L. Lovász and A. Schrijver. Cones of matrices and set-functions and 0–1 optimization. *SIAM Journal on Optimization*, 1(2):166–190, 1991.
- [117] Z.-Q. Luo, J.F. Sturm, and S. Zhang. Superlinear convergence of a Symmetric primal–dual path following algorithm for semidefinite programming. *SIAM Journal on Optimization*, 8(1):59–81, 1998.
- [118] Z.-Q. Luo, J.F. Sturm, and S. Zhang. Conic convex programming and self-dual embedding. *Optimization Methods and Software*, 14(3): 196–218, 2000.
- [119] I.J. Lustig, R.E. Marsten, and D.F. Shanno. Interior point methods : Computational state of the art. *ORSA Journal on Computing*, 6:1–15, 1994.
- [120] S. Mahajan and H. Ramesh. Derandomizing semidefinite programming based approximation algorithms. In *Proc. of the 36th Annual IEEE Symposium on Foundations of Computer Science*, pages 162–169. IEEE, 1995.
- [121] S. Mehrotra. On the implementation of a (primal–dual) interior point method. *SIAM Journal on Optimization*, 2:575–601, 1992.
- [122] J. Milnor. *Singular points of complex hypersurfaces*. Annals of mathematics studies. Princeton University Press, Princeton, New Jersey, 1968.
- [123] S. Mizuno, M.J. Todd, and Y. Ye. On adaptive step primal–dual interior–point algorithms for linear programming. *Mathematics of Operations Research*, 18:964–981, 1993.
- [124] R.D.C. Monteiro. Primal–dual path-following algorithms for semidefinite programming. *SIAM Journal on Optimization*, 7(3):663–678, 1997.
- [125] R.D.C. Monteiro, I. Adler, and M.G.C. Resende. A polynomial–time primal–dual affine scaling algorithm for linear and convex quadratic programming and its power series extension. *Mathematics of Operations Research*, 15:191–214, 1990.
- [126] R.D.C. Monteiro and P.R. Zanjácomo. A note on the existence of the Alizadeh-Haeberley-Overton direction in semidefinite programming. *Mathematical Programming*, 78(3):393–397, 1997.
- [127] R.D.C. Monteiro and P.R. Zanjácomo. Implementation of primal–dual methods for semidefinite programming based on Monteiro and Tsuchiya directions and their variants. *Optimization Methods and Software*, 11/12:91–140, 1999.
- [128] R. Motwani and P. Raghavan. *Randomized Algorithms*. Cambridge University Press, 1995.
- [129] T.S. Motzkin and E.G. Straus. Maxima for graphs and a new proof of a theorem of Turán. *Canadian J. Math.*, 17:533–540, 1965.
- [130] M. Muramatsu. Affine scaling algorithm fails for semidefinite programming. *Mathematical Programming*, 83(3):393–406, 1998.

- [131] M. Muramatsu and R.J. Vanderbei. Primal-dual affine-scaling algorithm fails for semidefinite programming. *Mathematics of Operations Research*, 24:149–175, 1999.
- [132] K.G. Murty and S.N. Kabadi. Some NP-complete problems in quadratic and linear programming. *Mathematical Programming*, 39:117–129, 1987.
- [133] A. Nemirovskii and P. Gahinet. The projective method for solving linear matrix inequalities. *Mathematical Programming, Series B*, 77(2): 163–190, 1997.
- [134] Y.E. Nesterov. Global quadratic optimization on the sets with simplex structure. Discussion paper 9915, CORE, Catholic University of Louvain, Belgium, 1999.
- [135] Yu. Nesterov. Long-step strategies in interior point potential reduction methods. *Mathematical Programming, Series B*, 76(1):47–94, 1997.
- [136] Yu. Nesterov. Quality of semidefinite relaxation for nonconvex quadratic optimization. Discussion paper 9719, CORE, Catholic University of Louvain, Belgium, March 1997.
- [137] Yu. Nesterov and A.S. Nemirovski. *Interior point polynomial algorithms in convex programming*. SIAM Studies in Applied Mathematics, Vol. 13. SIAM, Philadelphia, USA, 1994.
- [138] Yu. Nesterov and M.J. Todd. Self-scaled barriers and interior-point methods for convex programming. *Mathematics of Operations Research*, 22(1): 1–42, 1997.
- [139] Yu. Nesterov, H. Wolkowicz, and Y. Ye. Semidefinite programming relaxations of nonconvex quadratic optimization. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 361–419. Kluwer Academic Publishers, Norwell, MA, 2000.
- [140] J.A. Olkin. Using semi-definite programming for controller design in active noise control. *SIAG/OPT Views and News*, 8:1–5, Fall 1996.
- [141] P.A. Parillo. *Structured Semidefinite Programs and Semi-algebraic Geometry Methods in Robustness and Optimization*. PhD thesis, California Institute of Technology, Pasadena, California, USA, 2000. Available at: <http://www.cds.caltech.edu/~pablo/>.
- [142] D.V. Pasechnik. Bipartite sandwiches: bounding the size of a maximum biclique. ArXiv e-print, 1999. Available online at <http://xxx.lanl.gov/abs/math.CO/9907109>.
- [143] G. Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Mathematics of Operations Research*, 23(2):339–358, 1998.
- [144] G. Pataki. A Simple Derivation of a Facial Reduction Algorithm, and Extended Dual Systems. Technical Report, Columbia University, New York, USA, 2000.
- [145] G. Pataki. The geometry of semidefinite programming. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 29–66. Kluwer Academic Publishers, Norwell, MA, 2000.

- [146] G. Pólya. Über positive Darstellung von Polynomen. *Vierteljschr. Naturforsch. Ges. Zürich*, 73:141–145, 1928. (also Collected Papers, Vol. 2, 309–313, MIT Press, Cambridge, Mass., London 1974).
- [147] L. Porkolab and L. Khachiyan. On the complexity of semidefinite programs. *Journal of Global Optimization*, 10:351–365, 1997.
- [148] F.A. Potra and R. Sheng. A superlinearly convergent primal–dual infeasible–interior–point algorithm for semidefinite programming. *SIAM Journal on Optimization*, 8(4): 1007–1028, 1998.
- [149] F.A. Potra and R. Sheng. On homogeneous interior–point algorithms for semidefinite programming. *Optimization methods and software*, 9:161–184, 1998.
- [150] V. Powers and B. Reznick. A new bound for Pólya’s Theorem with applications to polynomials positive on polyhedra. In *Proceedings of MEGA 2000*, 2000. To appear in *J. Pure and Applied Algebra*.
- [151] W.R. Pulleyblank. The combinatorial optimization top 10 list. Plenary talk at the ISMP2000 conference, Atlanta, August 7-11, 2000.
- [152] A.J. Quist, E. de Klerk, C. Roos, and T. Terlaky. Copositive relaxation for general quadratic programming. *Optimization Methods and Software*, 9:185–209, 1998.
- [153] M. Ramana. An exact duality theory for semidefinite programming and its complexity implications. *Mathematical Programming, Series B*, 77(2): 129–162, 1997.
- [154] M. Ramana and R. Freund. On the ELSD duality theory for SDP. Technical report, Center of Applied Optimization, University of Florida, Gainesville, Florida, USA, 1996.
- [155] M. Ramana, L. Tunçel, and H. Wolkowicz. Strong duality for semidefinite programming. *SIAM Journal on Optimization*, 7(3):641–662, 1997.
- [156] M.V. Ramana and P.M. Pardalos. Semidefinite programming. In T. Terlaky, editor, *Interior point methods of mathematical programming*, pages 369–398. Kluwer, Dordrecht, The Netherlands, 1996.
- [157] J. Renegar. Condition numbers, the barrier method, and the conjugate-gradient method. *SIAM Journal on Optimization*, 6(4): 879–912, 1996.
- [158] J. Renegar. *A mathematical view of interior-point methods in convex optimization*. SIAM, Philadelphia, PA, 2001.
- [159] Bruce Reznick. Some Concrete Aspects of Hilbert’s 17th Problem. Preprint 98-002, Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA, 1998.
- [160] R.T. Rockafellar. *Convex analysis*. Princeton University Press, Princeton, New Jersey, 1970.
- [161] C. Roos, T. Terlaky, and J.-Ph. Vial. *Theory and Algorithms for Linear Optimization: An interior point approach*. John Wiley & Sons, New York, 1997.

- [162] C. Roos and J.-Ph. Vial. A polynomial method of approximate centers for linear programming. *Mathematical Programming*, 54:295–305, 1992.
- [163] Alexander Schrijver. A comparison of the Delsarte and Lovász bounds. *IEEE Trans. Inform. Theory*, 25(4):425–429, 1979.
- [164] H.D. Sherali and W.P. Adams. A hierarchy of relaxations between the continuous and convex hull representations for 0–1 programming problems. *SIAM Journal on Discrete Mathematics*, 3(3):411–430, 1990.
- [165] M. Shida, S. Shindoh, and M. Kojima. Existence of search directions in interior–point algorithms for the SDP and the monotone SDLCP. *SIAM Journal on Optimization*, 8(2):387–396, 1998.
- [166] N.Z. Shor. Quadratic optimization problems. *Soviet Journal of Computer and System Sciences*, 25:1–11, 1987.
- [167] J.F. Sturm. Using SeDuMi 1.02, a MATLAB toolbox for optimization over symmetric cones. *Optimization Methods and Software*, 11-12:625–653, 1999.
- [168] J.F. Sturm. Error bounds for linear matrix inequalities. *SIAM Journal on Optimization*, 10(4): 1228–1248, 2000.
- [169] J.F. Sturm and S. Zhang. On the long step path–following method for semidefinite programming. *OR Letters*, 22:145–150, 1998.
- [170] J.F. Sturm and S. Zhang. Symmetric primal–dual path following algorithms for semidefinite programming. *Applied Numerical Mathematics*, 29:301–316, 1999.
- [171] K. Tanabe. Centered Newton method for mathematical programming. In *System Modelling and Optimization: Proceedings of the 13th IFIP–Conference, Berlin, August/September 1987*, volume 113 of *Lecture Notes in Control and Information Sciences*, pages 197–206. Springer-Verlag, New York, 1988.
- [172] M.J. Todd. A study of search directions in primal–dual interior–point methods for semidefinite programming. *Optimization Methods and Software*, 11:1–46, 1999.
- [173] M.J. Todd, K.C. Toh, and R.H. Tütüncü. On the Nesterov–Todd direction in semidefinite programming. *SIAM Journal on Optimization*, 8(3):769–796, 1998.
- [174] M.J. Todd and Y. Ye. A centered projective algorithm for linear programming. *Mathematics of Operations Research*, 15:508–529, 1990.
- [175] K.C. Toh, M.J. Todd, and R.H. Tütüncü. SDPT3 — a Matlab software package for semidefinite programming. *Optimization Methods and Software*, 11:545–581, 1999.
- [176] L. Tunçel. Potential reduction and primal–dual methods. In H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors, *Handbook of semidefinite programming*, pages 235–266. Kluwer Academic Publishers, Norwell, MA, 2000.
- [177] A. Urquhart. Open problem posed at SAT’98 workshop, May 10–14, Paderborn, Germany, 1998.

- [178] H. van Maaren. Elliptic approximations of propositional formulae. *Discrete Applied Mathematics*, 96-97:223–244, 1999.
- [179] L. Vandenberghe and S. Boyd. SP: Software for semidefinite programming. User's Guide, Stanford University, Stanford, USA, 1994.
- [180] L. Vandenberghe and S. Boyd. A primal–dual potential reduction algorithm for problems involving matrix inequalities. *Math. Programming*, 69:205–236, 1995.
- [181] L. Vandenberghe and S. Boyd. Semidefinite programming. *SIAM Review*, 38:49–95, 1996.
- [182] R.J. Vanderbei, M.S. Meketon, and B.A. Freedman. A modification of Karmarkar's linear programming algorithm. *Algorithmica*, 1:395–407, 1986.
- [183] H. Wolkowicz, R. Saigal, and L. Vandenberghe, editors. *Handbook of semidefinite programming*. Kluwer Academic Publishers, Norwell, MA, 2000.
- [184] S.J. Wright. *Primal–dual interior point methods*. SIAM, Philadelphia, 1997.
- [185] X. Xu, P.-F. Hung, and Y. Ye. A simplified homogeneous and self–dual linear programming algorithm and its implementation. *Annals of OR*, 62:151–171, 1996.
- [186] H. Yang, Y. Ye, and J. Zhang. An approximation algorithm for the two-parallel machines scheduling problem with capacity constraints. Manuscript, Department of Management Sciences, Henry B. Tippie College of Business Administration, University of Iowa, 2000.
- [187] Y. Ye. Toward probabilistic analysis of interior–point algorithms for linear programming. *SIAM Journal on Optimization*, 19:38–52, 1994.
- [188] Y. Ye. *Interior point algorithms*. Discrete Mathematics and Optimization. Wiley-Interscience, New York, 1997.
- [189] Y. Ye. Approximating quadratic programming with bound and quadratic constraints. *Mathematical Programming*, 84:219–226, 1999.
- [190] Y. Ye. A .699 approximation algorithm for max-bisection. *Mathematical Programming*, 90:101–111, 2001.
- [191] Y. Ye, M.J. Todd, and S. Mizuno. An $\mathcal{O}(\sqrt{n}L)$ -iteration homogeneous and self–dual linear programming algorithm. *Mathematics of Operations Research*, 19:53–67, 1994.
- [192] Y. Zhang. On extending some primal-dual interior point algorithms from linear programming to semidefinite programming. *SIAM Journal on Optimization*, 8(2):365–386, 1998.
- [193] Q. Zhao, S.E. Karisch, F. Rendl, and H. Wolkowicz. Semidefinite programming relaxations for the quadratic assignment problem. *Journal of Combinatorial Optimization*, 2(1):71–109, 1998.
- [194] U. Zwick. Outward rotations: a new tool for rounding solutions of semidefinite programming relaxations, with applications to MAX CUT and other problems. In *Proc. 31st STOC*, pages 679–687, 1999.

Index

- (D) , 2, 22
- (P) , 2, 22
- (P_{gf}) , 262
- B , 50
- N , 50
- T , 50
- Φ , 134
- Ψ , 136, 137
- \mathcal{B} , 35
- \mathcal{D} , 22
- \mathcal{D}^* , 22
- \mathcal{N} , 35
- \mathcal{P} , 22
- \mathcal{P}^* , 22
- \mathcal{T} , 35
- $\delta(X, S, \mu)$, 117
- $\delta_d(S, \mu)$, 89
- $\delta_p(X, \mu)$, 78
- ϵ -optimality, 12
- μ -center, 115
- μ -update, 85, 123
 - adaptive, 87
- ψ , 43, 45, 140
- smat**, 249
- svec**, 249
- k-cut**, 169
- active constraint, 36
- adjacency matrix, 161
 - of the pentagon, 161
- affine-scaling, 15
 - primal, 81, 82
- AHO direction, 260
- algebraic set, 56
- analytic center, 51
 - of the duality gap level sets, 52
- analytic curve
 - central path as an, 46
- analytic extension, 248
 - of the central path, 48
- analytic function, 247
- approximate graph colouring, 183
- approximation algorithm, 174
 - for MAX-3-SAT, 223
 - for **MAX-k-CUT**, 175
- approximation guarantee, 174
 - for global optimization
 - implementable, 208
 - in the weak sense, 207
 - for MAX-3-SAT, 228
 - for **MAX-k-CUT**, 182
 - for standard quadratic optimization, 208
 - for the max. stable set problem, 188
- arithmetic-geometric mean inequality, 52, 53, 136, 234
- arithmetic-geometric inequality, 243
- arrow matrix, 4
- barrier parameter, 76
- big-M method, 16, 61
- bit model of computation, 17
- block diagonal matrix, 230
- Boolean quadratic (in)equality, 214
- Cauchy function, 247
- Cauchy-Schwartz inequality, 88, 108, 137
- centering component, 81
- centering parameter, 76
- central path, 13, 42, 52, 102
 - analyticity of, 46
 - convergence rate of, 58
 - derivatives of, 48
 - limit point, 48
 - of the embedding problem, 67
 - quadratic convergence to, 84, 120, 122
 - tangential direction to, 48
- centrality conditions, 41
- centrality function, 78
 - κ , 102

- of Faybusovich, 78
 - of Jiang, 117
- Choleski factorization, 224, 229
- chromatic number, 4, 158, 223
- clause, 7, 212
- clause–variable matrix, 214
- clique, 4, 222
- clique number, 4, 158, 223
- co-clique, 7
- co-clique number, 7
- combinatorial optimization, 3
- complementary graph, 158
- complementary solutions, 33
- condition number
 - of the embedding, 70
- cone
 - of completely positive matrices, 189
 - extreme ray of, 190, 239
 - of copositive matrices, 189
 - of nonnegative matrices, 189
 - of positive semidefinite matrices, 12
 - extreme ray of, 239
 - face of, 239
- conic duality theorem, 28
- conic problem formulation, 11, 24
- conjugate gradient method, 93
- conjunctive normal form (CNF), 212
- convex cone, 238
- convex function, 149, 237
- convex set, 237
- copositive matrix, 187
- curve selection lemma, 56
- cutting plane, 220
- cycle (in a graph), 203
- damped NT step, 118, 124
- de-randomization, 224
- defect edge, 169, 222
- degree of a vertex, 185
- deterministic rounding, 224
- diagonally dominant, 172, 232
- dihedral angle, 226
- Dikin ellipsoid, 82, 100
- Dikin-type primal–dual affine–scaling, 99
- directional derivative
 - of Φ , 137
 - of Ψ , 137
- dual barrier function, 76
- dual cone, 12, 178
- dual degeneracy, 37
- duality gap, 2, 25, 99, 137
 - after full NT step, 119
 - compact level set of, 43
 - upper bound in terms of $\Phi(X, S)$, 137
- ellipsoid method, 11, 17
- elliptic representations of clauses, 215
- epigraph, 151, 237
- extended Lagrange–Slater dual, 262
- extreme point, 190
- extreme ray, 190, 239
- face, 261
 - of a convex cone, 238
 - of the cone S_n^+ , 34
- feasible direction, 24, 98
- feasible set, 22
- feasible step, 24
- Frobenius norm, 2, 44, 233
- full NT step, 118
- gamma function, 183
- gap relaxation, 221
- gap-free primal problem, 262
- generalised eigenvalues, 135
- Goldman–Tucker theorem, 33
- gradient
 - of $\log \det(X)$, 243
- Gram matrix, 178
- heptagon, 167
- Hessian
 - of $\log \det(X)$, 244
- homogeneous embedding, 16, 64
- icosahedron, 201
- idempotent matrix, 177
- ill-posedness, 71
- implicit function theorem, 46, 48
- improving ray, 29, 69
- incidence vector, 158, 221
- inclusion–exclusion, 225
- independence number, 8
- independent set, 7
- infeasibility, 22
- infeasible–start method, 16
- inner iteration, 92, 124
- integer programming, 214
- interior point assumption, 23
- interior point methods, 11
- Jacobian, 47
- K–Z relaxation, 217, 228
- KKT conditions, 242
- KKT point, 228, 242
- Kronecker product, 47
 - properties of, 249
- Löwner partial order, 2, 235
- Lagrangian, 240
- Lagrangian dual, 2, 241
 - of (P) , 22
 - of SDP in conic form, 24
- Laplacian, 6, 93

- legal colouring, 170, 183
- lift-and-project method, 8
- linear programming (LP), 3, 11
- logarithmic barrier methods, 12
- logarithmic Chebyshev approximation, 9
- logical 'OR' operator, 211
- logical variables, 211
- long step method, 15
 - dual algorithm, 92
 - primal-dual algorithm with NT direction, 124
- Lorentz cone, 154
- Lovász ϑ -function, 4, 158, 172, 189
- Lovász sandwich theorem, 158, 223
- Lyapunov equation, 116, 131, 250
- machine scheduling, 8
- matrix
 - completely positive, 189
 - copositive, 10, 189
 - diagonally dominant, 232
 - idempotent, 177
 - nonnegative, 189
 - positive semidefinite, 2
 - skew-symmetric, 105
- MAX-2-SAT, 7
- MAX-3-SAT, 7
 - approximation guarantee for, 228
- MAX- k -CUT, 169
- MAX- k -SAT, 212
- MAX-BISECTION, 7
- MAX-CUT, 5
- maximal complementarity
 - of optimal solutions, 34
 - of the limit point of the central path, 48, 49
- maximum satisfiability problem, 7
- maximum stable set problem, 188
 - reduction from SAT, 212
- multinomial coefficient, 197
- multinomial theorem, 197
- mutilated chess board formula, 219, 222
- negation (of a logical variable), 211
- Nesterov-Todd (NT) direction, 116
- Nesterov-Todd (NT) scaling, 98
- non-defect edge, 5
- NP-complete, 212
- optimal partition, 35
- optimal solutions, 22
 - uniqueness of, 37, 38
- optimal value
 - of SDP problem, 22
- optimality conditions, 13, 42, 242
 - for (P) and (D) , 33
 - for projected Newton direction, 79
- optimization software, 18
 - CPLEX, 63
 - DSDP, 93
 - MOSEK, 63
 - SDTP3, 15, 131
 - SeDuMi, 15, 64, 131, 154
 - SP, 146
 - XPRESSMP, 63
- orthogonal projection, 79
- orthogonality property, 24, 67
- outer iteration, 92, 124
- Pólya's theorem, 196
- path-following method, 13
 - primal, 75
 - primal-dual, 115
- pentagon
 - $\vartheta(\bar{G})$ -number of, 161
 - approximations of the stability number of, 200
 - Shannon capacity of, 167
- perfect duality, 25, 73
- perfect graph, 162
- Petersen graph, 169
 - maximum stable set of, 188
- pigeonhole formula, 218, 222
- plane search, 135, 136
- pointed cone, 11
- positive semidefiniteness
 - characterizations of, 229
- potential reduction method, 16, 133
 - by Nesterov and Todd, 142
 - general framework, 134
- predictor-corrector method, 15, 115, 146
 - Mizuno-Todd-Ye type, 126
- primal barrier function, 42, 76
- primal degeneracy, 36
- primal-dual affine-scaling, 100, 109
- primal-dual barrier function, 13, 76, 117, 124
- primal-dual Dikin ellipsoid, 99
- primal-dual Dikin-type direction, 136
- primal-dual methods, 13
- projected Newton direction, 78, 89
- propositional formula, 212
- quadratic approximation, 149
 - multivariate case, 152
 - univariate case, 150
- quadratic assignment problem, 8
- quadratic least squares, 149
- quadratic programming (QP), 3
- Raleigh-Ritz theorem, 230
- randomized algorithm, 171, 174
 - for MAX- k -CUT, 175
 - for satisfiable instances of MAX-3-SAT, 224
- randomized rounding, 224

- for 2-clauses, 227
 - for 3-clauses, 227
- regularization, 29, 73, 261
- relative interior, 239
 - of the optimal set, 34
- relative volume, 228
- resolution, 220
- robustness, 11
- sandwich theorem, 172
- satisfiability problem (SAT), 7, 211
- scaled primal–dual direction, 98
- scaling matrix, 257
- Schrijver's ϑ' -function, 201
- Schur complement, 235
- Schur complement theorem, 216, 235
- SDP problems
 - complexity of, 17
 - standard form, 22
- SDTP3, 15, 131
- search directions, 67
- second order cone, 3
- second order solution
 - of centering system, 130
- SeDuMi, 15, 64, 131, 154
- self–dual, 65, 66
- self–dual embedding, 16
 - extended formulation, 64
 - search directions for, 257
- self-concordant barrier, 12
- semi-colouring, 184
- semidefinite feasibility problem, 18
 - complexity of, 18
- semidefinite programming, 1
 - applications in approximation theory, 8
 - applications in combinatorial optimization, 4
 - applications in engineering, 10
 - interior point algorithms for, 11
 - review papers on, 18
 - special cases of, 3
- separation theorem for convex sets, 26, 151, 239
- sequential minimization methods, 12
- Shannon capacity, 165
- Shor relaxation, 215
- simple graph, 169
- skew–symmetric matrix, 105
- Slater regularity, 23, 241
- solvability, 22
- SP, 146
- sparsity, 101
- spectral bundle method, 17
- spectral decomposition, 48
 - of optimal solutions, 35
- spectral norm, xvi, 232
- spectral radius, xvi, 143, 232
- spherical simplex, 178
- spherical tetrahedron, 227, 228
- spherical triangle, 226
 - angular excess of, 226
 - area of, 226
- stability number, 187
- stable set, 7, 188, 213
- stable set polytope, 8
- standard quadratic optimization, 204
 - copositive programming formulation of, 205
 - LP approximation of, 205
 - naive approximation of, 205
 - SDP approximation of, 207
- step length, 24
- strict complementarity, 33, 34, 38, 128
- strict feasibility, 23
- strictly convex function, 237
 - characterization of, 238
 - minimizers of, 238
- strong duality, 3, 25, 242
- strong feasibility, 23
- strong infeasibility
 - characterization of, 29
- strong product for graphs, 164
- structural design, 11
- sum of squares, 10
- superlinear convergence, 15, 128
- symmetric Kronecker product, 251
 - properties of, 252
- symmetric square root factorization, 230
- symmetrization strategy, 14
- system and control theory, 10, 133
- Tanabe–Todd–Ye potential function (Φ), 16, 134, 137
- Taylor expansion, 247
- trace, 2, 44, 99, 232
 - properties of, 232
- traveling salesman problem, 8
- triangle inequalities, 215
- triangle-free graph, 203
- truss, 11
- truth assignment, 211, 224
- Turing machine, 17
- unboundedness, 22
- unit hypersphere, 178, 227
- weak duality, 2, 241
- weak infeasibility, 29
 - characterization of, 31
- weakly improving ray, 31, 32
- weight of a cut, 5, 169
- worst-case iteration bound, 12
 - Dikin-type algorithm, 108
 - dual scaling method, 92
 - long step dual log-barrier method, 92

Mizuno-Todd-Ye predictor-corrector algorithm, 128

NT potential reduction method, 146

primal-dual affine-scaling algorithm, 112

primal-dual path-following algorithm with
full NT steps, 124
short step primal log-barrier method, 88

ZPP, 158

Applied Optimization

1. D.-Z. Du and D.F. Hsu (eds.): *Combinatorial Network Theory*. 1996
ISBN 0-7923-3777-8
2. M.J. Panik: *Linear Programming: Mathematics, Theory and Algorithms*. 1996
ISBN 0-7923-3782-4
3. R.B. Kearfott and V. Kreinovich (eds.): *Applications of Interval Computations*. 1996
ISBN 0-7923-3847-2
4. N. Hritonenko and Y. Yatsenko: *Modeling and Optimimization of the Lifetime of Technology*. 1996
ISBN 0-7923-4014-0
5. T. Terlaky (ed.): *Interior Point Methods of Mathematical Programming*. 1996
ISBN 0-7923-4201-1
6. B. Jansen: *Interior Point Techniques in Optimization. Complementarity, Sensitivity and Algorithms*. 1997
ISBN 0-7923-4430-8
7. A. Migdalas, P.M. Pardalos and S. Storøy (eds.): *Parallel Computing in Optimization*. 1997
ISBN 0-7923-4583-5
8. F.A. Lootsma: *Fuzzy Logic for Planning and Decision Making*. 1997
ISBN 0-7923-4681-5
9. J.A. dos Santos Gromicho: *Quasiconvex Optimization and Location Theory*. 1998
ISBN 0-7923-4694-7
10. V. Kreinovich, A. Lakeyev, J. Rohn and P. Kahl: *Computational Complexity and Feasibility of Data Processing and Interval Computations*. 1998
ISBN 0-7923-4865-6
11. J. Gil-Aluja: *The Interactive Management of Human Resources in Uncertainty*. 1998
ISBN 0-7923-4886-9
12. C. Zopounidis and A.I. Dimitras: *Multicriteria Decision Aid Methods for the Prediction of Business Failure*. 1998
ISBN 0-7923-4900-8
13. F. Giannessi, S. Komlósi and T. Rapcsák (eds.): *New Trends in Mathematical Programming*. Homage to Steven Vajda. 1998
ISBN 0-7923-5036-7
14. Ya-xiang Yuan (ed.): *Advances in Nonlinear Programming*. Proceedings of the '96 International Conference on Nonlinear Programming. 1998
ISBN 0-7923-5053-7
15. W.W. Hager and P.M. Pardalos: *Optimal Control. Theory, Algorithms, and Applications*. 1998
ISBN 0-7923-5067-7
16. Gang Yu (ed.): *Industrial Applications of Combinatorial Optimization*. 1998
ISBN 0-7923-5073-1
17. D. Braha and O. Maimon (eds.): *A Mathematical Theory of Design: Foundations, Algorithms and Applications*. 1998
ISBN 0-7923-5079-0

Applied Optimization

18. O. Maimon, E. Khmelnitsky and K. Kogan: *Optimal Flow Control in Manufacturing. Production Planning and Scheduling.* 1998 ISBN 0-7923-5106-1
19. C. Zopounidis and P.M. Pardalos (eds.): *Managing in Uncertainty: Theory and Practice.* 1998 ISBN 0-7923-5110-X
20. A.S. Belenky: *Operations Research in Transportation Systems: Ideas and Schemes of Optimization Methods for Strategic Planning and Operations Management.* 1998 ISBN 0-7923-5157-6
21. J. Gil-Aluja: *Investment in Uncertainty.* 1999 ISBN 0-7923-5296-3
22. M. Fukushima and L. Qi (eds.): *Reformulation: Nonsmooth, Piecewise Smooth, Semismooth and Smoothing Methods.* 1999 ISBN 0-7923-5320-X
23. M. Patriksson: *Nonlinear Programming and Variational Inequality Problems. A Unified Approach.* 1999 ISBN 0-7923-5455-9
24. R. De Leone, A. Murli, P.M. Pardalos and G. Toraldo (eds.): *High Performance Algorithms and Software in Nonlinear Optimization.* 1999 ISBN 0-7923-5483-4
25. A. Schöbel: *Locating Lines and Hyperplanes. Theory and Algorithms.* 1999 ISBN 0-7923-5559-8
26. R.B. Statnikov: *Multicriteria Design. Optimization and Identification.* 1999 ISBN 0-7923-5560-1
27. V. Tsurkov and A. Mironov: *Minimax under Transportation Constrains.* 1999 ISBN 0-7923-5609-8
28. V.I. Ivanov: *Model Development and Optimization.* 1999 ISBN 0-7923-5610-1
29. F.A. Lootsma: *Multi- Criteria Decision Analysis via Ratio and Difference Judgement.* 1999 ISBN 0-7923-5669-1
30. A. Eberhard, R. Hill, D. Ralph and B.M. Glover (eds.): *Progress in Optimization. Contributions from Australasia.* 1999 ISBN 0-7923-5733-7
31. T. Hürlimann: *Mathematical Modeling and Optimization. An Essay for the Design of Computer-Based Modeling Tools.* 1999 ISBN 0-7923-5927-5
32. J. Gil-Aluja: *Elements for a Theory of Decision in Uncertainty.* 1999 ISBN 0-7923-5987-9
33. H. Frenk, K. Roos, T. Terlaky and S. Zhang (eds.): *High Performance Optimization.* 1999 ISBN 0-7923-6013-3
34. N. Hritonenko and Y. Yatsenko: *Mathematical Modeling in Economics, Ecology and the Environment.* 1999 ISBN 0-7923-6015-X
35. J. Virant: *Design Considerations of Time in Fuzzy Systems.* 2000 ISBN 0-7923-6100-8

Applied Optimization

36. G. Di Pillo and F. Giannessi (eds.): *Nonlinear Optimization and Related Topics*. 2000
ISBN 0-7923-6109-1
37. V. Tsurkov: *Hierarchical Optimization and Mathematical Physics*. 2000
ISBN 0-7923-6175-X
38. C. Zopounidis and M. Doumpos: *Intelligent Decision Aiding Systems Based on Multiple Criteria for Financial Engineering*. 2000
ISBN 0-7923-6273-X
39. X. Yang, A.I. Mees, M. Fisher and L. Jennings (eds.): *Progress in Optimization*. Contributions from Australasia. 2000
ISBN 0-7923-6286-1
40. D. Butnariu and A.N. Iusem: *Totally Convex Functions for Fixed Points Computation and Infinite Dimensional Optimization*. 2000
ISBN 0-7923-6287-X
41. J. Mockus: *A Set of Examples of Global and Discrete Optimization*. Applications of Bayesian Heuristic Approach. 2000
ISBN 0-7923-6359-0
42. H. Neunzert and A.H. Siddiqi: *Topics in Industrial Mathematics*. Case Studies and Related Mathematical Methods. 2000
ISBN 0-7923-6417-1
43. K. Kogan and E. Khmelnitsky: *Scheduling: Control-Based Theory and Polynomial-Time Algorithms*. 2000
ISBN 0-7923-6486-4
44. E. Triantaphyllou: *Multi-Criteria Decision Making Methods*. A Comparative Study. 2000
ISBN 0-7923-6607-7
45. S.H. Zanakis, G. Doukidis and C. Zopounidis (eds.): *Decision Making: Recent Developments and Worldwide Applications*. 2000
ISBN 0-7923-6621-2
46. G.E. Stavroulakis: *Inverse and Crack Identification Problems in Engineering Mechanics*. 2000
ISBN 0-7923-6690-5
47. A. Rubinov and B. Glover (eds.): *Optimization and Related Topics*. 2001
ISBN 0-7923-6732-4
48. M. Pursula and J. Niittymäki (eds.): *Mathematical Methods on Optimization in Transportation Systems*. 2000
ISBN 0-7923-6774-X
49. E. Cascetta: *Transportation Systems Engineering: Theory and Methods*. 2001
ISBN 0-7923-6792-8
50. M.C. Ferris, O.L. Mangasarian and J.-S. Pang (eds.): *Complementarity: Applications, Algorithms and Extensions*. 2001
ISBN 0-7923-6816-9
51. V. Tsurkov: *Large-scale Optimization – Problems and Methods*. 2001
ISBN 0-7923-6817-7
52. X. Yang, K.L. Teo and L. Caccetta (eds.): *Optimization Methods and Applications*. 2001
ISBN 0-7923-6866-5
53. S.M. Stefanov: *Separable Programming Theory and Methods*. 2001
ISBN 0-7923-6882-7

Applied Optimization

- 54. S.P. Uryasev and P.M. Pardalos (eds.): *Stochastic Optimization: Algorithms and Applications*. 2001 ISBN 0-7923-6951 -3
- 55. J. Gil-Aluja (ed.): *Handbook of Management under Uncertainty*. 2001 ISBN 0-7923-7025-2
- 56. B.-N. Vo, A. Cantoni and K.L. Teo: *Filter Design with Time Domain Mask Constraints: Theory and Applications*. 2001 ISBN 0-7923-7138-0
- 57. S. Zlobec: *Stable Parametric Programming*. 2001 ISBN 0-7923-7139-9
- 58. M.G. Nicholls, S. Clarke and B. Lehaney (eds.): *Mixed-Mode Modelling: Mixing Methodologies for Organisational Intervention*. 2001 ISBN 0-7923-7151-8
- 59. F. Giannessi, P.M. Pardalos and T. Rapcsák (eds.): *Optimization Theory. Recent Developments from Mátraháza*. 2001 ISBN 1 -4020-0009-X
- 60. K.M. Hangos, R. Lakner and M. Gerzson: *Intelligent Control Systems. An Introduction with Examples*. 2001 ISBN 1-4020-0134-7
- 61. D. Gstach: *Estimating Output-Specific Efficiencies*. 2002 ISBN 1-4020-0483-4
- 62. J. Geunes, P.M. Pardalos and H.E. Romeijn (eds.): *Supply Chain Management: Models, Applications, and Research Directions*. 2002 ISBN 1-4020-0487-7
- 63. M. Gendreau and P. Marcotte (eds.): *Transportation and Network Analysis: Current Trends*. Miscellanea in Honor of Michael Florian. 2002 ISBN 1-4020-0488-5
- 64. M. Patriksson and M. Labbé (eds.): *Transportation Planning*. State of the Art. 2002 ISBN 1-4020-0546-6
- 65. E. de Klerk: *Aspects of Semidefinite Programming*. Interior Point Algorithms and Selected Applications. 2002 ISBN 1-4020-0547-4